

Evaluation Antidiabetic Activity of Various Leaf Extracts of *Pluchea indica* Less

by Tarsisius Dwi Wibawa Budianta

Submission date: 02-Jul-2022 12:22PM (UTC+0700)

Submission ID: 1865671932

File name: Extracts_of_Pluchea_indica_Less_IJPPR,Vol7,Issue3,Article35.pdf (466.61K)

Word count: 3568

Character count: 19625

Evaluation Antidiabetic Activity of Various Leaf Extracts of *Pluchea indica* Less

Paini Sri Widyawati, Tarsisius Dwi Wibawa Budianta, Dian Ivan Gunawan, Ribka Stefanie Wongso

Food Technology Study Programme, Agricultural Technology Faculty, Widya Mandala Catholic University of Surabaya, East Java, Indonesia

Available Online: 23rd May, 2015

ABSTRACT

Pluchea indica Less leaf extracts of various solvents (methanol, water, ethyl acetate) proven have antioxidant ability using DPPH radical scavenging effect, reducing power, total phenolic content and total flavonoid. However anti-diabetic ability and toxicity level of each extract have not even determined. This study was conducted to determine the effect of different polarity of solvent to phytochemical screening that cause reducing of blood glucose and influence toxicity level of laboratory animal. Our results showed that the water extract was the most potential as anti-diabetic agent. Its blood glucose reducing capability of Rattus Norvegicus Rats Wistar strain was 56.37 % that was the higher than glibenclamide (49.59%) and the other extracts (ethyl acetate extract = 19.11% and methanol extract = 24.27%). The water extract was also safe to the body healthy of Mus Musculus Mice (20-30 g), Balb/c strain. The male mice oral administration of water extract at 2.6 mg/20g b.w. increased motoric activity, however female mice at 1.3 and 2.6 mg/20 g b.w., respectively oral administration experienced motoric activity increasing and reestablishment capacity decreasing. The potency of *Pluchea* leaf water extract as anti-diabetic agent was predicted that there was cardiac glycoside compounds contribution to reduce blood glucose.

Keywords: *Pluchea indica* Less, Anti-diabetic agent, Various Solvent

INTRODUCTION

Type 2 diabetes mellitus (DM) is a disease of carbohydrate metabolism characterized by a number of metabolic abnormalities, including impaired pancreatic beta cell functions and insulin resistance in the skeletal muscles, adipose tissue, and liver^{19,15,8}. This disease can be a major health problem and one of the leading causes of death. Recent record indicates that in 2010, an estimated 285 million people had diabetes of which type 2 were about 90% of the case^{18,2}. WHO predicts that diabetes mellitus patients in Indonesia increase from 8.4 million at 2000 to around 21.3 million at 2030. According to International diabetes Federation (IDF) is predicted diabetes mellitus patients increasing to be 12 million at 2030. Hyperglycemia is a widely known cause of enhanced plasma free radical concentration and development of oxidative stress. Impaired generation of naturally occurring antioxidants in diabetes can also be expected to result in increased oxidative cell damage¹⁹. The dietary antioxidants can reduce toxicity of free radicals and protect the body from oxidative damage¹⁵. Herb plants are usually used as anti-hyperglycemic agent because they are composed phytochemical compounds acted as antioxidant. Bioactive compounds of plants can be effective anti-diabetic agents if they have capacity as anti-hyperglycemic and antioxidant agents^{5,6}. Leaf of *Pluchea*

indica Less has been used by people as traditional medicine because this herb plant has been proven containing many phytochemical compounds as lignin, terpene, phenylpropanoid, benzoid, alkanes¹⁷, sterol, 2-(pro-1-ynyl)-5-(5,6-dihydroxy hexa-1,3-diynyl)-thiophene, (-)-catechin⁷, alkaloid⁴, saponin, tannin, phenol, hydroquinone, flavonoid^{23,24,25}, cardiac glycoside²⁵, flavonol (quercetin, kaempferol, myricetin)³. Widyawati⁹ et al. (2014) informed that different polarity of solvent (methanol, ethanol, ethyl acetate, hexanes, and water) can extract different phytochemical compounds that have various antioxidant activities. Methanolic extract of *Pluchea* leaf is the highest potential as antioxidant agent (DPPH scavenging activity and iron ion reducing power, 794.9 mg/GAE/g samples dry base and 2.14 mg GAE/g samples dry base, respectively) because contain the biggest phenolic and flavonoid contents, 1185.2 mg GAE/g samples dry base and 911.9 mg CE/g samples dry base). Pramanik et al. (2006) also said that methanolic extract of *Pluchea* can reduce in blood glucose in normal (35.12% and 36.01% for 200 and 400 mg/kg, respectively) and in streptozotocin induced diabetic rats (36.10% and 41.87% for 200 and 400 mg/kg, respectively). Srisook et al. (2012) has studied that hot water extract from *Pluchea*, herbal tea shows good antioxidant activity based on DPPH, superoxide and hydroxyl scavenging activities, ferric

*Author for Correspondence




Figure 1. The effect of various extracts of *Pluchea indica* Less leaf of blood glucose level

ion reducing power, and ferrous ion chelating. Thus, the objective of this research was to determine the effect of different polarity of solvent to phytochemical screening that cause reducing of blood glucose and influence toxicity level laboratory animal.

MATERIALS AND METHODS

Plant material

Leaves of *Pluchea indica* Less at 1-6 segment levels were collected from areas at East Coast, Bendul Merisi, Keputih, and Wiyungin Surabaya and Kertosono, East Java, Indonesia.

Chemicals and reagents

Glibenclamide was purchased from medicine store Surabaya, Indonesia, glucose assay kits (Sigma, USA). All the solvents and chemicals used were analytical grade.

Preparation of the plant extract

1-6 segment level of *Pluchea* leaves from peak was used as sample. These leaves were dried at ambient temperature and grinded with 45 mesh size. Dried flour of *Pluchea* leaves was measured moisture content. And then this flour was extracted by different polarity solvent (water, methanol, and ethyl acetate) with soxhlet extractor at a boiling point for three hours. Extract was evaporated by rotary evaporator. The extract was stored at 4°C in black glass bottle until analysis further.

Phytochemical screening

The freshly prepared crude extracts were qualitatively tested for presence of chemical constituents. These were identified by characteristic color changes using standard procedure. Phytochemical compounds in *Pluchea* leaves extract detected included alkaloid, flavonoid, phenolic, sterol, saponin, tannin, and cardiac glycoside.

Experimental animals

All the experiments were carried out using *Mus Musculus* Mice (20-30 g), Balb/c strain and *Rattus Norvegicus* Rats (125-150 g), Wistar strain of either sex that obtained from Rachmadi Priyadi's mice/rat husbandry, Surabaya, East Java, Indonesia. The animals were placed at random and allocated to treatment groups in polypropylene cages with paddy husk as bedding. Animals were housed at a temperature of $24 \pm 2^\circ\text{C}$ and relative humidity of 30-70%. A 12:12 light:day cycle was followed. All animals were allowed free access to water (*ad libitum*) and fed. The animals were used accordingly to guidelines of the Committee on Care and use of Experimental Animal Resources of LPPT-10 M, Jogjakarta, Indonesia.

Subchronic toxicity studies

Subchronic toxicity study was performed as per OECD-4 (1995). Albino mice was divided into 7 groups (n=8) of either sex selected by random sampling technique were used for the study. The animals were adapted in environmental and fed for one week before they were used. The animals were kept fasting for 8 hours providing only water, after that the suspended extract (methanol, ethyl acetate, and water) with CMC-Na 0.5% were administered orally at the dose level 1.3 mg/20g b.w. by intragastric tube and observed for 28 days. The effect of clinical of the experiment animal was observed for 4 hours at first day and the next day observation was done for 2 hours including deviation activity related to pharmacological screening assay and body healthy of the experiment animal. Morbidity and mortality of animals were observed at least 2 times a day including motoric activity, haging activity, retabliment, flexion, cornea condition, pineal, haffner, posture, catalepsy, and straub. If mortality was observed in one animal, then the same dose was repeated again to confirm the toxic dose. If mortality was not

Table 1: Phytochemical compounds in various *Pluchea* leaves extracts

Extract types	Color intensitas of phytochemical compounds							
	Terpenoid	Sterol	Flavonoid	Saponin	Tannin	Phenolic	Alkaloid	Cardiac glycoside
Aquadest	-	-	+++	++	+	+++	++++++	+++++
Methanol	-	+++	+++++	++++	++++	+++++	+++++	+++
Ethyl acetate	-	+++	++	+	-	++	+++	+

Note: + detected based on color intensity, - not detected based on color intensity

Table 2. Effect of *Pluchea indica* Less leaf extract on mortality in experimental animals

Groups	Sample	Day									Mortality
		1	2	3	4	5	6	7	8	9-28	
Male mice											
I	Control	0	0	0	0	0	0	0	0	0	0
II	Water extract at 1.3 mg/20g b.w.	0	0	0	0	0	0	0	0	0	0
III	Water extract at 2.6 mg/20g b.w.	0	0	0	0	0	0	0	0	0	0
IV	Methanol extract at 1.3 mg/20g b.w.	0	0	2	0	1	0	1	0	0	4
V	Methanol extract at 2.6 mg/20g b.w.	0	0	1	2	1	2	0	2	-	8
VI	Ethyl acetate extract at 1.3 mg/20g b.w.	0	1	3	1	2	1	-	-	-	8
VII	Ethyl acetate at 2,6 mg/20g b.w.	0	5	3	-	-	-	-	-	-	8
Female mice											
VIII	Control	0	0	0	0	0	0	0	0	0	0
IX	Water extract at 1.3 mg/20g b.w.	0	0	0	0	0	0	0	0	0	0
X	Water extract at 2.6 mg/20g b.w.	0	0	0	0	0	0	0	0	0	0
XI	Methanol extract at 1,3 mg/20g b.w.	0	0	2	0	0	0	1	0	2	5
XII	Methanol extract at 2.6 mg/20g b.w.	0	3	5	-	-	-	-	-	-	8
XIII	Ethyl acetate extract at 1.3 mg/20g b.w.	0	2	5	1	-	-	-	-	-	8
XIV	Ethyl acetate at 2,6 mg/20g b.w.	0	4	3	1	-	-	-	-	-	8

observed, the procedure was repeated for further higher such as 2.6 mg/20g b.w.

Effect of *Pluchea* extracts on normal blood glucose level

Rats were divided into five groups (n = 5) and fasted for 10 h before the experiment. Rats were administered 50% (5 ml/kg b.w.) load followed by suspended extracts (with CMC-Na 0.5%) of *Pluchea indica* Less leaves and reference drug suspended glibenclamide (with CMC-Na 0.5%) orally by a cannula just after checking the fasting blood glucose. First group received the control vehicle 0.5% w/v CMC-Na. The

methanol, ethyl acetate, and water extracts of *Pluchea indica* Less leaves (9 mg/200g animal body weight) were administered to second until fourth groups, respectively. The fifth group received the standard drug glibenclamide (0.126 mg/200 gb.w.) for assessing the comparative pharmacological significance^{21,11}. Blood was withdrawn from the tail vein by cutting the tip of the tail at 0, 15, 30, 60, 90 and 120 minutes of glucose administration and glucose levels were estimated immediately using compatible blood glucose test strips of glucometer (GlucoDr).

Table 3. Pharmacological screening assay of experimental animal after oral administration

Groups	Number	Motoric activity	Hanging activity	Retalishment	Flexion	Cornea Condition	Pineal	Haffner	Posture	Catal epsy	Straub
Male mice											
Control	1	+	+	+	+	+	+	-	+	-	-
	2	+	+	+	+	+	+	-	+	-	-
	3	+	+	+	+	+	+	-	+	-	-
	4	+	+	+	+	+	+	-	+	-	-
	5	+	+	+	+	+	+	-	+	-	-
	6	+	+	+	+	+	+	-	+	-	-
	7	+	+	+	+	+	+	-	+	-	-
	8	+	+	+	+	+	+	-	+	-	-
	1	+	+	+	+	+	+	-	+	-	-

Table 3. Pharmacological screening assay of experimental animal after oral administration

Groups	Number	Moto ric activi ty	Hagi ng activi ty	Retablis ment	Flexi on	Cornea Condition	Pineal	Haffner	Posture	Catal epsy	Straub
Water extract at 1.3 mg/20g b.w.	2	+	+	+	+	+	+	-	+	-	-
	3	+	+	+	+	+	+	-	+	-	-
	4	+	+	+	+	+	+	-	+	-	-
	5	+	+	+	+	+	+	-	+	-	-
	6	+	+	+	+	+	+	-	+	-	-
	7	+	+	+	+	+	+	-	+	-	-
	8	+	+	+	+	+	+	-	+	-	+
Water extract at 2.6 mg/20g b.w.	1	++	+	+	+	+	+	-	+	-	-
	2	++	+	+	+	+	+	-	+	-	-
	3	+++	-	+	+	+	+	+	+	-	-
	4	+	+	+	+	+	+	-	+	-	-
	5	+++	+	-	+	+	+	-	+	-	-
	6	+	+	+	+	+	+	-	+	-	-
	7	+++	+	+	+	+	+	-	+	-	-
	8	+++	+	+	+	+	+	-	+	-	-
Methanol extract at 1.3 mg/20g b.w.	1	+	+	+	+	+	+	-	+	-	-
	2	+	+	+	+	+	+	-	+	-	-
	3	+	+	+	+	+	+	-	+	-	-
	4	+	+	+	+	+	+	-	+	-	-
	5	+	+	+	+	+	+	-	+	-	-
	6	+	+	+	+	+	+	-	+	-	-
	7	+	+	+	+	+	+	-	+	-	-
	8	+	+	+	+	+	+	-	+	-	-
Methanol extract at 2.6 mg/20g b.w.	1-8	†	†	†	†	†	†	†	†	†	†
Ethyl acetate extract at 1.3 mg/20g b.w.	1-8	†	†	†	†	†	†	†	†	†	†
Ethyl acetate at 2.6 mg/20g b.w.	1-8	†	†	†	†	†	†	†	†	†	†
Female mice Control	1	+	+	+	+	+	+	-	+	-	-
	2	+	+	+	+	+	+	-	+	-	-
	3	+	+	+	+	+	+	-	+	-	-
	4	+	+	+	+	+	+	-	+	-	-
	5	+	+	+	+	+	+	-	+	-	-
	6	+	+	+	+	+	+	-	+	-	-
	7	+	+	+	+	+	+	-	+	-	-
	8	+	+	+	+	+	+	-	+	-	-
Water extract at 1.3 mg/20g b.w.	1	++	+	-	+	+	+	-	+	-	-
	2	++	+	-	+	+	+	-	+	-	-
	3	++	+	-	+	+	+	-	+	-	-
	4	++	+	-	+	+	+	-	+	-	-
	5	++	+	+	+	+	+	-	+	-	-
	6	++	+	-	+	+	+	-	+	-	-

Table 3. Pharmacological screening assay of experimental animal after oral administration

Groups	Number	Moto ric activi ty	Hagi ng activi ty	Retablis ment	Flexi on	Cornea Condition	Pineal	Haffner	Posture	Catal epsy	Straub
Water extract at 2.6 mg/20g b.w.	7	++	+	-	+	+	+	-	+	-	-
	8	+	+	-	+	+	+	-	+	-	-
	1	++	+	-	+	+	+	-	+	-	-
	2	++	+	-	+	+	+	-	+	-	-
	3	++	+	--	+	+	+	-	+	-	-
	4	++	+	-	+	+	+	-	+	-	-
	5	++	+	--	+	+	+	+	-	-	-
	6	++	+	-	+	+	+	+	-	-	-
Methanol extract at 1.3 mg/20g b.w.	7	++	+	-	+	+	+	+	-	-	-
	8	++	+	-	+	+	+	+	-	-	-
	1	+	+	-	+	+	+	+	-	+	-
	2	+	+	--	+	+	+	+	-	-	-
				(drop after large time)							
	3	+	+	---	+	+	+	+	-	-	-
				(drop directly)							
	4	+	+	---	+	+	+	+	-	-	-
Methanol extract at 2.6 mg/20g b.w.				(drop directly)							
	5	+	+	---	+	+	+	+	-	-	-
				(drop directly)							
	6	+	+	---	+	+	+	+	-	-	-
				(drop directly)							
	7	+	+	---	+	+	+	+	-	-	-
				(drop directly)							
	8	+	+	---	+	+	+	+	-	-	-
Ethyl acetate extract at 1.3 mg/20g b.w.				(drop directly)							
	1-8	†	†	†	†	†	†	†	†	†	†
	1-8	†	†	†	†	†	†	†	†	†	†
Ethyl acetate at 2,6 mg/20g b.w.											
	1-8	†	†	†	†	†	†	†	†	†	†

Note :+ or - = deviation of experimental animal, + = normal condition at motoric and hanging activities, reestablishment, posture, flexion, cornea condition, and pineal, - = normal condition at haffner, straub, catalepsy, and straub, † = mortality

RESULTS

Phytochemical compositions of various extracts of *Pluchea* leaves were showed at Table 1. Data informed that phytochemical compositions of three types of *Pluchea* leaf extract were different. The qualitative analysis based on color intensity showed that methanol solvent was the most effective to extract phytochemical compounds compared with the other solvents. Their compounds detected in methanol extract included sterol, flavonoid, saponin, tannin, phenolic, alkaloid and cardiac glycoside. Whereas water extract wasn't detected sterol and ethyl acetate extract wasn't detected tannin. Sub chronic study showed that methanol and ethyl acetate extracts of *Pluchea* leaves at 1.3 and 2.6 mg/20g b.w. caused mortality effect whereas the water extract are safe for long term administration (Table 2). Pharmacological screening effect (Table 3) showed that the oral administration of water extract at male mice was safe, but its consumption at 2.6 mg/20g b.w. could cause motoric activity increasing. Whereas female mice that was given water extract of *Pluchea* leaves at 1.3 and 2.6 mg/20 g b.w., respectively showed motoric activity increasing and reestablishment capacity decreasing. Methanol extract given at 1.3 mg/20g b.w. was no clinical effect at male experimental animals but it gave effect at female experimental animals. Methanol extract at 2.6 mg/20g b.w. and ethyl acetate at 1.3 and 2.6 mg/20g b.w., respectively caused mortality. Blood glucose level at experimental animal after oral administration from various extract of *Pluchea indica* Less showed at Figure 1. Water extract of *Pluchea* leaves was highest potentially to reduce blood glucose level compared with the other extract.

DISCUSSION

Our study showed that *Pluchea* leaves extract with various polarity of solvent possessed antidiabetic activity based on blood glucose reducing capacity. Adesegun et al. (2013); Kwon et al. (2006) informed that postprandial hyperglycemia is one of the risk factors associated with type 2 diabetes mellitus. Mohammed et al. (2006) said that hyperglycemia is caused by free radical accumulation on plasma and development of oxidative stress. *Pluchea indica* Less leaves contained phytochemical compounds so that could act as natural sources of antioxidants useful to health. Widyawati et al. (2014) has showed that leaves extract of *Pluchea* with various polarity of solvent can scavenge DPPH free radical and reduce iron ion. The observed activity may be due to the presence of chemical constituents such as phenolic compounds (tannins and flavonoids) in the extract. This argument is supported by^{10,14}. Phenolics have been reported to inhibit α -amylase activities. They also have anti-hyperglycemic activity and inhibit the development of diabetes^{12,26}. The antidiabetic activity of water extract was the highest, notwithstanding Widyawati et al. (2014) said that its DPPH scavenging activity is lower than methanol extract. It means that antidiabetic activity is not only

depended by hydrogen donating capacity of phenolic compounds. It was predicted that there were involvement constituent chemical reducing glucose (cardiac glycoside test). The oral administration test also showed that water extract was safe that it was potentially as antidiabetic agent.

CONFLICTS OF INTEREST

All contributing authors declare no conflicts of interest.

CONCLUSION

This study showed that the *Pluchea* leaf water extract possessed anti-diabetic activity based on blood glucose reducing capability. Its ability was 56.37 % that was the higher than glibenclamide (49.59%) and the other extracts (ethyl acetate extract = 19.11% and methanol extract = 24.27%). The water extract was safe to the body healthy. The anti-diabetic activity of this extract was supported by cardiac glucose compounds content.

ACKNOWLEDGEMENTS

The authors would like to thank the Higher Education Directorate (DIKTI) of the Indonesian's Government for the financial support (The competitive research grant) and Widya Mandala Catholic University in Surabaya for this research.

REFERENCES

- Adesegun SA, Fayemiwo O, Odufuye B, Coker, HAB. α -amylase inhibition and antioxidant activity of *Pterocarpus* Craib. Journal of Natural Products 2013; 6: 90-95.
- Akter F, Rahman MM, Mostofa M, Chowdhury EH. Anti-diabetic effect of neem and spirulina in alloxan induced diabetic mice. International Journal of Current Research and Academic Review 2014; 2(4):124-134.
- Andarwulan N, Batari R, Sandrasari D A, Bolling B, Wijaya H. Short communication flavonoid content and antioxidant activity of vegetables from Indonesia. Food Chemistry 2010; 121 : 1231-1235.
- Ardiansyah, Nuraida L, Andarwulan N. Aktivitas antimikroba daun beluntas (*Pluchea indica* Less) dan stabilitas aktivitasnya pada berbagai konsentrasi garam dan tingkat pH. Jurnal Teknologi dan Industri Pangan 2003; 14(2) : 90-97.
- Atawodi SE. Antioxidant potential of African medicinal plants. African Journal of Biotechnology 2005; 4(2): 128-129.
- Bello A, Aliero AA, Saidu Y, Muhammad S. Phytochemical screening, polyphenolic content and α -glucosidase inhibitory potential of *Leptadenia hastata* (Pers.) Decne. Nigerian Journal of Basic and Applied Science 2011; 19 (2): 181-186.
- Biswas R, Dutta PK, Achari B, Bandyopadhyay D, Mirsha M, Pramanik KC, Catterjee TK. Isolation, purification and characterization of four pure compounds from the root extract of *Pluchea indica* Less and the

- potentiality of the root extract and the pure compounds for antimicrobial activity. *European Bulletin of Drug Research* 2005; 13 : 63-70.
8. Bhutkar MA and Bhise SB. In vitro assay of alpha amylase inhibitory activity of some indigenous plants. *International Journal Chemical Science* 2012; 10(1): 457-462
9. Dorman HJD, Hiltunen R. Fe (III) reductive and free radical-scavenging properties of summer savory (*Satureja hortensis* L.) extract and sub fractions. *Food Chemistry* 2004; 88 : 193-199.
10. Ebi GC, Ofoefule SI. Antimicrobial activity of *Pterocarpus* stems. *Fitoterapia* 2006;71(4): 433-435.
11. Fahri C, Sutarno, Listyawati S. Blood glucose and total cholesterol content of hyperglycemic white male rat (*Rattus norvegicus* L.) after orally intakes of methanol meniran (*Phyllanthus Niruri* L.) root extract. *Biofarmasi* 2005; 3 (1): 1-6.
12. Hanamura T, Mayama C, Aoki H, Hirayama Y, Shimizu M. Antihyperglycemic effect of polyphenols from *Acerola* (*Malpighia marginata* DC.) fruit. *Bioscience Biotechnology Biochemical* 2006; 70: 1813-1820.
13. Harbone JB. Metode fitokimia. Padmawinata K, Soediro I, penerjemah, Bandung :Institut Teknologi Bandung, 1996.
14. Kim JS, Kwon CS, Son KH. Inhibition of alpha-glucosidase and amylase by luteolin, a flavonoid. *Bioscience Biotechnology Biochemical* 2000; 64: 2458-2461.
15. Kumar S, Rashmi, Kumar D. Evaluation of antidiabetic activity of *Euphorbia hirta* Linn. In streptozotocin induced diabetic mice. *Indian Journal of Natural Products and Resources* 2010; 1(2):200-203.
16. Kwon YI, Vattam DA, Shetty K. Evaluation of clonal herbs of Lamiaceae species for management of diabetes and hypertension. *Asia Pacific Journal Clinical and Nutrition* 2006; 15:107-118.
17. Luger P, Weber M, Dung XZ, Ngoc PH, Tuong DT, Rang DD. The crystal structure of hop-17(21)-en-3 β -yl asetat of *Pluchea pteropoda* Hemsl. from Vietnam. *Crystal Res Technology* 2000; 35(3) : 355-362.
18. Melmed S, Polonsky KS, Lasen PR, Kronenberg HM. Williams textbook of endocrinology (12th-ed). Philadelphia: Elsevier/Saunders 2012; pp. 1371-1435.
19. Mohammed HA, Shamma GAA, Hashim HM. β -carotene, glycemic control and dyslipidemia in type 2 diabetes mellitus. *Journal of the Faculty of Medicine, Baghdad* 2006; 48(4):435-441.
20. Pramanik KC, Brattacharya P, Biswas R, Bandyopadhyaya D, Mirsha M, Chatterjee TK. Hypoglycemic and antihyperglycemic activity of leaf extract of *Pluchea indica* Less. *Oriental Pharmacy and Experimental Medicine* 2006; 6(3):232-236.
21. Sharma SR, Dwivedi SK, Swarup D. Hypoglycaemic and hypolipidemic activity of *Caesalpinia bonducella* seeds in rats. *Journal of Ethnopharmacology* 1997; 58: 39-44.
22. Srisook K, Buapool D, Boonbai R, Simmasut P, Charoensuk Y, Srisook E. Antioxidant and anti-inflammatory activities of hot water extract from *Pluchea indica* Less herbal tea. *Journal of Medicinal Plants research* 2012; 6(23):4077-4081.
23. Widyawati PS, Wijaya CH, Harjosworo PS, Sajuthi D. Pengaruh ekstraksi dan fraksinasi terhadap aktivitas antioksidatif daun beluntas (*Pluchea indica* Less). *Jurnal Agroteknologi* 2010; 4(2):183-193.
24. Widyawati PS, Wijaya CH, Harjosworo PS, Sajuthi D. Evaluasi aktivitas antioksidatif ekstrak daun beluntas (*Pluchea indica* Less) berdasarkan perbedaan ruas daun. *Rekapangan Jurnal Teknologi Pangan* 2011; 5(1):1-14.
25. Widyawati PS, Budianta TDW, Kusuma FA, Wijaya EL. Difference of solvent polarity to phytochemical content and antioxidant activity of *Pluchea indica* Less leaves extracts *International of Pharmacology and Phytochemical Research* 2014; 6(4):850-855.
26. Zunino SJ, Storms DH, Stephensen CB. Diet rich in polyphenols and vitamin A inhibit the development of type1 autoimmune diabetes in nonobese diabetic mice. *Journal of Nutrition* 2007; 137: 1216-1221.

Evaluation Antidiabetic Activity of Various Leaf Extracts of *Pluchea indica* Less

ORIGINALITY REPORT

20%
SIMILARITY INDEX

18%
INTERNET SOURCES

12%
PUBLICATIONS

7%
STUDENT PAPERS

PRIMARY SOURCES

1 iqjmc.uobaghdad.edu.iq **2%**
Internet Source

2 journalofnaturalproducts.com **2%**
Internet Source

3 rjpbcs.com **2%**
Internet Source

4 chimie-biologie.ubm.ro **2%**
Internet Source

5 repository-tnmgrmu.ac.in **2%**
Internet Source

6 www.ijcrar.com **1%**
Internet Source

7 www.koreascience.or.kr **1%**
Internet Source

8 www.degruyter.com **1%**
Internet Source

9 www.neliti.com **1%**
Internet Source

10	tnsroindia.org.in Internet Source	1 %
11	ijnhs.net Internet Source	1 %
12	www.sapili.org Internet Source	1 %
13	Muhammad Zahoor, Wasim Ul Bari, Alam Zeb, Irfan Khan. "Toxicological, anticholinesterase, antilipidemic, antidiabetic and antioxidant potentials of Grewia optiva Drummond ex Burret extracts", Journal of Basic and Clinical Physiology and Pharmacology, 2020 Publication	1 %
14	Submitted to Universitas Katolik Widya Mandala Student Paper	1 %
15	www.sphinxsai.com Internet Source	1 %
16	C.T. Kumarappan, Subhash C. Mandal. " Polyphenolic Extract of Attenuates Diabetic Complications in Streptozotocin-Treated Diabetic Rats ", Renal Failure, 2009 Publication	<1 %
17	repository.maranatha.edu Internet Source	<1 %

18

PAINI SRI WIDYAWATI, CHRISTOFORA HANNY
WIJAYA, PENI SUPRPTI HARDJOSWORO,
DONDIN SAJUTHI. "Volatile Compounds of
Pluchea indica Less and Ocimum basilicum
Linn Essential Oil and Potency as Antioxidant",
HAYATI Journal of Biosciences, 2013

Publication

<1 %

19

core.ac.uk
Internet Source

<1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On