

BAB 7

KESIMPULAN DAN SARAN

7.1 Kesimpulan

Berdasarkan hasil penelitian mengenai Korelasi Lama Menderita DM dengan Skor MMSE yang dilaksanakan di Rumah Sakit Gotong Royong Surabaya, dapat disimpulkan sebagai berikut:

- Pasien DM dengan jenis kelamin perempuan lebih banyak dibandingkan dengan laki-laki.
- Pasien DM terbanyak ada pada rentang usia 61-70 tahun.
- Sebagian besar pasien DM yang berobat telah menderita DM kurang dari 5 tahun.
- Terdapat korelasi negatif, lemah, tidak bermakna antara lama DM dengan skor MMSE.
- Terdapat korelasi lemah, tidak bermakna antara lama DM dengan skor MMSE dengan mengendalikan variabel usia.

7.2 Saran

Berdasarkan hasil dan kesimpulan penelitian ini, peneliti menyarankan untuk melakukan edukasi tentang pentingnya deteksi dini DM, khususnya bagi mereka yang memiliki faktor risiko menderita DM.

Peneliti merekomendasikan kepada peneliti selanjutnya untuk meneliti lebih dalam mengenai DM dan HbA1c, serta hubungannya dengan fungsi kognitif khususnya pada populasi lansia, agar pencegahan dan deteksi dini demensia dapat terlaksana dengan baik.

DAFTAR PUSTAKA

1. PERKENI. Konsensus Pengendalian dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia 2015. Perkeni. 2015. 78 p.
2. International Diabetes Federation. Annual Report 2015. 2015;1–29.
3. Tuomilehto J, Schwarz PEH. Preventing diabetes: Early versus late preventive interventions. *Diabetes Care*. 2016;39(August):S115–20.
4. Goldman L, Shafer I A. Goldman-Cecil Medicine. *Goldman's Cecil Medicine*. 2016. 3024 p.
5. Tombaugh TN, McIntyre NJ. The Mini-Mental State Examination: A Comprehensive Review. *J Am Geriatr Soc*. 1992;40:922–35.
6. Larner A. Cognitive Screening Instruments. 2013.
7. World Health Organization. Global Report on Diabetes. Isbn. 2016;978:88.
8. Harrison TR. *Harrison's Manual of Medicine*. 18th ed. New York, N.Y.: McGraw-Hill Education LLC.; 2013.
9. Alzheimer Society of Canada. About Dementia | Risk factors [Internet]. 2017 [cited 2017 Dec 28]. Available from: <http://www.alzheimer.ca/en/Home/About-dementia/Alzheimer-s-disease/Risk-factors>
10. Kasper D, Fauci A, Hauser S, Longo D, Jameson J, Loscalzo J. *Harrison's Principles of Internal Medicine*. Mc GrawHill. 2015. 2958 p.
11. Rudijanto A, Soewondo P, Waspadji S, Yunir E, Purnamasari D. The Indonesian Society of Endocrinology's Summary Article of Diabetes Mellitus National Clinical Practice Guidelines. *J ASEAN Fed Endocr Soc*. 2011 May 31;26(1):17–9.
12. Barrett K, Barman S, Boitano S, Brooks H. *Ganong's Review of Medical Physiology*. New York, NY: McGraw-Hill. 2016.
13. John Edward Hall ACG. *Guyton and Hall Textbook of Medical Physiology*. Elsevier,. 2011. p. 1091 pages.
14. Elsevier. *Dorland's Illustrated Medical Dictionary* 32nd Edition. Saunders. 2012.
15. Khachiyants, Nina; Kim KY. Mini-mental status examination mapping to the corresponding brain areas in dementia. *Appl Technol Innov*. 2012;7(2):55–8.
16. Wrighten SA, Piroli GG, Grillo CA, Reagan LP. A look inside the diabetic brain: Contributors to diabetes-induced brain aging. *Biochim Biophys Acta - Mol Basis Dis* [Internet]. 2009;1792(5):444–53. Available from: <http://dx.doi.org/10.1016/j.bbadi.2008.10.013>
17. Nunley KA, Rosano C, Ryan CM, Jennings JR, Aizenstein HJ, Zgibor JC, et al. Clinically relevant cognitive impairment in Middle-Aged adults with childhood-onset type 1 diabetes. *Diabetes Care*. 2015;38(9):1768–76.
18. Mahfud RA, Lyrawati D, Sarwono I. Effect of Alpha Lipoic Acid on MDA Levels and Histology of Brain in Type 1 DM. *MNJ (Malang Neurol Journal)*. 2017 Jan 1;3(1):23–9.
19. Munshi MN. Cognitive dysfunction in older adults with diabetes: What a clinician needs to know. *Diabetes Care*. 2017;40(4):461–7.
20. Zheng T, Qin L, Chen B, Hu X, Zhang X, Liu Y, et al. Association of plasma

- DPP4 activity with mild cognitive impairment in elderly patients with type 2 diabetes: Results from the GDMD study in China. *Diabetes Care*. 2016;39(9):1594–601.
- 21. Tiehuis AM, van der Graaf Y, Visseren FL, Vincken KL, Biessels GJ, Appelman APA, et al. Diabetes increases atrophy and vascular lesions on brain MRI in patients with symptomatic arterial disease. *Stroke*. 2008;39(5):1600–3.
 - 22. Zhang T, Shaw M, Humphries J, Sachdev P, Anstey KJ, Cherbuin N. Higher fasting plasma glucose is associated with striatal and hippocampal shape differences: the 2sweet project. *BMJ Open Diabetes Res Care*. 2016;4(1):e000175.
 - 23. Alzheimer's Association. Mild Cognitive Impairment (MCI) | Signs, Symptoms, & Diagnosis [Internet]. [cited 2017 Dec 28]. Available from: <https://www.alz.org/dementia/mild-cognitive-impairment-mci.asp>
 - 24. Hjelm K, Atwine F. Health-care seeking behaviour among persons with diabetes in Uganda: an interview study. *BMC Int Health Hum Rights*. 2011;11(1):11.
 - 25. Vlassoff C. Gender differences in determinants and consequences of health and illness. *J Heal Popul Nutr*. 2007;25(1):47–61.
 - 26. Centers for Disease Control and Prevention. National Diabetes Statistics Report: Estimates of Diabetes and Its Burden in the United States. Atlanta, GA: Centers for Disease Control and Prevention; 2014. US Dep Heal Hum Serv. 2017;(Cdc):2009–12.
 - 27. Foundation THJKF. Percentage of Adults with Diagnosed Diabetes by Age Group [Internet]. 2014 [cited 2017 Nov 28]. Available from: <https://www.kff.org/other/state-indicator/adults-with-diabetes-by-age/>
 - 28. Zoungas S, Woodward M, Li Q, Cooper ME, Hamet P, Harrap S, et al. Impact of age, age at diagnosis and duration of diabetes on the risk of macrovascular and microvascular complications and death in type 2 diabetes. *Diabetologia*. 2014;57(12):2465–74.
 - 29. Alencar R, Cobas R, Gomes M. Assessment of cognitive status in patients with type 2 diabetes through the mini-mental status examination: a cross-sectional study. *Diabetol Metab Syndr* [Internet]. 2010;2(1):10. Available from: <http://www.dmsjournal.com/content/2/1/10>