

BUKTI KORESPONDENSI

Judul Artikel	:	Separation and analysis of <i>Monascus</i> yellow pigments produced on durian seed substrate
Jurnal	:	Food Research Volume 4, Issue 4, August 2020
Penulis	:	*Srianta, I. , Nugerahani, I. and Ristiarini, S.

No	Perihal	Tanggal
1	Bukti konfirmasi submit artikel beserta cover letter, formulir submission, dan artikel yang di-submit	10 Januari 2020
2	Bukti konfirmasi respon penerimaan artikel beserta letter to author	15 Januari 2020
3	Bukti konfirmasi hasil review dan permintaan revisi artikel beserta manuscript evaluation form dan hasil review dari ketiga reviewer	7 Februari 2020
4	Bukti konfirmasi submit revisi artikel beserta respon kepada reviewer dan artikel yang di-resubmit	2 Maret 2020
5	Bukti konfirmasi artikel diterima dan acceptance letter	7 Maret 2020
6	Bukti konfirmasi permintaan klarifikasi beberapa poin di artikel sebelum masuk tahap produksi	23 Maret 2020
7	Bukti konfirmasi respon atas permintaan klarifikasi beberapa poin di artikel sebelum masuk tahap produksi	24 Maret 2020
8	Bukti konfirmasi respon penerimaan klarifikasi beberapa poin di artikel sebelum masuk tahap produksi	24 Maret 2020
9	Bukti konfirmasi galley proof artikel	26 Maret 2020
10	Bukti konfirmasi submit permintaan revisi galley proof artikel pertama dan artikel yang di-resubmit	26 Maret 2020
11	Bukti konfirmasi respon atas permintaan revisi galley proof artikel pertama	27 Maret 2020
12	Bukti konfirmasi submit permintaan revisi galley proof artikel kedua dan artikel yang di-resubmit	27 Maret 2020
13	Bukti konfirmasi persetujuan galley proof artikel	27 Maret 2020
14	Bukti konfirmasi artikel published online dan artikel yang diterbitkan	1 April 2020

1

**Bukti konfirmasi submit artikel beserta cover letter,
formulir submission, dan artikel yang di-submit**

10 Januari 2020

Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Srianta, I. *, Nugerahani, I. and Ristiarini, S.

*Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala Catholic University
Surabaya, Jalan Dinoyo 42-44 Surabaya, Indonesia 60265*

*Corresponding author: srianta_wm@yahoo.com

Author No.1: <https://orcid.org/0000-0001-9810-7735>

Author No. 2: <https://orcid.org/0000-0002-9917-5649>

Author No.3: <https://orcid.org/0000-0003-2387-3819>

Abstract

This research was aimed to separate and analysis of monascus yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate:ethanol:water=90:25:4. The separated MYP then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has maximum absorption at 386 nm. Five MYP compounds detected by using LC-MS/MS, 2 of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in the MYP. The MYP is being further studied for the antidiabetic bioactivity.

Keywords: monascus, yellow pigment, durian seed, separation, analysis

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through polyketide pathway. It has been used as natural food colourant, food supplement, and in traditional medicine. The monascus pigments consists of many pigment compounds which are categorized into 3 groups i.e. red, orange and yellow pigments. Six well known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srianta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, 39 new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid state fermentation substrate in monascus pigments production. Non-rice substrates have been developed in the monascus pigment e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srianta *et al.*, 2012; Srianta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of monascus yellow pigments is still limited. MYP is

not only potential as natural colorant for biscuit, mayonaisse, wheat noodle, etc. (Srianta *et al.*, 2014), but also as functional food ingredient and in medicine because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), anti-obesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) show that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in monascus pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various monascus pigments because its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analysis MYP produced on durian seed substrate. This is the first investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and Methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid state fermentation on durian seed substrate. The culture was maintain periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya. The durian seed is by product of the food industry, and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid state fermentation with durian seed substrate

Pigments production was performed through solid state fermentation on durian seed substrate according to our previous research (Srianta *et al.*, 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 minutes and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hours, ground and sieved.

2.3 Monascus pigment separation

The monascus pigment was extracted with soxhletation method in ethanolic solvent at ratio of 1:50 for 2 hours. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre eluted with ethyl acetate:ethanol:water=90:25:4. The extract was put into the column and eluted with ethyl acetate:ethanol:water=90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate:ethanol:water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer

(Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through 0.2 μ m PTFE filter membrane, put on to the auto sampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with an isocratic program of mobile phase consist of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and Discussion

In the pigments separation by using column chromatography, 4 fractions (F1 F2, F3 and F4) derived with different color. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that the MYP was the pigments with the lowest polarity among the others. Although in the chemical structure of monascus pigments are similar each other which consist of β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments. The presence of $-\text{NH}_3$ group can increase the polarity of red pigments.

Figure 2 show the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and Yellow II. Monascin and Yellow II were the major pigment compounds in the MYP with relative amount of 59,81% and 27,80%. The MYP composition depends on the pigment composition in MFDS, extraction and separation processes.

Pigments in MFDS is produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but still unclear and controversy. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment Rubropunctatin biosynthesis, it consist of β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the monascus pigment composition. Ethanolic extract contain higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated the MYP from the red and orange pigments.

Monascin, the major MYP compound in this research, has attracted for their strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and antiatherosclerosis activities, also improve memory and learning ability. On the other hand, there is no report on the yellow II bioactivity. In conclusion, the separated MYP has great potential as functional ingredient in food and pharmaceuticals. The MYP is being further studied for the antidiabetic bioactivity.

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in the MYP.

Conflict of Interest

Authors declare no conflict of interest.

Acknowledgments


This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [grant number 200X/WM01.5/N/2019].

References

- Agboyibor, C., Kong, W.B., Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447.
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565.
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98, 1554-1560.
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136.
- Feng, Y., Shao, Y. and Chen, F. (2012). *Monascus* pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440.
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented *dioscorea*. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518.
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264, 117–124.

- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology* 49, 2609–2617.
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18, 387-399.
- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143.
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390.
- Pattanagul, P., Pinthong, R., Phianmongkhol, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.
- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19, 941-945.
- Srianta, I., Ristiarini, S., Nugerahani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. *International Food Research Journal*, 21, 1-12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatalysis and Agricultural Biotechnology*, 7, 181-186.
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from *Monascus* fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *Journal of Agricultural and Food Chemistry*, 53, 1949-1954.
- Vidyalakshmi, R., Paranthaman, R., Muruges, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. *Global Journal of Biotechnology & Biochemistry*, 4, 84–87.
- Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. *Marine Drugs*, 11, 3902-3925.

Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




(A)




(B)


Figure 2. Scanning results of separated Monascus pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




F1



F2



F3



F4

Mass spectrum of the sample showing relative intensity versus m/z . The base peak is at m/z 359.5. Other significant peaks are labeled at m/z 101.0, 143.4, 167.0, 192.2, 279.3, 295.5, 320.5, 399.6, 422.4, 463.6, 503.8, and 519.5.

Mass spectrum showing relative intensity (Y-axis, 0.00 to 1.75) versus mass-to-charge ratio (X-axis, 100 to 600 m/z). The spectrum displays several peaks, with the base peak at m/z 385.1. Other labeled peaks include m/z 105.1, 279.0, and 423.0.

m/z	Relative Intensity (approx.)
105.1	0.25
279.0	0.10
385.1	1.00
423.0	0.15

(C)

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected



srianta ignatius <srianta2601@gmail.com>

Short communication manuscript submission

srianta ignatius <srianta2601@gmail.com>

10 Januari 2020 pukul 23.06

Kepada: Food Research <foodresearch.my@outlook.com>

Dear Prof. Son Radu
Editor in Chief of Food Research

Please kindly find the attached files of cover letter, submission form and manuscript titled "separation and analysis of monascus yellow pigment produced on durian seed substrate" (Authors: Ignatius Srianta, Susana Ristiarini and Ira Nugerahani) for your consideration and possible publication in Food Research journal.,

I am looking forward to being informed the evaluation process

Best regards
Srianta

3 lampiran



Cover letter_Srianta_FR.docx
18K



manuscript_submission_form_filled.docx
60K



Short communication ms_monascus yellow pigment_Srianta.docx
158K

Please fill in your manuscript details in 'New Manuscript Submission Section' and submit this form together with your manuscript.

Authors are requested to suggest at least 3 – 4 potential reviewers by filling in their particulars in the 'Suggested Reviewer' section.

NEW MANUSCRIPT SUBMISSION

Manuscript Title	Separation and analysis of Monascus yellow pigments produced on durian seed substrate		
Manuscript Type <i>(Please Bold)</i>	Original Article	Review	
	Short Communication	Technical Notes	
Authors	Ignatius Srianta, Susana Ristiarini and Ira Nugerahani		
Corresponding Author <i>(Only one)</i>	Ignatius Srianta		
Email address of the Corresponding Author	Srianta_wm@yahoo.com		

SUGGESTED REVIEWERS

Name	Philippe J. Blanc
Salutation	PhD
Area of Expertise <i>(Please use comma for more than one)</i>	Bioprocess, biotechnology, fungal pigment
Email	philippe.blanc1@univ-tlse2.fr

Name	Erning Indrastuti
Salutation	Dr
Area of Expertise <i>(Please use comma for more than one)</i>	Agricultural product technology
Email	indrastuti_erning@yahoo.com

Name	Ihab Tewfik
Salutation	PhD
Area of Expertise <i>(Please use comma for more than one)</i>	Nutrition, nutraceutical, functional food, food analysis
Email	I.Tewfik@westminster.ac.uk

Name	Chatarina Yayuk Trisnawati
Salutation	MSc
Area of Expertise <i>(Please use comma for more than one)</i>	Food science, food chemistry
Email	tchatarinayayuk@gmail.com

January 10, 2020

Professor Dr. Son Radu
Chief Editor
Food Research
foodresearch.my@outlook.com

Dear Sir,

COVER LETTER FOR MANUSCRIPT SUBMISSION TO FOOD RESEARCH

Please include your manuscript title and your significant results and impact of your research to be considered for publication in Food Research.

On behalf of my colleagues, I wish to submit the attached short communication manuscript titled "Separation and analysis of *Monascus* yellow pigments produced on durian seed substrate" for your consideration and possible publication in the Food Research journal.

The manuscript is on separation of *Monascus* yellow pigments produced on durian seed substrate with solid state fermentation, and the pigments analysis by using spectrophotometry, thin layer chromatography (TLC) and liquid chromatography-mass spectrometry (LC-MS).

We produced the pigments with *Monascus purpureus* M9 on durian seed according to our previous method. The *Monascus* yellow pigments was separated on a silica gel column, eluted with ethyl acetate:ethanol:water=90:25:4. This method was developed in our laboratory. The profile and composition of the *Monascus* yellow pigment were investigated.

The study on the separation and analysis of *Monascus* yellow pigments produced on durian seed substrate has never been carried out before. The *Monascus* yellow pigments contained several yellow pigment compounds that potential as functional natural food colorant. These would give benefit to your journal and the readers.

I also would like to suggest 4 potential reviewers as mentioned in the manuscript submission form.

I thank you for considering this manuscript. Correspondence should be addressed to Dr. Ignatius Srianata, email: srianata_wm@yahoo.com . I look forward to being updated on the progress of the manuscript in the evaluation process.

Your sincerely,



Dr. Ignatius Srianata
Department of Food Technology
Faculty of Agricultural Technology
Widya Mandala Catholic University Surabaya
Jalan Dinoyo 42-44, Surabaya
Indonesia 60625,

2

**Bukti konfirmasi respon penerimaan artikel beserta letter
to author**

15 Januari 2020



srianta ignatius <srianta2601@gmail.com>

FR-2020-020

Food Research <foodresearch.my@outlook.com>
Kepada: srianta ignatius <srianta2601@gmail.com>

15 Januari 2020 pukul 17.42

Dear Srianta,

This message is to acknowledge the receipt of the above manuscript that you submitted via email to Food Research. Your manuscript has been successfully checked-in. Please refer to the assigned manuscript ID number in any correspondence with the Food Research Editorial Office or with the editor.

Your paper will be reviewed by three or more reviewers assigned by the Food Research editorial board and final decision made by the editor will be informed by email in due course. Reviewers' suggestions and editor's comments will be then made available via email attached file. You can monitor the review process for your paper by emailing us on the "Status of my manuscript".

If your manuscript is accepted for publication, Food Research editorial office will contact you for the production of your manuscript.

Thank you very much for submitting your manuscript to Food Research.

Sincerely,

Professor Dr. Son Radu
Chief Editor
Email: foodresearch.my@outlook.com

From: srianta ignatius <srianta2601@gmail.com>
Sent: Saturday, 11 January, 2020 12:06 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Short communication manuscript submission

Dear Prof. Son Radu
Editor in Chief of Food Research

Please kindly find the attached files of cover letter, submission form and manuscript titled "separation and analysis of monascus yellow pigment produced on durian seed substrate" (Authors: Ignatius Srinta, Susana Ristiarini and Ira Nugerahani) for your consideration and possible publication in Food Research journal.,

I am looking forward to being informed the evaluation process

Best regards
Srinta

 **Letter to Author FR-2020-020.pdf**
22K

15th January 2020

Authors: Srianta, I., Nugerahani, I. and Ristiarini, S.

Manuscript title: Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Manuscript ID: FR-2020-020

Dear Srianta,

This message is to acknowledge the receipt of the above manuscript that you submitted via email to Food Research. Your manuscript has been successfully checked-in. Please refer to the assigned manuscript ID number in any correspondence with the Food Research Editorial Office or with the editor.

Your paper will be reviewed by three or more reviewers assigned by the Food Research editorial board and final decision made by the editor will be informed by email in due course. Reviewers' suggestions and editor's comments will be then made available via email attached file. You can monitor the review process for your paper by emailing us on the "Status of my manuscript".

If your manuscript is accepted for publication, Food Research editorial office will contact you for the production of your manuscript.

Thank you very much for submitting your manuscript to Food Research.

Sincerely,



Professor Dr. Son Radu
Chief Editor
Email: foodresearch.my@outlook.com

3

**Bukti konfirmasi hasil review dan permintaan revisi
artikel beserta manuscript evaluation form dan hasil
review dari ketiga reviewer**

7 Februari 2020

Separation and analysis of *Monascus* yellow pigment produced on durian seed substrate

Abstract

This research was aimed to separate and analysis of monascus yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate:ethanol:water=90:25:4. The separated MYP then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has maximum absorption at 386 nm. Five MYP compounds detected by using LC-MS/MS, 2 of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in the MYP. The MYP is being further studied for the antidiabetic bioactivity.

Keywords: monascus, yellow pigment, durian seed, separation, analysis

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through polyketide pathway. It has been used as natural food colourant, food supplement, and in traditional medicine. The monascus pigments consists of many pigment compounds which are categorized into 3 groups i.e. red, orange and yellow pigments. Six well known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srinta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, 39 new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid state fermentation substrate in monascus pigments production. Non-rice substrates have been developed in the monascus pigment e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srinta *et al.*, 2012; Srinta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of monascus yellow pigments is still limited. MYP is not only potential as natural colorant for biscuit, mayonnaise, wheat noodle, etc. (Srinta *et al.*, 2014), but also as functional food ingredient and in medicine because MYP has various possitive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), anti-obesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) show that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in monascus pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various monascus pigments because its high sensitivity and

Commented [t1]: Because of

multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analysis MYP produced on durian seed substrate. This is the first investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and Methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid state fermentation on durian seed substrate. The culture was maintain periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya. The durian seed is by product of the food industry, and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid state fermentation with durian seed substrate

Pigments production was performed through solid state fermentation on durian seed substrate according to our previous research (Srianta *et al.*, 2012). After peeling, cutting, soaking, sterilization at 121oC for 10 minutes and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hours, ground and sieved.

2.3 Monascus pigment separation

The monascus pigment was extracted with soxhletation method in ethanolic solvent at ratio of 1:50 for 2 hours. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre eluted with ethyl acetate:ethanol:water=90:25:4. The extract was put into the column and eluted with ethyl acetate:ethanol:water=90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate:ethanol:water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through 0.2 um PTFE filter membrane, put on to the auto sampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with an isocratic program of mobile phase consist of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and Discussion

In the pigments separation by using column chromatography, 4 fractions (F1 F2, F3 and F4) derived with different color. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and

Commented [t2]: from

pigments profile on TLC shown in Figure 1. Those profiles describe that the MYP was the pigments with the lowest polarity among the others. Although in the chemical structure of monascus pigments are similar each other which consist of β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments. The presence of $-\text{NH}_3$ group can increase the polarity of red pigments.

Figure 2 show the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and Yellow II. Monascin and Yellow II were the major pigment compounds in the MYP with relative amount of 59,81% and 27,80%. The MYP composition depends on the pigment composition in *Monascus Fermented Durian Seeds* (MFDS), extraction and separation processes.

Pigments in MFDS is produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but still unclear and controversy. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment Rubropunctatin biosynthesis, it consist of β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the monascus pigment composition. Ethanolic extract contain higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated the MYP from the red and orange pigments.

Monascin, the major MYP compound in this research, has attracted for their strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and antiatherosclerosis activities, also improve memory and learning ability. On the other hand, there is no report on the yellow II bioactivity. In conclusion, the separated MYP has great potential as functional ingredient in food and pharmaceuticals. The MYP is being further studied for the antidiabetic bioactivity.

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in the MYP.

Conflict of Interest

Authors declare no conflict of interest.

Acknowledgments

This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik


Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [grant number 200X/WM01.5/N/2019].

References

- Agboyibor, C., Kong, W.B., Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447.
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565.
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98, 1554-1560.
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136.
- Feng, Y., Shao, Y. and Chen, F. (2012). *Monascus* pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440.
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518.
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264, 117–124.
- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology* 49, 2609–2617.
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18, 387-399.
- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143.
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390.

- Pattanagul, P., Pinthong, R., Phianmongkhon, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). Chiang Mai Journal of Science, 34, 319–328.
- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. International Food Research Journal, 19, 941-945.
- Srianta, I., Ristiarini, S., Nugerahani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. International Food Research Journal, 21, 1-12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. Biocatalysis and Agricultural Biotechnology, 7, 181-186.
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from *Monascus* fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. Journal of Agricultural and Food Chemistry, 53, 1949-1954.
- Vidyalakshmi, R., Paranthaman, R., Muruges, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. Global Journal of Biotechnology & Biochemistry, 4, 84–87.
- Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. Marine Drugs, 11, 3902-3925.

Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




(A)




(B)


Figure 2. Scanning results of separated Monascus pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




F1



F2




F3




F4


Figure 3. LC-MS analysis results: Total Ion Chromatogram of MYP (A) and representatives of chromatogram of major MYP monascin (B) and yellow II (C).



(A)



(B)



(C)

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

FR-2020-020

1 message

Food Research <foodresearch.my@outlook.com>
To: ignatius srianta <srianta_wm@yahoo.com>

Fri, 7 Feb 2020 at 23:55

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research

6 attachments

- ☐ **Evaluation Form FR-2020-020 .doc**
261 KB
- ☐ **Evaluation Form FR-2020-020.doc**
260 KB
- ☐ **Evaluation Form FR-2020-020-1.doc**
260 KB
- ☐ **FR-2020-020.docx**
210 KB
- ☐ **FR-2020-020-1.docx**
211 KB
- ☐ **FR-2020-020-2.docx**
213 KB

MANUSCRIPT EVALUATION FORM

Date : 15th January 2020

Manuscript ID : FR-2020-020

Please return by : 15th February 2020

Title of Manuscript : Separation and analysis of Monascus yellow pigment
produced on durian seed substrate

1. IF YOU CANNOT REVIEW THIS MANUSCRIPT OR MEET THE DEADLINE, PLEASE INFORM US WITHOUT DELAY.
2. Your review should consider the article's scholarly merit including originality of the research issue and/or methodology, adequacy and rigor of the research methodology and techniques used, quality and rigor of data analysis, comprehensiveness of literature review, and the readability and presentation of the article. Please provide detailed and specific comments to all items. Also, where appropriate please provide suggestions for revision.

COMMENT SHEET

Using item 2 in page 1 as a guideline, please indicate the reasons for your recommendations. Most author(s) will appreciate frankness, combined with a modicum of tact. Even if you recommend that the manuscript be accepted for publication, please provide some general comments to the author(s).

Evaluation Criteria	Grade				
	A (Excellent)	B	C	D	E (Worst)
1. Appropriateness of Contents		X			
2. Originality of Topic		X			
3. Manuscript Format		X			
4. Research Methodology		X			
5. Data Analysis		X			
6. Relevance to the Journal		X			

FOOD RESEARCH

(REVIEWER'S SECTION)		(AUTHOR'S SECTION)
REVIEWER'S COMMENTS/SUGGESTIONS		AUTHOR'S ACTION/RESPONSE
		*NOTE FOR AUTHOR: Please state your response to the reviewer's comments/suggestion below
1.	Title <i>"Separation and analysis of Monascus yellow pigment produced on durian seed substrate "</i> <i>This title reflects the article</i>	
2.	Abstract <i>This research was aimed to separate and analyse Monascus yellow pigment with chromatography. Monascin and yellow II were the major pigment compounds in MYP.</i>	
3.	Keywords <i>Monascus, yellow pigment, durian seed</i>	
4.	Introduction <i>The introduction is concise with sufficient references to the topic.</i>	
5.	Research design/Methodology <i>Materials and Methods are well described and probably reproducible</i>	
6.	Data Analysis <i>The results are well presented and discussed. The main results is that monascin and yellow II were the major pigments in MYP. The indication of further studies to evaluate antidiabetic bioactivity is a little premature and useless.</i>	
7.	Conclusion <i>The conclusion is short and is a clear summary of the study</i>	
8.	References	
9.	English Proficiency <i>It has really to be improved and corrected by an English-speaking person.</i>	

10.	Additional comments/suggestions by the reviewer about the article <i>It's a pity not to have information on the pigment bioactivity.</i>	
-----	--	--

Overall Evaluation

Please choose one.

Accept		Major Revision	
Minor Revision	X	Reject	

This article is relevant as it identified two yellow pigments produced by Monascus on an unusual substrate, the durian seed. It should be accepted if English proficiency is improved.

Please return Manuscript and/or Review Comments to:

Professor Dr. Son Radu

Food Research

Email: foodresearch.my@outlook.com

MANUSCRIPT EVALUATION FORM

Date : 15th January 2020

Manuscript ID : FR-2020-020

Please return by : 15th February 2020

Title of Manuscript : Separation and analysis of Monascus yellow pigment
produced on durian seed substrate

1. IF YOU CANNOT REVIEW THIS MANUSCRIPT OR MEET THE DEADLINE, PLEASE INFORM US WITHOUT DELAY.
2. Your review should consider the article's scholarly merit including originality of the research issue and/or methodology, adequacy and rigor of the research methodology and techniques used, quality and rigor of data analysis, comprehensiveness of literature review, and the readability and presentation of the article. Please provide detailed and specific comments to all items. Also, where appropriate please provide suggestions for revision.

COMMENT SHEET

Using item 2 in page 1 as a guideline, please indicate the reasons for your recommendations. Most author(s) will appreciate frankness, combined with a modicum of tact. Even if you recommend that the manuscript be accepted for publication, please provide some general comments to the author(s).

Evaluation Criteria	Grade				
	A (Excellent)	B	C	D	E (Worst)
1. Appropriateness of Contents	A				
2. Originality of Topic	A				
3. Manuscript Format		B			
4. Research Methodology	A				
5. Data Analysis	A				
6. Relevance to the Journal	A				

FOOD RESEARCH

(REVIEWER'S SECTION)		(AUTHOR'S SECTION)
REVIEWER'S COMMENTS/SUGGESTIONS		AUTHOR'S ACTION/RESPONSE
		*NOTE FOR AUTHOR: Please state your response to the reviewer's comments/suggestion below
1.	Title <i>It should reflect the article</i>	Fine
2.	Abstract <i>Background, Aim, Methodology and Conclusion</i>	Fine, just in once sentence add the significance of your study to the field of Food Research
3.	Keywords <i>Min. 3 and Max. 6</i>	Fine
4.	Introduction <i>Concise with sufficient background</i>	Fine
5.	Research design/Methodology <i>Clearly described and reproducible</i>	Fine
6.	Data Analysis <i>Results well presented and discussed</i>	Fine
7.	Conclusion <i>A clear summary of the study</i>	Please revisit [see attached document]
8.	References <i>References should follow the journal's format</i>	Good relevant set of references
9.	English Proficiency	Please pay attention to some typos

10.	Additional comments/suggestions by the reviewer about the article	
-----	---	--

Overall Evaluation

Please choose one.

Accept		Major Revision	
Minor Revision	X	Reject	

Please return Manuscript and/or Review Comments to:

Professor Dr. Son Radu

Food Research

Email: foodresearch.my@outlook.com

MANUSCRIPT EVALUATION FORM

Date : 15th January 2020

Manuscript ID : FR-2020-020

Please return by : 15th February 2020

Title of Manuscript : Separation and analysis of Monascus yellow pigment produced on durian seed substrate

1. IF YOU CANNOT REVIEW THIS MANUSCRIPT OR MEET THE DEADLINE, PLEASE INFORM US WITHOUT DELAY.
2. Your review should consider the article's scholarly merit including originality of the research issue and/or methodology, adequacy and rigor of the research methodology and techniques used, quality and rigor of data analysis, comprehensiveness of literature review, and the readability and presentation of the article. Please provide detailed and specific comments to all items. Also, where appropriate please provide suggestions for revision.

COMMENT SHEET

Using item 2 in page 1 as a guideline, please indicate the reasons for your recommendations. Most author(s) will appreciate frankness, combined with a modicum of tact. Even if you recommend that the manuscript be accepted for publication, please provide some general comments to the author(s).

Evaluation Criteria	Grade				
	A (Excellent)	B	C	D	E (Worst)
1. Appropriateness of Contents	√				
2. Originality of Topic	√				
3. Manuscript Format		√			
4. Research Methodology	√				
5. Data Analysis	√				
6. Relevance to the Journal	√				

FOOD RESEARCH

(REVIEWER'S SECTION)		(AUTHOR'S SECTION)
		AUTHOR'S ACTION/RESPONSE
		*NOTE FOR AUTHOR: Please state your response to the reviewer's comments/suggestion below
1.	Title <i>It should reflect the article</i>	
2.	Abstract <i>Background, Aim, Methodology and Conclusion</i>	
3.	Keywords <i>Min. 3 and Max. 6</i>	
4.	Introduction <i>Concise with sufficient background</i>	
5.	Research design/Methodology <i>Clearly described and reproducible</i>	
6.	Data Analysis <i>Results well presented and discussed</i>	
7.	Conclusion <i>A clear summary of the study</i>	
8.	References <i>References should follow the journal's format</i>	Minor revision on the writing the citation and references. Writing the journal's name is italicized. A space in one reference is one.
9.	English Proficiency	

FOOD RESEARCH

10.	Additional comments/suggestions by the reviewer about the article	The word <i>Monascus</i> is italicized.
-----	--	---

Overall Evaluation

Please choose one.

Accept		Major Revision	
Minor Revision	v	Reject	

Please return Manuscript and/or Review Comments to:

Professor Dr. Son Radu

Food Research

Email: foodresearch.my@outlook.com

Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Abstract

This research was aimed to separate and analysis of monascus yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate:ethanol:water=90:25:4. The separated MYP then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has maximum absorption at 386 nm. Five MYP compounds detected by using LC-MS/MS, 2 of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in the MYP. The MYP is being further studied for the antidiabetic bioactivity.

Keywords: monascus, yellow pigment, durian seed, separation, analysis

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through polyketide pathway. It has been used as natural food colourant, food supplement, and in traditional medicine. The monascus pigments consists of many pigment compounds which are categorized into 3 groups i.e. red, orange and yellow pigments. Six well known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srinta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, 39 new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid state fermentation substrate in monascus pigments production. Non-rice substrates have been developed in the monascus pigment e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srinta *et al.*, 2012; Srinta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of monascus yellow pigments is still limited. MYP is not only potential as natural colorant for biscuit, mayonnaise, wheat noodle, etc. (Srinta *et al.*, 2014), but also as functional food ingredient and in medicine because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), anti-obesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) show that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in monascus pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various monascus pigments ~~because~~because its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analysis MYP produced on durian seed substrate. This is the first investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and Methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the ~~solid-state~~solid-state fermentation on durian seed substrate. The culture was ~~mantain~~maintain periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya. The durian seed is by product of the food industry, and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid state fermentation with durian seed substrate

Pigments production was performed through solid state fermentation on durian seed substrate according to our previous research (Srianta *et al.*, 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 minutes and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hours, ground and sieved.

2.3 Monascus pigment separation

The monascus pigment was extracted with soxhletation method in ethanolic solvent at ratio of 1:50 for 2 hours. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre eluted with ethyl acetate:ethanol:water=90:25:4. The extract was put into the column and eluted with ethyl acetate:ethanol:water=90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate:ethanol:water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through 0.2 μ m PTFE filter membrane, put on to the auto sampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with an isocratic program of mobile phase consist of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and Discussion

In the pigments separation by using column chromatography, 4 fractions (F1 F2, F3 and F4) derived with different color. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that the MYP was the pigments with the lowest polarity among the others. Although in the chemical structure of monascus pigments are similar each other which consist of β -ketoacid and chromophore, MYP compounds contain less

Formatted: Superscript

polar groups than those of red pigments. The presence of $-NH_3$ group can increase the polarity of red pigments.

Figure 2 shows the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and Yellow II. Monascin and Yellow II were the major pigment compounds in the MYP with relative amount of 59,81% and 27,80%. The MYP composition depends on the pigment composition in MFDS, extraction and separation processes.

Pigments in MFDS is produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but still unclear and controversy. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment Rubropunctatin biosynthesis, it consist of β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the monascus pigment composition. Ethanolic extract contain higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated the MYP from the red and orange pigments.

Monascin, the major MYP compound in this research, has attracted for their strong bioactivities such as anticancer, anti-tumor, anti-diabetic, antioxidative stress, anti-inflammatory, anti-obesity and showed anti-atherosclerosis-atherogenic activities, also improve memory and learning ability. On the other hand, there is no report on the yellow II bioactivity. In conclusion, the separated MYP has great potential as functional ingredient in food and pharmaceuticals. The MYP is being further studied for the anti-diabetic bioactivity.

Formatted: Font: Bold, Underline, Highlight

4. Conclusion

Formatted: Underline, Highlight

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in the MYP.

Formatted: Underline, Highlight

Conflict of Interest

Authors declare no conflict of interest.

Acknowledgments

This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [grant number 200X/WM01.5/N/2019].

References

- Agboyibor, C., Kong, W.B. , Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447.
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565.
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98, 1554-1560.
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136.
- Feng, Y., Shao, Y. and Chen, F. (2012). *Monascus* pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440.
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518.
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264, 117–124.
- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology* 49, 2609–2617.
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18, 387-399.
- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143.
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390.
- Pattanagul, P., Pinthong, R., Phianmongkhon, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.
- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19, 941-945.

Formatted: Indonesian


Formatted: Indonesian

Formatted: Indonesian

Formatted: Indonesian

- Srianta, I., Ristiarini, S., Nugerahani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. *International Food Research Journal*, 21, 1-12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatalysis and Agricultural Biotechnology*, 7, 181-186.
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from *Monascus* fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *Journal of Agricultural and Food Chemistry*, 53, 1949-1954.
- Vidyalakshmi, R., Paranthaman, R., Muruges, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. *Global Journal of Biotechnology & Biochemistry*, 4, 84-87.
- Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. *Marine Drugs*, 11, 3902-3925.

Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




(A)




(B)


Figure 2. Scanning results of separated Monascus pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




F1



F2




F3




F4


Figure 3. LC-MS analysis results: Total Ion Chromatogram of MYP (A) and representatives of chromatogram of major MYP monascin (B) and yellow II (C).



(A)



(B)



(C)

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Abstract

This research was aimed to separate and analysis ~~of M~~onascus yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate:ethanol:water=90:25:4. The separated MYP ~~were~~ then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has ~~a~~ maximum absorption at 386 nm. ~~On f~~ive MYP compounds detected by using LC-MS/MS, 2 of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in ~~the~~ MYP. ~~The MYP is being further studied for the antidiabetic bioactivity.~~

Keywords: ~~M~~onascus, yellow pigment, durian seed, separation, analysis

Formatted: English (United States)

1. Introduction

Monascus pigments are secondary metabolites of Monascus fungi biosynthesized through polyketide pathway. It has been used as natural food colourant, food supplement, and in traditional medicine. ~~M~~The monascus pigments consists of many pigment compounds which are categorized into 3 groups i.e. red, orange and yellow pigments. Six well known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srinta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, 39 new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid state fermentation substrate in ~~monascus-Monascus~~ pigments production. Non-rice substrates have been developed in the ~~monascus-Monascus~~ pigment e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srinta *et al.*, 2012; Srinta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of ~~monascus-Monascus~~ yellow pigments is still limited. MYP is not only potential as natural colorant for biscuit, mayonnaise, wheat noodle, etc. (Srinta *et al.*, 2014), but also as functional food ingredient and in medicine because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), anti-obesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) show that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in ~~monascus-Monascus~~ pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various ~~monascus-Monascus~~ pigments because its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

Formatted: English (United States)

Formatted: English (United States)

This research was aimed to separate and analyse MYP produced on durian seed substrate. This is the first investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and Methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid state fermentation on durian seed substrate. The culture was maintained periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya (Indonesia). The durian seed is a by product of the food industry, and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid state fermentation with durian seed substrate

Pigments production was performed through solid state fermentation on durian seed substrate according to our previous research (Srianta *et al.*, 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 minutes and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hours, ground and sieved.

2.3 Monascus pigment separation

The monascus pigment was extracted with soxhletation method in ethanolic solvent at ratio of 1:50 for 2 hours. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre-eluted with ethyl acetate:ethanol:water=90:25:4. The extract was put into the column and eluted with ethyl acetate:ethanol:water=90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate:ethanol:water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through 0.2 µm PTFE filter membrane, put on to the auto sampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with an isocratic program of mobile phase consist of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and Discussion

In the pigments separation by using column chromatography, 4 fractions (F1 F2, F3 and F4) derived with different color. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that the MYP was the pigments with the lowest polarity among the others. Although in the chemical structure of *Monascus* pigments are similar each other which consist for all of them of β-ketoacid and chromophore, MYP

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

compounds contain less polar groups than those of red pigments. The presence of –NH₃ group can increase the polarity of red pigments.

Figure 2 show the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and Yellow II. Monascin and Yellow II were the major pigment compounds in the MYP with relative amount of 59,81% and 27,80%. The MYP composition depends on the pigment composition in MFDS, extraction and separation processes.

Pigments in MFDS is produced through a complex polyketide biosynthesis pathway in the Monascus fungi, but it is still unclear and controversy. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment rubropunctatin biosynthesis, it consist of β-ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the Monascus pigment composition. Ethanolic extract contain higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated the MYP from the red and orange pigments.

Monascin, the major MYP compound in this research, has-is attractiveed for their-its strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and antiatherosclerosis activities, and also improves memory and learning ability. On the other hand, there is no report on the yellow II bioactivity. In conclusion, the separated MYP has great potential as a functional ingredient in food and pharmaceuticals. The MYP is-being-will be further studied for the antidiabetic bioactivity.

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in the MYP.

Conflict of Interest

Authors declare no conflict of interest.

Acknowledgments

This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [grant number 200X/WM01.5/N/2019].

References

Formatted: English (United States)

Formatted: English (United States)

Commented [MOU1]: What is MFDS

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Agboyibor, C., Kong, W.B., Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447.

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562-565.

Formatted

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98, 1554-1560.

Formatted

Formatted

Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136.

Formatted

Feng, Y., Shao, Y. and Chen, F. (2012). *Monascus* pigments. *Applied Microbiology and Biotechnology*, 96, 1421-1440.

Formatted

Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512-4518.

Formatted

Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264, 117-124.

Formatted

Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology* 49, 2609-2617.

Formatted

Formatted

Formatted

Field Code Changed

Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18, 387-399.

Formatted

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143.

Formatted

Field Code Changed

Field Code Changed

Nimnoi, P. and Lumyong, S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384-1390.

Field Code Changed

Field Code Changed

Pattanaagul, P., Pinthong, R., Phianmongkhon, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319-328.

Field Code Changed

Formatted

Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19, 941-945.


Field Code Changed

Formatted

Formatted

Formatted

Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




(A)




(B)


Figure 2. Scanning results of separated Monascus pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




F1



F2



F3



F4



Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

4

**Bukti konfirmasi submit revisi artikel beserta respon
kepada reviewer dan artikel yang di-resubmit**

2 Maret 2020

Re: FR-2020-020 (Revised)

Dari: srianta_wm@yahoo.com

Kepada: foodresearch.my@outlook.com

Tanggal: Senin, 2 Maret 2020 pukul 08.00 GMT+7

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards
Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research



FR-2020-020_Revised version.docx

196.8kB



FR-2020-020_Response to The Reviewers Comments.docx

16.6kB

Response to The Reviewers Comments
FR-2020-020

No.	Section	Page	Line	Reviewer Comment	Response
1.	Abstract	1	1	Analysis Monascus	It has been revised: Analyse <i>Monascus</i>
		1	3	The separated MYP then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS.	It has been modified: The separated MYP were then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS.
		1	4	Five MYP compounds detected by using LC-MS/MS, 2 of them were well known yellow pigment compounds, monascin and ankaflavin	It has been modified: On five MYP compounds detected by using LC-MS/MS, 2 of them were well known yellow pigment compounds, monascin and ankaflavin
		1	5	Monascin and yellow II were the major pigment compounds in the MYP.	It has been revised: Monascin and yellow II were the major pigment compounds in MYP.
		1	6	Reviewer suggestion: "The MYP is being further studied for the antidiabetic bioactivity" to be deleted	It has been deleted
2.	Introduction	1	3	The monascus pigments	It has been revised: <i>Monascus</i> pigments
		1	6,7,9,13, 15	monascus pigments	It has been revised: <i>Monascus</i> pigments
		1	10	mayonnaise	It has been revised: mayonnaise
		1	10	possitive	It has been revised: positive
		1	11	anti-obesity	It has been revised: antiobesity activity
		1	15	because its high	It has been modified: because of its high
		2	16	analysis	It has been revised: analyse
3,	Material and Methods	2	1	solid state	It has been revised: solid-state
		2 2	2 3	mantain Durian seed was obtained from a durian processing unit in Surabaya	It has been revised: maintain It has been modified: Durian seed was obtained from a durian processing unit in Surabaya (Indonesia)
4.	Result and Discussion	2	1	In the pigments separation by using column	It has been modified: In the pigments separation by using column chromatography, 4

		2	5	chromatography, 4 fractions (F1 F2, F3 and F4) derived with different color monascus pigments	fractions (F1 F2, F3 and F4) derived from different color
		3	5	Although in the chemical structure of <i>Monascus</i> pigments are similar each other which consist of β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments.	It has been revised: <i>Monascus</i> pigments It has been modified: Although the chemical structure of <i>Monascus</i> pigments consist for all of them of β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments.
		3	13	The MYP composition depends on the pigment composition in MFDS	It has been revised: The MYP composition depends on the pigment composition in <i>Monascus</i> Fermented Durian Seed (MFDS)
		3	16	Rubropunctatin	It has been revised: rubropunctatin
		3	18	monascus pigments	It has been revised: <i>Monascus</i> pigments
		3	21	Monascin, the major MYP compound in this research, has attracted for their strong bioactivities such as anticancer, anti-tumor, anti-diabetic, antioxidative stress, anti-inflammatory, anti-obesity and anti-atherosclerosis activities, also improve memory and learning ability.	It has been modified: Monascin, the major MYP compound in this research, is attractive for its strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and showed antiatherogenic activities, and also improves memory and learning ability.
		3	23, 24	Reviewer suggestion: "In conclusion, the separated MYP has great potential as functional ingredient in food and pharmaceuticals. The MYP is being further studied for the antidiabetic bioactivity" to be deleted	It has been deleted
	English			It has really to be	It has been done

	proficiency			improved and corrected by an English-speaking person	
--	-------------	--	--	--	--

Separation and analysis of *Monascus* yellow pigment produced on durian seed substrate

Abstract

This research was aimed to separate and analyse *Monascus* yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate:ethanol:water=90:25:4. The separated MYP were then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has a maximum absorption at 386 nm. On five MYP compounds detected by using LC-MS/MS, 2 of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in MYP.

Keywords: *Monascus*, yellow pigment, durian seed, separation, analysis

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through polyketide pathway. It has been used as natural food colourant, food supplement, and in traditional medicine. *Monascus* pigments consists of many pigment compounds which are categorized into 3 groups i.e. red, orange and yellow pigments. Six well known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srianta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, 39 new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid state fermentation substrate in *Monascus* pigments production. Non-rice substrates have been developed in the *Monascus* pigment e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srianta *et al.*, 2012; Srianta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of *Monascus* yellow pigments is still limited. MYP is not only potential as natural colorant for biscuit, mayonnaise, wheat noodle, etc. (Srianta *et al.*, 2014), but also as functional food ingredient and in medicine because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), antiobesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) show that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in *Monascus* pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various *Monascus* pigments because of its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analyse MYP produced on durian seed substrate. This is the first investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and Methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid-state fermentation on durian seed substrate. The culture was maintained periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya (Indonesia). The durian seed is a by-product of the food industry, and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid state fermentation with durian seed substrate

Pigments production was performed through solid state fermentation on durian seed substrate according to our previous research (Srianta *et al.*, 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 minutes and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hours, ground and sieved.

2.3 Monascus pigment separation

The monascus pigment was extracted with soxhlet method in ethanolic solvent at ratio of 1:50 for 2 hours. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre-eluted with ethyl acetate:ethanol:water=90:25:4. The extract was put into the column and eluted with ethyl acetate:ethanol:water=90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate:ethanol:water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through 0.2 µm PTFE filter membrane, put on to the auto sampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with an isocratic program of mobile phase consist of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and Discussion

In the pigments separation by using column chromatography, 4 fractions (F1 F2, F3 and F4) derived from different color. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that MYP was the pigments with the lowest polarity among the others. Although the chemical structure of *Monascus* pigments

consist for all of them of β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments. The presence of $-\text{NH}_3$ group can increase the polarity of red pigments. Figure 2 show the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and Yellow II. Monascin and Yellow II were the major pigment compounds in the MYP with relative amount of 59,81% and 27,80%. The MYP composition depends on the pigment composition in *Monascus* Fermented Durian Seed (MFDS), extraction and separation processes. Pigments in MFDS produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but it is still unclear and controversy. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment rubropunctatin biosynthesis, it consist of β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the *Monascus* pigment composition. Ethanolic extract contain higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated MYP from the red and orange pigments. Monascin, the major MYP compound in this research, is attractive for its strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and showed antiatherogenic activities, and also improves memory and learning ability. On the other hand, there is no report on the yellow II bioactivity.

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in MYP.

Conflict of Interest

Authors declare no conflict of interest.

Acknowledgments


This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [grant number 200X/WM01.5/N/2019].

References

- Agboyibor, C., Kong, W.B. , Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447.
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565.
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98, 1554-1560.
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136.
- Feng, Y., Shao, Y. and Chen, F. (2012). *Monascus* pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440.
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518.
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264, 117–124.
- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology* 49, 2609–2617.
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18, 387-399.
- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143.
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390.
- Pattanagul, P., Pinthong, R., Phianmongkhol, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.

- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19, 941-945.
- Srianta, I., Ristiarini, S., Nugerahani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. *International Food Research Journal*, 21, 1-12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatalysis and Agricultural Biotechnology*, 7, 181-186.
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from *Monascus* fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *Journal of Agricultural and Food Chemistry*, 53, 1949-1954.
- Vidyalakshmi, R., Paranthaman, R., Muruges, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. *Global Journal of Biotechnology & Biochemistry*, 4, 84–87.
- Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. *Marine Drugs*, 11, 3902-3925.

Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.



(A)



(B)

Figure 2. Scanning results of separated Monascus pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.







Figure 3. LC-MS analysis results: Total Ion Chromatogram of MYP (A) and representatives of chromatogram of major MYP monascin (B) and yellow II (C).



(A)



(B)



(C)

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

5

Bukti konfirmasi artikel diterima dan acceptance letter

7 Maret 2020

Re: FR-2020-020 - Decision on your manuscript

Dari: Food Research (foodresearch.my@outlook.com)

Kepada: srianta_wm@yahoo.com

Tanggal: Sabtu, 7 Maret 2020 pukul 11.52 GMT+7

Dear Prof Srianta,

It is a pleasure to accept your manuscript for publication in Food Research journal. Please refer to the attachment for your acceptance letter. I will contact you again once the galley proof is ready for viewing and approval.

Thank you for your fine contribution. We look forward to your continued contributions to the Journal.

Sincerely,
Dr. Vivian New
Editor
Food Research

From: srianta_wm@yahoo.com <srianta_wm@yahoo.com>

Sent: Monday, 2 March, 2020 9:00 AM

To: Food Research <foodresearch.my@outlook.com>

Subject: Re: FR-2020-020 (Revised)

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards
Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate "

which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research



FR-2020-020 Acceptance Letter.pdf
31.1kB

7th March 2020

Dear Prof Dr Srianta, I.,

ACCEPTANCE LETTER

Food Research, is pleased to inform you that the following manuscript has been accepted for publication in Food Research journal.

Manuscript Title : Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Authors : Srianta, I., Nugerahani, I. and Ristiarini, S.

We thank you for your fine contribution to the Food Research journal and encourage you to submit other articles to the Journal.

Yours sincerely,



Professor Dr. Son Radu
Chief Editor
Food Research



6

**Bukti konfirmasi permintaan klarifikasi beberapa poin di
artikel sebelum masuk tahap produksi**

23 Maret 2020

Re: FR-2020-020 - Article Production

Dari: Food Research (foodresearch.my@outlook.com)

Kepada: srianta_wm@yahoo.com

Tanggal: Senin, 23 Maret 2020 pukul 16.25 GMT+7

Dear Prof Srianta,

Manuscript ID: FR-2020-020

Manuscript Title: Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,

Vivian New

Editor

Food Research

From: srianta_wm@yahoo.com <srianta_wm@yahoo.com>

Sent: Monday, 2 March, 2020 9:00 AM

To: Food Research <foodresearch.my@outlook.com>

Subject: Re: FR-2020-020 (Revised)

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards

Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research



FR-2020-020.docx
212.1kB

Separation and analysis of *Monascus* yellow pigment produced on durian seed substrate

*Srianta, I., Nugerahani, I. and Ristiarini, S.

*Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala
Catholic University Surabaya, Jalan Dinoyo 42-44 Surabaya, Indonesia 60265*

*Corresponding author: srianta_wm@yahoo.com

Article history:

Received: 15 January 2020

Received in revised form: 2 March 2020

Accepted: 5 March 2020

Abstract

This research was aimed to separate and analyse *Monascus* yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate: ethanol: water = 90: 25: 4. The separated MYP were then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has a maximum absorption at 386 nm. Out of five MYP compounds detected by using LC-MS/MS, two of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in MYP.

Keywords: *Monascus*, Yellow pigment, Durian seed, Separation, Analysis

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through the polyketide pathway. It has been used as a natural food colourant, food supplement, and in traditional

medicine. *Monascus* pigments consist of many pigment compounds which are categorized into 3 groups i.e. red, orange and yellow pigments. A total of six well-known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srinta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, 39 new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid-state fermentation substrate in *Monascus* pigments production. Non-rice substrates have been developed in the *Monascus* pigment e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srinta *et al.*, 2012; Srinta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of *Monascus* yellow pigments is still limited. MYP is not only a potential natural colorant for biscuit, mayonnaise, wheat noodle, etc. (Srinta *et al.*, 2014) but also as functional food ingredient and in medicine, because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), antiobesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) show that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in *Monascus* pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various *Monascus* pigments because of its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analyse MYP produced on durian seed substrate. This is the first investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid-state fermentation on durian seed substrate. The culture was maintained periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya (Indonesia). The durian seed is a by-product of the food industry and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid-state fermentation with durian seed substrate

Pigments production was performed through solid-state fermentation on durian seed substrate according to our previous research (Srianta *et al.*, 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 mins and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hrs, ground and sieved.

2.3 *Monascus* pigment separation

The monascus pigment was extracted with soxhletation method in ethanolic solvent at a ratio of 1:50 for 2 hrs. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre eluted with ethyl acetate: ethanol: water = 90:25:4. The extract was put into the column and eluted with ethyl acetate: ethanol: water = 90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate: ethanol: water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through a 0.2 µm PTFE filter membrane, put on to the autosampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with an isocratic program of mobile phase consists of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and discussion

In the separation of the pigments by using column chromatography, 4 fractions (F1 F2, F3 and F4) derived from different color. The first fraction was MYP target, while other fractions were orange and

red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that MYP was the pigments with the lowest polarity among the others. Although the chemical structure of *Monascus* pigments consists of all of them of β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments. The presence of $-\text{NH}_3$ group can increase the polarity of red pigments.

Figure 2 shows the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in the orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and Yellow II. Monascin and Yellow II were the major pigment compounds in the MYP with the relative amount of 59.81% and 27.80%. The MYP composition depends on the pigment composition in *Monascus* Fermented Durian Seed (MFDS), extraction and separation processes.

Pigments in MFDS produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but it is still unclear and controversy. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment rubropunctatin biosynthesis, it consist of the β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the *Monascus* pigment composition. The ethanolic extract contains higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated MYP from the red and orange pigments.

Monascin, the major MYP compound in this research, is attractive for its strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and showed antiatherogenic activities, and also improves memory and learning ability. On the other hand, there is no report on the yellow II bioactivity.

Commented [VN1]: Please rephrase. The sentence is very confusing.

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in MYP.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [grant number 200X/WM01.5/N/2019].


References

- Agboyibor, C., Kong, W.B., Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447.
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565.
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98, 1554-1560.
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136.

- Feng, Y., Shao, Y. and Chen, F. (2012). Monascus pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440.
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518.
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264, 117–124.
- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology*, 49, 2609–2617.
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18, 387–399.
- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143.
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390.
- Pattanagul, P., Pinthong, R., Phianmongkhon, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.
- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19, 941–945.
- Srianta, I., Ristiarini, S., Nugerhani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. *International Food Research Journal*, 21, 1–12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatalysis and Agricultural Biotechnology*, 7, 181–186.
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from *Monascus* fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *Journal of Agricultural and Food Chemistry*, 53, 1949–1954.
- Vidyalakshmi, R., Paranthaman, R., Muruges, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. *Global Journal of Biotechnology and Biochemistry*, 4, 84–87.

Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. *Marine Drugs*, 11, 3902-3925.

Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.



(A)



(B)

Figure 2. Scanning results of separated *Monascus* pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.







Figure 3. LC-MS analysis results: Total Ion Chromatogram of MYP (A) and representatives of the chromatogram of major MYP monascin (B) and yellow II (C).



(A)



(B)



(C)

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

7

**Bukti konfirmasi respon atas permintaan klarifikasi
beberapa poin di artikel sebelum masuk tahap produksi**

24 Maret 2020

Re: FR-2020-020 - Article Production

Dari: ignatius srianta (srianta_wm@yahoo.com)

Kepada: foodresearch.my@outlook.com

Tanggal: Selasa, 24 Maret 2020 pukul 01.18 GMT+7

Dear Dr. Vivian
Please kindly the attached file of corrected FR-2020-020. Thank you
Best regards
Srianta

Pada Senin, 23 Maret 2020 16.25.24 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Manuscript ID: FR-2020-020

Manuscript Title: Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,
Vivian New
Editor
Food Research

From: srianta_wm@yahoo.com <srianta_wm@yahoo.com>
Sent: Monday, 2 March, 2020 9:00 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 (Revised)

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards
Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research



FR-2020-020_corrected.docx
211.5kB

Separation and analysis of *Monascus* yellow pigment produced on durian seed substrate

*Srianta, I., Nugerahani, I. and Ristiarini, S.

*Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala
Catholic University Surabaya, Jalan Dinoyo 42-44 Surabaya, Indonesia 60265*

*Corresponding author: srianta_wm@yahoo.com

Article history:

Received: 15 January 2020

Received in revised form: 2 March 2020

Accepted: 5 March 2020

Abstract

This research was aimed to separate and analyse *Monascus* yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate: ethanol: water = 90: 25: 4. The separated MYP were then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has a maximum absorption at 386 nm. Out of five MYP compounds detected by using LC-MS/MS, two of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in MYP.

Keywords: *Monascus*, Yellow pigment, Durian seed, Separation, Analysis

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through the polyketide pathway. It has been used as a natural food colourant, food supplement, and in traditional

medicine. *Monascus* pigments consist of many pigment compounds which are categorized into 3 groups i.e. red, orange and yellow pigments. A total of six well-known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srinta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, 39 new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid-state fermentation substrate in *Monascus* pigments production. Non-rice substrates have been developed in the *Monascus* pigment e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srinta *et al.*, 2012; Srinta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of *Monascus* yellow pigments is still limited. MYP is not only a potential natural colorant for biscuit, mayonnaise, wheat noodle, etc. (Srinta *et al.*, 2014) but also as functional food ingredient and in medicine, because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), antiobesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) show that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in *Monascus* pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various *Monascus* pigments because of its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analyse MYP produced on durian seed substrate. This is the first investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid-state fermentation on durian seed substrate. The culture was maintained periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya (Indonesia). The durian seed is a by-product of the food industry and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid-state fermentation with durian seed substrate

Pigments production was performed through solid-state fermentation on durian seed substrate according to our previous research (Srianta *et al.*, 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 mins and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hrs, ground and sieved.

2.3 *Monascus* pigment separation

The monascus pigment was extracted with soxhletation method in ethanolic solvent at a ratio of 1:50 for 2 hrs. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre eluted with ethyl acetate: ethanol: water = 90:25:4. The extract was put into the column and eluted with ethyl acetate: ethanol: water = 90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate: ethanol: water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through a 0.2 µm PTFE filter membrane, put on to the autosampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with an isocratic program of mobile phase consists of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and discussion

In the separation of the pigments by using column chromatography, 4 fractions (F1 F2, F3 and F4) with different color have been derived. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that MYP was the pigments with the lowest polarity among the others. Although the chemical structure of *Monascus* pigments consists of all of them of β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments. The presence of $-\text{NH}_3$ group can increase the polarity of red pigments. Figure 2 shows the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in the orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and Yellow II. Monascin and Yellow II were the major pigment compounds in the MYP with the relative amount of 59.81% and 27.80%. The MYP composition depends on the pigment composition in *Monascus* Fermented Durian Seed (MFDS), extraction and separation processes. Pigments in MFDS produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but it is still unclear and controversy. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment rubropunctatin biosynthesis, it consist of the β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the *Monascus* pigment composition. The ethanolic extract contains higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated MYP from the red and orange pigments. Monascin, the major MYP compound in this research, is attractive for its strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and showed antiatherogenic activities, and also improves memory and learning ability. On the other hand, there is no report on the yellow II bioactivity.

Commented [VN1]: Please rephrase. The sentence is very confusing.

Commented [A2R1]:

Commented [A3R1]:

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in MYP.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [grant number 200X/WM01.5/N/2019].


References

- Agboyibor, C., Kong, W.B., Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447.
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565.
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98, 1554-1560.
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136.


- Feng, Y., Shao, Y. and Chen, F. (2012). Monascus pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440.
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from Monascus-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518.
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of Monascus pigments. *FEMS Microbiology Letters*, 264, 117–124.
- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The Monascus metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology*, 49, 2609–2617.
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). Monascus-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18, 387–399.
- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native Monascus pigments. *Frontiers in Microbiology*, 9, 3143.
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390.
- Pattanagul, P., Pinthong, R., Phianmongkhon, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.
- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19, 941–945.
- Srianta, I., Ristiarini, S., Nugerahani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. *International Food Research Journal*, 21, 1–12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatalysis and Agricultural Biotechnology*, 7, 181–186.
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from Monascus fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *Journal of Agricultural and Food Chemistry*, 53, 1949–1954.
- Vidyalakshmi, R., Paranthaman, R., Murugesh, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. *Global Journal of Biotechnology and Biochemistry*, 4, 84–87.

Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. *Marine Drugs*, 11, 3902-3925.

Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.



(A)



(B)

Figure 2. Scanning results of separated *Monascus* pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.







Figure 3. LC-MS analysis results: Total Ion Chromatogram of MYP (A) and representatives of the chromatogram of major MYP monascin (B) and yellow II (C).



(A)



(B)



(C)

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

8

**Bukti konfirmasi respon penerimaan klarifikasi beberapa
poin di artikel sebelum masuk tahap produksi**

24 Maret 2020

Re: FR-2020-020 - Article Production

Dari: Food Research (foodresearch.my@outlook.com)

Kepada: srianta_wm@yahoo.com

Tanggal: Selasa, 24 Maret 2020 pukul 11.30 GMT+7

Dear Prof Srianta,

Received with thanks.

Thanks & Regards,
Vivian New
Editor
Food Research

From: ignatius srianta <srianta_wm@yahoo.com>
Sent: Tuesday, 24 March, 2020 2:18 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian
Please kindly the attached file of corrected FR-2020-020. Thank you
Best regards
Srianta

Pada Senin, 23 Maret 2020 16.25.24 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Manuscript ID: FR-2020-020

Manuscript Title: Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,
Vivian New
Editor
Food Research

From: srianta_wm@yahoo.com <srianta_wm@yahoo.com>
Sent: Monday, 2 March, 2020 9:00 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 (Revised)

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards
Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,

Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research

9

Bukti konfirmasi galley proof artikel

26 Maret 2020

Re: FR-2020-020 - Article Production

Dari: Food Research (foodresearch.my@outlook.com)

Kepada: srianta_wm@yahoo.com

Tanggal: Kamis, 26 Maret 2020 pukul 11.47 GMT+7

Dear Prof Srianta,

Please refer to the attachment for the galley proof of your manuscript FR-2020-020 entitled 'Title' . Please check the content of the galley proof. If there are any mistakes, please comment and highlight in the PDF itself and revert to us within two (2) days of receipt. Once we have finalized the PDF version, your manuscript will be published online for early viewing.

Please see the attachment for the invoice INV20060. We hope that you can make the payment as soon as possible before 22 July 2020 for us to complete the publication of your manuscript. The manuscript information e.g. volume, issue, page numbers and DOI, will be provided once we have received the payment.

Thanks & Regards,
Vivian New
Editor
Food Research

From: ignatius srianta <srianta_wm@yahoo.com>
Sent: Tuesday, 24 March, 2020 2:18 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian
Please kindly the attached file of corrected FR-2020-020. Thank you
Best regards
Srianta

Pada Senin, 23 Maret 2020 16.25.24 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Manuscript ID: FR-2020-020

Manuscript Title: Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,
Vivian New
Editor
Food Research

From: srianta_wm@yahoo.com <srianta_wm@yahoo.com>

Sent: Monday, 2 March, 2020 9:00 AM

To: Food Research <foodresearch.my@outlook.com>

Subject: Re: FR-2020-020 (Revised)

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards
Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript

should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research



INV20060.pdf
165.4kB



FR-2020-020.pdf
565.1kB

Separation and analysis of *Monascus* yellow pigment produced on durian seed substrate

*Srianta, I., Nugerahani, I. and Ristiarini, S.

Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala Catholic University
Surabaya, Jalan Dinoyo 42-44 Surabaya, Indonesia 60265

Article history:

Received: 15 January 2020

Received in revised form: 2 March 2020

Accepted: 5 March 2020

Available Online:

Keywords:

Monascus,
Yellow pigment,
Durian seed,
Separation,
Analysis

DOI:

dfid

Abstract

This research was aimed to separate and analyse *Monascus* yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate: ethanol: water = 90: 25: 4. The separated MYP were then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has a maximum absorption at 386 nm. Out of five MYP compounds detected by using LC-MS/MS, two of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in MYP.

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through the polyketide pathway. It has been used as a natural food colourant, food supplement, and in traditional medicine. *Monascus* pigments consist of many pigment compounds which are categorized into three groups i.e. red, orange and yellow pigments. A total of six well-known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srianta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, thirty-nine new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid-state fermentation substrate in *Monascus* pigments production. Non-rice substrates have been developed in the *Monascus* pigment e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srianta *et al.*, 2012; Srianta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of *Monascus* yellow pigments is still limited. MYP is not only a potential natural colorant for biscuit, mayonnaise, wheat noodle, etc. (Srianta *et al.*, 2014) but also as functional food ingredient and in medicine,

because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), antiobesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) show that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in *Monascus* pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various *Monascus* pigments because of its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analyse MYP produced on durian seed substrate. This is the first

*Corresponding author.

Email: srianta_wm@yahoo.com

investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid-state fermentation on durian seed substrate. The culture was maintained periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya (Indonesia). The durian seed is a by-product of the food industry and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid-state fermentation with durian seed substrate

Pigments production was performed through solid-state fermentation on durian seed substrate according to our previous research (Srianta et al., 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 mins and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hrs, ground and sieved.

2.3 Monascus pigment separation

The *Monascus* pigment was extracted with soxhletation method in ethanolic solvent at a ratio of 1:50 for 2 hrs. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre eluted with ethyl acetate: ethanol: water = 90:25:4. The extract was put into the column and eluted with ethyl acetate: ethanol: water = 90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate: ethanol: water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through a 0.2 µm PTFE filter membrane, put on to the autosampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with

an isocratic program of mobile phase consists of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and discussion

In the separation of the pigments by using column chromatography, 4 fractions (F1 F2, F3 and F4) with different colors have been derived. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that MYP was the pigments with the lowest polarity among the others. Although the chemical structure of *Monascus* pigments consists of all of them of β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments. The presence of $-NH_3$ group can increase the polarity of red pigments.




Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.

Figure 2 shows the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in the orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin,




Figure 2. Scanning results of separated *Monascus* pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




Figure 3. LC-MS analysis results: Total Ion Chromatogram of MYP (A) and representatives of the chromatogram of major MYP monascin (B) and yellow II (C).

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

xanthomonasin B, monascopyridine B and Yellow II. Monascin and Yellow II were the major pigment compounds in the MYP with the relative amount of 59.81% and 27.80%. The MYP composition depends on the pigment composition in *Monascus* Fermented Durian Seed (MFDS), extraction and separation processes.

Pigments in MFDS produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but it is still unclear and controversy. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment rubropunctatin biosynthesis, it consist of the β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the *Monascus* pigment composition. The ethanolic extract contains higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated MYP from the red and orange pigments.

Monascin, the major MYP compound in this research, is attractive for its strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and showed antiatherogenic activities, and also improves memory and learning ability. On the other hand, there is no report on the yellow II bioactivity.

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in MYP.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [grant number 200X/WM01.5/N/2019].

References

- Agboyibor, C., Kong, W.B., Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447. <https://doi.org/10.1016/j.bcab.2018.09.012>
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565. <https://doi.org/10.1021/jf040199p>
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98(8), 1554-1560. <https://doi.org/10.1016/j.biortech.2006.06.005>
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136. <https://doi.org/10.1007/s11274-016-2082-8>
- Feng, Y., Shao, Y. and Chen, F. (2012). *Monascus* pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440. <https://doi.org/10.1007/s00253-012-4504-3>
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518. <https://doi.org/10.1021/jf1045987>
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264(1), 117–124. <https://doi.org/10.1111/j.1574-6968.2006.00451.x>
- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology*, 49(10), 2609–2617. <https://doi.org/10.1016/j.fct.2011.07.005>
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18(Part A), 387-399. <https://doi.org/10.1016/j.jff.2015.08.002>

- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143. <https://doi.org/10.3389/fmicb.2018.03143>
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390. <https://doi.org/10.1007/s11947-009-0233-8>
- Pattanagul, P., Pinthong, R., Phianmongkhon, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.
- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19(3), 941-945.
- Srianta, I., Ristiarini, S., Nugerahani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. *International Food Research Journal*, 21(1), 1-12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatalysis and Agricultural Biotechnology*, 7, 181-186. <https://doi.org/10.1016/j.bcab.2016.05.011>
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from *Monascus* fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *Journal of Agricultural and Food Chemistry*, 53, 1949-1954. <https://doi.org/10.1021/jf048310e>
- Vidyalakshmi, R., Paranthaman, R., Muruges, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. *Global Journal of Biotechnology and Biochemistry*, 4, 84–87.
- Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. *Marine Drugs*, 11(10), 3902-3925. <https://doi.org/10.3390/md11103902>

10

**Bukti konfirmasi submit permintaan revisi galley proof
artikel pertama dan artikel yang di-resubmit**

26 Maret 2020

Separation and analysis of *Monascus* yellow pigment produced on durian seed substrate

*Srianta, I., Nugerahani, I. and Ristiarini, S.

Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala Catholic University
Surabaya, Jalan Dinoyo 42-44 Surabaya, Indonesia 60265

Article history:

Received: 15 January 2020

Received in revised form: 2 March 2020

Accepted: 5 March 2020

Available Online:

Keywords:

Monascus,
Yellow pigment,
Durian seed,
Separation,
Analysis

DOI:

dfid

Abstract

This research was aimed to separate and analyse *Monascus* yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate: ethanol: water = 90: 25: 4. The separated MYP were then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has a maximum absorption at 386 nm. Out of five MYP compounds detected by using LC-MS/MS, two of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in MYP.

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through the polyketide pathway. It has been used as a natural food colourant, food supplement, and in traditional medicine. *Monascus* pigments consist of many pigment compounds which are categorized into three groups i.e. red, orange and yellow pigments. A total of six well-known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srianta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, thirty-nine new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid-state fermentation substrate in *Monascus* pigments production. Non-rice substrates have been developed in the *Monascus* pigment e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srianta *et al.*, 2012; Srianta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of *Monascus* yellow pigments is still limited. MYP is not only a potential natural colorant for biscuit, mayonnaise, wheat noodle, etc. (Srianta *et al.*, 2014) but also as functional food ingredient and in medicine,

because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), antiobesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) show that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in *Monascus* pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various *Monascus* pigments because of its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analyse MYP produced on durian seed substrate. This is the first

*Corresponding author.

Email: srianta_wm@yahoo.com

investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid-state fermentation on durian seed substrate. The culture was maintained periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya (Indonesia). The durian seed is a by-product of the food industry and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid-state fermentation with durian seed substrate

Pigments production was performed through solid-state fermentation on durian seed substrate according to our previous research (Srianta et al., 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 mins and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hrs, ground and sieved.

2.3 Monascus pigment separation

The *Monascus* pigment was extracted with soxhletation method in ethanolic solvent at a ratio of 1:50 for 2 hrs. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre eluted with ethyl acetate: ethanol: water = 90:25:4. The extract was put into the column and eluted with ethyl acetate: ethanol: water = 90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate: ethanol: water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through a 0.2 μ m PTFE filter membrane, put on to the autosampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with

an isocratic program of mobile phase consists of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and discussion

In the separation of the pigments by using column chromatography, 4 fractions (F1 F2, F3 and F4) with different colors have been derived. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that MYP was the pigments with the lowest polarity among the others. Although the chemical structure of *Monascus* pigments consists of **all of them of** β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments. The presence of $-NH_3$ group can increase the polarity of red pigments.




Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.

Figure 2 shows the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in the orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin,




Figure 2. Scanning results of separated *Monascus* pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




Figure 3. LC-MS analysis results: Total Ion Chromatogram of MYP (A) and representatives of the chromatogram of major MYP monascin (B) and yellow II (C).

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

xanthomonasin B, monascopyridine B and Yellow II. Monascin and Yellow II were the major pigment compounds in the MYP with the relative amount of 59.81% and 27.80%. The MYP composition depends on the pigment composition in *Monascus* Fermented Durian Seed (MFDS), extraction and separation processes.

Pigments in MFDS produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but it is still unclear and controversy. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment rubropunctatin biosynthesis, it consist of the β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the *Monascus* pigment composition. The ethanolic extract contains higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated MYP from the red and orange pigments.

Monascin, the major MYP compound in this research, is attractive for its strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and showed antiatherogenic activities, and also improves memory and learning ability. On the other hand, there is no report on the yellow II bioactivity.

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in MYP.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [grant number 200X/WM01.5/N/2019].

References

- Agboyibor, C., Kong, W.B., Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447. <https://doi.org/10.1016/j.bcab.2018.09.012>
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565. <https://doi.org/10.1021/jf040199p>
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98(8), 1554-1560. <https://doi.org/10.1016/j.biortech.2006.06.005>
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136. <https://doi.org/10.1007/s11274-016-2082-8>
- Feng, Y., Shao, Y. and Chen, F. (2012). *Monascus* pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440. <https://doi.org/10.1007/s00253-012-4504-3>
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518. <https://doi.org/10.1021/jf1045987>
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264(1), 117–124. <https://doi.org/10.1111/j.1574-6968.2006.00451.x>
- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology*, 49(10), 2609–2617. <https://doi.org/10.1016/j.fct.2011.07.005>
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18(Part A), 387-399. <https://doi.org/10.1016/j.jff.2015.08.002>

- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143. <https://doi.org/10.3389/fmicb.2018.03143>
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390. <https://doi.org/10.1007/s11947-009-0233-8>
- Pattanagul, P., Pinthong, R., Phianmongkhon, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.
- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19(3), 941-945.
- Srianta, I., Ristiarini, S., Nugerahani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. *International Food Research Journal*, 21(1), 1-12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatalysis and Agricultural Biotechnology*, 7, 181-186. <https://doi.org/10.1016/j.bcab.2016.05.011>
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from *Monascus* fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *Journal of Agricultural and Food Chemistry*, 53, 1949-1954. <https://doi.org/10.1021/jf048310e>
- Vidyalakshmi, R., Paranthaman, R., Muruges, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. *Global Journal of Biotechnology and Biochemistry*, 4, 84–87.
- Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. *Marine Drugs*, 11(10), 3902-3925. <https://doi.org/10.3390/md11103902>

Re: FR-2020-020 - Article Production

Dari: ignatius srianta (srianta_wm@yahoo.com)

Kepada: foodresearch.my@outlook.com

Tanggal: Kamis, 26 Maret 2020 pukul 21.41 GMT+7

Dear Dr. Vivian
Please kindly find the attached file for correction of FR-2020-020. Thank you
Best regards
Srianta

Pada Kamis, 26 Maret 2020 11.47.45 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Please refer to the attachment for the galley proof of your manuscript FR-2020-020 entitled 'Title' . Please check the content of the galley proof. If there are any mistakes, please comment and highlight in the PDF itself and revert to us within two (2) days of receipt. Once we have finalized the PDF version, your manuscript will be published online for early viewing.

Please see the attachment for the invoice INV20060. We hope that you can make the payment as soon as possible before 22 July 2020 for us to complete the publication of your manuscript. The manuscript information e.g. volume, issue, page numbers and DOI, will be provided once we have received the payment.

Thanks & Regards,
Vivian New
Editor
Food Research

From: ignatius srianta <srianta_wm@yahoo.com>
Sent: Tuesday, 24 March, 2020 2:18 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian
Please kindly the attached file of corrected FR-2020-020. Thank you
Best regards

Srianta

Pada Senin, 23 Maret 2020 16.25.24 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Manuscript ID: FR-2020-020

Manuscript Title: Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,

Vivian New

Editor

Food Research

From: srianta_wm@yahoo.com <srianta_wm@yahoo.com>

Sent: Monday, 2 March, 2020 9:00 AM

To: Food Research <foodresearch.my@outlook.com>

Subject: Re: FR-2020-020 (Revised)

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards

Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research



FR-2020-020_correction.pdf
575kB

11

**Bukti konfirmasi respon atas permintaan revisi galley
proof artikel pertama**

27 Maret 2020

Re: FR-2020-020 - Article Production

Dari: Food Research (foodresearch.my@outlook.com)

Kepada: srianta_wm@yahoo.com

Tanggal: Jumat, 27 Maret 2020 pukul 09.42 GMT+7

Dear Prof Srianta,

Please refer to the attachment for the edited galley proof. Please note that colourant is the right spelling as your article is using the English (UK), not English (US). Colorant is the spelling for English (US).

If the galley proof is alright, please approve the galley proof.

Thanks & Regards,

Vivian New

Editor

Food Research

From: ignatius srianta <srianta_wm@yahoo.com>

Sent: Thursday, 26 March, 2020 10:41 PM

To: Food Research <foodresearch.my@outlook.com>

Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian

Please kindly find the attached file for correction of FR-2020-020. Thank you

Best regards

Srianta

Pada Kamis, 26 Maret 2020 11.47.45 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Please refer to the attachment for the galley proof of your manuscript FR-2020-020 entitled 'Title'. Please check the content of the galley proof. If there are any mistakes, please comment and highlight in the PDF itself and revert to us within two (2) days of receipt. Once we have finalized the PDF version, your manuscript will be published online for early viewing.

Please see the attachment for the invoice INV20060. We hope that you can make the payment as soon as possible before 22 July 2020 for us to complete the publication of your manuscript. The manuscript information e.g. volume, issue, page numbers and DOI, will be provided once we have received the payment.

Thanks & Regards,
Vivian New
Editor
Food Research

From: ignatius srianta <srianta_wm@yahoo.com>
Sent: Tuesday, 24 March, 2020 2:18 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian
Please kindly the attached file of corrected FR-2020-020. Thank you
Best regards
Srianta

Pada Senin, 23 Maret 2020 16.25.24 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Manuscript ID: FR-2020-020

Manuscript Title: Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,
Vivian New
Editor
Food Research

From: srianta_wm@yahoo.com <srianta_wm@yahoo.com>
Sent: Monday, 2 March, 2020 9:00 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 (Revised)

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards
Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research



FR-2020-020.pdf
565.4kB

12

**Bukti konfirmasi submit permintaan revisi galley proof
artikel kedua dan artikel yang di-resubmit**

27 Maret 2020

Re: FR-2020-020 - Article Production

Dari: ignatius srianta (srianta_wm@yahoo.com)

Kepada: foodresearch.my@outlook.com

Tanggal: Jumat, 27 Maret 2020 pukul 10.52 GMT+7

Dear Dr. Vivian
Please one more correction, thank you
Best regards
Srianta

Pada Jumat, 27 Maret 2020 09.42.09 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Please refer to the attachment for the edited galley proof. Please note that colourant is the right spelling as your article is using the English (UK), not English (US). Colorant is the spelling for English (US).
If the galley proof is alright, please approve the galley proof.

Thanks & Regards,
Vivian New
Editor
Food Research

From: ignatius srianta <srianta_wm@yahoo.com>
Sent: Thursday, 26 March, 2020 10:41 PM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian
Please kindly find the attached file for correction of FR-2020-020. Thank you
Best regards
Srianta

Pada Kamis, 26 Maret 2020 11.47.45 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Sriantha,

Please refer to the attachment for the galley proof of your manuscript FR-2020-020 entitled 'Title' . Please check the content of the galley proof. If there are any mistakes, please comment and highlight in the PDF itself and revert to us within two (2) days of receipt. Once we have finalized the PDF version, your manuscript will be published online for early viewing.

Please see the attachment for the invoice INV20060. We hope that you can make the payment as soon as possible before 22 July 2020 for us to complete the publication of your manuscript. The manuscript information e.g. volume, issue, page numbers and DOI, will be provided once we have received the payment.

Thanks & Regards,
Vivian New
Editor
Food Research

From: ignatius sriantha <sriantha_wm@yahoo.com>
Sent: Tuesday, 24 March, 2020 2:18 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian
Please kindly the attached file of corrected FR-2020-020. Thank you
Best regards
Sriantha

Pada Senin, 23 Maret 2020 16.25.24 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Sriantha,

Manuscript ID: FR-2020-020

Manuscript Title: Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,
Vivian New
Editor
Food Research

From: srianta_wm@yahoo.com <srianta_wm@yahoo.com>
Sent: Monday, 2 March, 2020 9:00 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 (Revised)

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards
Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research



FR-2020-020_correction2.pdf
568.9kB

Separation and analysis of *Monascus* yellow pigment produced on durian seed substrate

*Srianta, I., Nugerahani, I. and Ristiarini, S.

Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala Catholic University
Surabaya, Jalan Dinoyo 42-44 Surabaya, Indonesia 60265

Article history:

Received: 15 January 2020

Received in revised form: 2 March 2020

Accepted: 5 March 2020

Available Online:

Keywords:

Monascus,
Yellow pigment,
Durian seed,
Separation,
Analysis

DOI:

dfid

Abstract

This research was aimed to separate and analyse *Monascus* yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate: ethanol: water = 90: 25: 4. The separated MYP were then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has a maximum absorption at 386 nm. Out of five MYP compounds detected by using LC-MS/MS, two of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in MYP.

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through the polyketide pathway. It has been used as a natural food colourant, food supplement, and in traditional medicine. *Monascus* pigments consist of many pigment compounds which are categorized into three groups i.e. red, orange and yellow pigments. A total of six well-known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srianta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, thirty-nine new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid-state fermentation substrate in *Monascus* pigments production. Non-rice substrates have been developed in the *Monascus* pigment production e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srianta *et al.*, 2012; Srianta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of *Monascus* yellow pigments is still limited. MYP is not only a potential natural colorant for biscuit, mayonnaise, wheat noodle, etc. (Srianta *et al.*, 2014) but also as functional food ingredient and in medicine,

because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), antiobesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) reported that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in *Monascus* pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various *Monascus* pigments because of its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analyse MYP produced on durian seed substrate. This is the first

*Corresponding author.

Email: srianta_wm@yahoo.com

investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid-state fermentation on durian seed substrate. The culture was maintained periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya (Indonesia). The durian seed is a by-product of the food industry and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid-state fermentation with durian seed substrate

Pigments production was performed through solid-state fermentation on durian seed substrate according to our previous research (Srianta et al., 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 mins and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hrs, ground and sieved.

2.3 Monascus pigment separation

The *Monascus* pigment was extracted with soxhletation method in ethanolic solvent at a ratio of 1:50 for 2 hrs. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre eluted with ethyl acetate: ethanol: water = 90:25:4. The extract was put into the column and eluted with ethyl acetate: ethanol: water = 90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate: ethanol: water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through a 0.2 µm PTFE filter membrane, put on to the autosampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with

an isocratic program of mobile phase consists of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and discussion

In the separation of the pigments by using column chromatography, 4 fractions (F1 F2, F3 and F4) with different colors have been derived. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that MYP was the pigments with the lowest polarity among the others. Although the chemical structure of *Monascus* pigments consists of β-ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments. The presence of -NH₃ group can increase the polarity of red pigments.




Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.

Figure 2 shows the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in the orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and Yellow II.




Figure 2. Scanning results of separated *Monascus* pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




Figure 3. LC-MS analysis results: Total Ion Chromatogram of MYP (A) and representatives of the chromatogram of major MYP monascin (B) and yellow II (C).

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

Monascin and Yellow II were the major pigment compounds in the MYP with the relative amount of 59.81% and 27.80%, respectively. The MYP composition depends on the pigment composition in *Monascus* Fermented Durian Seed (MFDS), extraction and separation processes.

Pigments in MFDS produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but it is still unclear and controversial. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment rubropunctatin biosynthesis, it consists of the β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the *Monascus* pigment composition. The ethanolic extract contains higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated MYP from the red and orange pigments.

Monascin, the major MYP compound in this research, is attractive for its strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and showed antiatherogenic activities, and also improves memory and learning ability. On the other hand, there is no report on the yellow II bioactivity.

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in MYP.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [contract number 200X/WM01.5/

N/2019].

References

- Agboyibor, C., Kong, W.B., Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447. <https://doi.org/10.1016/j.bcab.2018.09.012>
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565. <https://doi.org/10.1021/jf040199p>
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98(8), 1554-1560. <https://doi.org/10.1016/j.biortech.2006.06.005>
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136. <https://doi.org/10.1007/s11274-016-2082-8>
- Feng, Y., Shao, Y. and Chen, F. (2012). *Monascus* pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440. <https://doi.org/10.1007/s00253-012-4504-3>
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518. <https://doi.org/10.1021/jf1045987>
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264(1), 117–124. <https://doi.org/10.1111/j.1574-6968.2006.00451.x>
- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology*, 49(10), 2609–2617. <https://doi.org/10.1016/j.fct.2011.07.005>
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18(Part A), 387-399.

- <https://doi.org/10.1016/j.jff.2015.08.002>
- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143. <https://doi.org/10.3389/fmicb.2018.03143>
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390. <https://doi.org/10.1007/s11947-009-0233-8>
- Pattanagul, P., Pinthong, R., Phianmongkhol, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.
- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19(3), 941-945.
- Srianta, I., Ristiarini, S., Nugerahani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. *International Food Research Journal*, 21(1), 1-12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatalysis and Agricultural Biotechnology*, 7, 181-186. <https://doi.org/10.1016/j.bcab.2016.05.011>
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from *Monascus* fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *Journal of Agricultural and Food Chemistry*, 53, 1949-1954. <https://doi.org/10.1021/jf048310e>
- Vidyalakshmi, R., Paranthaman, R., Muruges, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. *Global Journal of Biotechnology and Biochemistry*, 4, 84–87.
- Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. *Marine Drugs*, 11(10), 3902-3925. <https://doi.org/10.3390/md11103902>

13

Bukti konfirmasi persetujuan galley proof artikel

27 Maret 2020

Re: FR-2020-020 - Article Published

Dari: Food Research (foodresearch.my@outlook.com)

Kepada: srianta_wm@yahoo.com

Tanggal: Jumat, 27 Maret 2020 pukul 20.56 GMT+7

Dear Prof Srianta,

Your manuscript is currently available online and in press on our website <https://www.myfoodresearch.com>.

Please note that the version in press is the 'Corrected Proof' version. The manuscript information e.g. volume, issue, page number and DOI, will be provided once we have received the payment. We will then inform you of the update of your manuscript details.

Thank you for your fine contribution. We hope that you continue to submit other articles to the Journal.

Thanks & Regards,

Vivian New

Editor

Food Research

From: ignatius srianta <srianta_wm@yahoo.com>
Sent: Friday, 27 March, 2020 11:52 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian
Please one more correction, thank you
Best regards
Srianta

Pada Jumat, 27 Maret 2020 09.42.09 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Please refer to the attachment for the edited galley proof. Please note that colourant is the right spelling as your article is using the English (UK), not English (US). Colorant is the spelling for English (US).

If the galley proof is alright, please approve the galley proof.

Thanks & Regards,
Vivian New
Editor
Food Research

From: ignatius srianta <srianta_wm@yahoo.com>
Sent: Thursday, 26 March, 2020 10:41 PM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian
Please kindly find the attached file for correction of FR-2020-020. Thank you
Best regards
Srianta

Pada Kamis, 26 Maret 2020 11.47.45 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Please refer to the attachment for the galley proof of your manuscript FR-2020-020 entitled 'Title' . Please check the content of the galley proof. If there are any mistakes, please comment and highlight in the PDF itself and revert to us within two (2) days of receipt. Once we have finalized the PDF version, your manuscript will be published online for early viewing.

Please see the attachment for the invoice INV20060. We hope that you can make the payment as soon as possible before 22 July 2020 for us to complete the publication of your manuscript. The manuscript information e.g. volume, issue, page numbers and DOI, will be provided once we have received the payment.

Thanks & Regards,
Vivian New
Editor
Food Research

From: ignatius srianta <srianta_wm@yahoo.com>
Sent: Tuesday, 24 March, 2020 2:18 AM

To: Food Research <foodresearch.my@outlook.com>

Subject: Re: FR-2020-020 - Article Production

Dear Dr. Vivian
Please kindly the attached file of corrected FR-2020-020. Thank you
Best regards
Srianta

Pada Senin, 23 Maret 2020 16.25.24 GMT+7, Food Research <foodresearch.my@outlook.com> menulis:

Dear Prof Srianta,

Manuscript ID: FR-2020-020

Manuscript Title: Separation and analysis of Monascus yellow pigment produced on durian seed substrate

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,
Vivian New
Editor
Food Research

From: srianta_wm@yahoo.com <srianta_wm@yahoo.com>

Sent: Monday, 2 March, 2020 9:00 AM

To: Food Research <foodresearch.my@outlook.com>

Subject: Re: FR-2020-020 (Revised)

Dear Prof. Son Radu

Please kindly find the attached files of FR-2020-020_revised version and response to reviewers comments. Thank you for your kind attention

Best regards
Srianta

On Feb 7, 2020 23:55, Food Research <foodresearch.my@outlook.com> wrote:

Dear Srianta,

Manuscript FR-2020-020 entitled " Separation and analysis of Monascus yellow pigment produced on durian seed substrate " which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted before or by 8th March 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,
Professor Dr. Son Radu
foodresearch.my@outlook.com

Chief Editor, Food Research

14

**Bukti konfirmasi artikel published online dan artikel yang
diterbitkan**

1 April 2020

FR-2020-020 - Article Updated

Dari: Food Research (foodresearch.my@outlook.com)

Kepada: srianta_wm@yahoo.com

Tanggal: Rabu, 1 April 2020 pukul 14.34 GMT+7

Dear Prof Srianta,

Thank you for the payment.

Kindly be informed that your manuscript has been assigned to Food Research 2020, Vol. 4, Issue 4 (August). Your manuscript is currently available online and in press on our website <https://www.myfoodresearch.com>. Alternatively, you can download a copy of the manuscript by clicking on the following link:

[https://doi.org/10.26656/fr.2017.4\(4\).020](https://doi.org/10.26656/fr.2017.4(4).020)

We encourage you to share your published work with your colleagues. Thank you for your fine contribution. We hope that you continue to submit other articles to the Journal.

Thanks & Regards,

Dr. Vivian New

Editor

Food Research

Separation and analysis of *Monascus* yellow pigment produced on durian seed substrate

*Srianta, I., Nugerahani, I. and Ristiarini, S.

Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala Catholic University
Surabaya, Jalan Dinoyo 42-44 Surabaya, Indonesia 60265

Article history:

Received: 15 January 2020

Received in revised form: 2 March 2020

Accepted: 5 March 2020

Available Online: 27 March 2020

Keywords:

Monascus,
Yellow pigment,
Durian seed,
Separation,
Analysis

Abstract

This research was aimed to separate and analyse *Monascus* yellow pigment (MYP) with chromatography. MYP was in a mixture with red and orange pigments. MYP separation was performed in a silica gel column with eluent of ethyl acetate: ethanol: water = 90: 25: 4. The separated MYP were then subjected to analysis by using spectrophotometer, TLC, LC-PDA and LC-MS/MS. The results revealed that the separated MYP has a maximum absorption at 386 nm. Out of five MYP compounds detected by using LC-MS/MS, two of them were well known yellow pigment compounds, monascin and ankaflavin. Monascin and yellow II were the major pigment compounds in MYP.

DOI:

[https://doi.org/10.26656/fr.2017.4\(4\).020](https://doi.org/10.26656/fr.2017.4(4).020)

1. Introduction

Monascus pigments are secondary metabolites of *Monascus* fungi biosynthesized through the polyketide pathway. It has been used as a natural food colourant, food supplement, and in traditional medicine. *Monascus* pigments consist of many pigment compounds which are categorized into three groups i.e. red, orange and yellow pigments. A total of six well-known pigment compounds are monascorubramine and rubropunctamine (red pigments), monascorubrin and rubropunctatin (orange pigments), monascin and ankaflavin (yellow pigments) (Feng *et al.*, 2012; Srianta *et al.*, 2014; Agboyibor *et al.*, 2018). Until 2011, thirty-nine new pigment compounds have been identified (Feng *et al.*, 2012).

Rice is commonly used as solid-state fermentation substrate in *Monascus* pigments production. Non-rice substrates have been developed in the *Monascus* pigment production e.g. corn, adlay, sorghum, dioscorea, jackfruit seed and durian seed (Pattanagul *et al.*, 2007; Babitha *et al.*, 2007; Hsu *et al.*, 2011; Srianta *et al.*, 2012; Srianta *et al.*, 2016). Our previous studies revealed that *Monascus purpureus* produced red, orange and yellow pigments. Exploration of *Monascus* yellow pigments is still limited. MYP is not only a potential natural colourant for biscuit, mayonnaise, wheat noodle, etc. (Srianta *et al.*, 2014) but also as functional food ingredient and in medicine,

because MYP has various positive health effects i.e. antioxidant, antidiabetes and anticancer (Hsu *et al.*, 2011; Chen and Wu, 2016). Monascin and ankaflavin have antitumor activity (Akihisa *et al.*, 2005; Su *et al.*, 2005), antiobesity activity, anti-atherosclerosis effect (Wang *et al.*, 2013; Liu *et al.*, 2018), improve memory and learning ability (Lee *et al.*, 2015). Chen and Wu (2016) reported that MYPs have promising potential for application in food and pharmaceutical industries based on their bioactivities.

Silica gel has been applied in *Monascus* pigments mixture separation through thin layer chromatography and column chromatography systems with various eluents i.e. chloroform, methanol, benzene, acetone and petroleum ether (Kim *et al.*, 2006; Vidyalakshmi *et al.*, 2009; Hsu *et al.*, 2011; Lee *et al.*, 2011). In practice, ethyl acetate, ethanol and water are often more preferred for food and pharmaceutical processing compared to other solvents due to their safety and affordability. HPLC is utilized to analyze various *Monascus* pigments because of its high sensitivity and multiple detection systems including ultraviolet, fluorescent, photodiode array and mass spectrometer detectors (Feng *et al.*, 2012).

This research was aimed to separate and analyse MYP produced on durian seed substrate. This is the first

*Corresponding author.

Email: srianta_wm@yahoo.com

investigation in MYP separation using ethyl acetate, ethanol and water.

2. Materials and methods

2.1 Microorganism, substrate and chemicals

Monascus purpureus M9 culture was used in the solid-state fermentation on durian seed substrate. The culture was maintained periodically on Potato Dextrose Agar. Durian seed was obtained from a durian processing unit in Surabaya (Indonesia). The durian seed is a by-product of the food industry and was used as the SSF substrate. All chemicals were chromatographic and analytical grades.

2.2 Pigments production through solid-state fermentation with durian seed substrate

Pigments production was performed through solid-state fermentation on durian seed substrate according to our previous research (Srianta et al., 2012). After peeling, cutting, soaking, sterilization at 121°C for 10 mins and cooling, the durian seed substrate was inoculated with *Monascus purpureus* starter culture, and then incubated at 30°C for 14 days. The fermented matter was dried at 45°C for 24 hrs, ground and sieved.

2.3 Monascus pigment separation

The *Monascus* pigment was extracted with soxhlet method in ethanolic solvent at a ratio of 1:50 for 2 hrs. The extract was then evaporated in a rotary evaporator. The evaporated extract was subjected to separation process. MYP separation was performed in a column chromatography (10 mm x 500 mm) with silica gel 60. The column was packed with silica gel by slurry method, then pre eluted with ethyl acetate: ethanol: water = 90:25:4. The extract was put into the column and eluted with ethyl acetate: ethanol: water = 90:25:4. The obtained fractions were subjected to analysis by using TLC and spectrophotometer. The MYP fraction was analyzed the pigment composition by using LC-MS/MS.

2.4 Pigment analysis

The separated pigments were subjected to the TLC analysis according to Nimnoi and Lumyong (2011) with modification, on silica gel F60 plate, with ethyl acetate: ethanol: water at 90:25:4. The separated MYP was scanned in a range of wavelength 380-600 nm with UV-Vis spectrophotometer (Shimadzu UV-1900, Japan). Pigment composition analysis was conducted by using LC-MS/MS (Shimadzu, Kyoto, Japan) with Q1 scan positive mode and SIM positive mode. The extract was filtered through a 0.2 µm PTFE filter membrane, put on to the autosampler. Pigments were separated on a Cosmosil 5 C18-MS-II (4.6 i.d. x 150 mm) column, with

an isocratic program of mobile phase consists of acetonitrile containing 0.1% formic acid and water containing 0.1% formic acid (60:40 vol/vol), flow rate of 0.5 mL/min; oven temperature of 40°C, run time for 30 min. Pigment compounds were detected by MS/MS using electro-spray ionization in the positive ion mode, scan range: m/z 100 to 1000.

3. Results and discussion

In the separation of the pigments by using column chromatography, 4 fractions (F1 F2, F3 and F4) with different colours have been derived. The first fraction was MYP target, while other fractions were orange and red pigments. The MYP has the highest retention factor, followed by F2, F3 and F4. Those fractions and pigments profile on TLC shown in Figure 1. Those profiles describe that MYP was the pigments with the lowest polarity among the others. Although the chemical structure of *Monascus* pigments consists of β -ketoacid and chromophore, MYP compounds contain less polar groups than those of red pigments. The presence of $-NH_3$ group can increase the polarity of red pigments.




Figure 1. Picture of 4 separated fractions of *Monascus* pigments (A) and their TLC chromatogram (B). F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.

Figure 2 shows the fractions scanning by using spectrophotometer at 380-700 nm. F1, the MYP, has a λ max at 386 nm that was in yellow spectrum. F3 and F4 have 2 peaks in the orange and red spectrum. Those reflect that contain a mixture of orange and red pigments. This is the first investigation on the pigment compounds composition of MYP by using the LC-MS/MS. The LC-MS chromatograms and pigment composition of the extracts are presented in Figure 3 and Table 1, respectively. Five MYP compounds were detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and Yellow II.




Figure 2. Scanning results of separated *Monascus* pigment fractions by using spectrophotometer. F1: Fraction 1; F2: Fraction 2; F3: Fraction 3; F4: Fraction 4.




Figure 3. LC-MS analysis results: Total Ion Chromatogram of MYP (A) and representatives of the chromatogram of major MYP monascin (B) and yellow II (C).

Table 1. MYP composition

MYP compound	Content (%)
Monascin	59.81
Ankaflavin	6.32
XanthomonasinA	n.d.
XanthomonasinB	0.67
Monascopyridin A	n.d.
Monascopyridin B	5.39
Yellow II	27.80

n.d. = not detected

Monascin and Yellow II were the major pigment compounds in the MYP with the relative amount of 59.81% and 27.80%, respectively. The MYP composition depends on the pigment composition in *Monascus* Fermented Durian Seed (MFDS), extraction and separation processes.

Pigments in MFDS produced through a complex polyketide biosynthesis pathway in the *Monascus* fungi, but it is still unclear and controversial. Some researchers proposed that only orange pigments (rubropunctatin and monascorubramine) were biosynthetic, and the others were transformed from them chemically. In orange pigment rubropunctatin biosynthesis, it consists of the β -ketide pathway to form acetate and malonate and another pathway to form n-hexanoyl acetyl residue. Yellow pigments can be transformed from orange pigments via oxidation. Our previous research revealed that solvent type affected the *Monascus* pigment composition. The ethanolic extract contains higher MYP than those in the mixture solvent of ethanol and water. This research used ethanol as the extraction solvent. In the separation process, the chromatographic system used has successfully separated MYP from the red and orange pigments.

Monascin, the major MYP compound in this research, is attractive for its strong bioactivities such as anticancer, antitumor, antidiabetic, antioxidative stress, antiinflammatory, antiobesity and showed antiatherogenic activities, and also improves memory and learning ability. On the other hand, there is no report on the yellow II bioactivity.

4. Conclusion

The MYP has been successfully separated in a column chromatography packed with silica gel 60 and eluent of ethyl acetate:ethanol:water=90:25:4. Five MYP compounds have been detected by using LC-MS/MS i.e. monascin, ankaflavin, xanthomonasin B, monascopyridine B and yellow II. Monascin and yellow II were the major pigment compounds in MYP.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

This work was supported by Direktorat Riset dan Pengabdian Masyarakat, Direktorat Jenderal Penguatan Riset dan Pengembangan, Kementerian Riset, Teknologi, dan Pendidikan Tinggi, Republik Indonesia through the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019 [contract number 200X/WM01.5/

N/2019].

References

- Agboyibor, C., Kong, W.B., Chen, D., Zhang, A.M. and Niu, S.Q. (2018). *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatalysis and Agricultural Biotechnology*, 16, 433-447. <https://doi.org/10.1016/j.bcab.2018.09.012>
- Akihisa, T., Tokuda, H., Yasukawa, K., Ukiya, M., Kiyota, A., Sakamoto, N., Suzuki, T., Tanabe, N. and Nishino, H. (2005). Azaphilones, furanoisophthalides, and amino acids from the extracts of *Monascus pilosus*-fermented rice (red-mold rice) and their chemopreventive effects. *Journal of Agricultural and Food Chemistry*, 53, 562–565. <https://doi.org/10.1021/jf040199p>
- Babitha, S., Soccol, C.R. and Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98(8), 1554-1560. <https://doi.org/10.1016/j.biortech.2006.06.005>
- Chen, G. and Wu, Z. (2016). Production and biological activities of yellow pigments from *Monascus* fungi. *World Journal of Microbiology and Biotechnology*, 32, 136. <https://doi.org/10.1007/s11274-016-2082-8>
- Feng, Y., Shao, Y. and Chen, F. (2012). *Monascus* pigments. *Applied Microbiology and Biotechnology*, 96, 1421–1440. <https://doi.org/10.1007/s00253-012-4504-3>
- Hsu, Y.W., Hsu, L.C., Liang, Y.H., Kuo, Y.H. and T.M. Pan. (2011). New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented dioscorea. *Journal of Agricultural and Food Chemistry*, 59, 4512–4518. <https://doi.org/10.1021/jf1045987>
- Kim, C., Jung, H., Kim, Y.O. and Shin, C.S. (2006). Antimicrobial activities of amino acid derivatives of *Monascus* pigments. *FEMS Microbiology Letters*, 264(1), 117–124. <https://doi.org/10.1111/j.1574-6968.2006.00451.x>
- Lee, B.H., Hsu, W.H., Liao, T.H. and Pan, T.M. (2011). The *Monascus* metabolite monascin against TNF- α -induced insulin resistance via suppressing PPAR- γ phosphorylation in C2C12 myotubes. *Food and Chemical Toxicology*, 49(10), 2609–2617. <https://doi.org/10.1016/j.fct.2011.07.005>
- Lee, C.L., Lin, P.Y., Hsu, Y.W. and Pan, T.M. (2015). *Monascus*-fermented monascin and ankaflavin improve the memory and learning ability in amyloid β -protein intracerebroventricular-infused rat via the suppression of Alzheimer's disease risk factors. *Journal of Functional Foods*, 18(Part A), 387-399.

- <https://doi.org/10.1016/j.jff.2015.08.002>
- Liu, L., Zhao, J., Huang, Y., Xin, Q. and Wang, Z. (2018). Diversifying of chemical structure of native *Monascus* pigments. *Frontiers in Microbiology*, 9, 3143. <https://doi.org/10.3389/fmicb.2018.03143>
- Nimnoi, P. and Lumyong S. (2011). Improving solid-state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food Bioprocess and Technology*, 4, 1384–1390. <https://doi.org/10.1007/s11947-009-0233-8>
- Pattanagul, P., Pinthong, R., Phianmongkhol, A. and Leksawasdi, N. (2007) Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.
- Srianta, I., Hendrawan, B., Kusumawati, N. and Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19(3), 941-945.
- Srianta, I., Ristiarini, S., Nugerahani, I., Sen, S.K., Zhang, B.B., Xu, G.R. and Blanc, P.J. (2014). Recent research and development of *Monascus* fermentation products. *International Food Research Journal*, 21(1), 1-12.
- Srianta, I., Zubaidah, E., Estiasih, T., Yamada, M. and Harijono (2016). Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatalysis and Agricultural Biotechnology*, 7, 181-186. <https://doi.org/10.1016/j.bcab.2016.05.011>
- Su, N.W., Lin, Y.L., Lee, M.H. and Ho, C.Y. (2005). Ankaflavin from *Monascus* fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *Journal of Agricultural and Food Chemistry*, 53, 1949-1954. <https://doi.org/10.1021/jf048310e>
- Vidyalakshmi, R., Paranthaman, R., Muruges, S. and Singaravadivel, K. (2009). Microbial bioconversion of rice broken to food grade pigments. *Global Journal of Biotechnology and Biochemistry*, 4, 84–87.
- Wang, L.C., Lung, T.Y., Kung, Y.H., Wang, J.J., Tsai, T.Y., Wei, B.L., Pan, T.M. and Lee, C.L. (2013). Enhanced anti-obesity activities of red mold dioscorea when fermented using deep ocean water as the culture water. *Marine Drugs*, 11(10), 3902-3925. <https://doi.org/10.3390/md11103902>