

Effect of Lactobacillus plantarum and Leuconostoc mesenteroides

by Ignatius Srianta

Submission date: 15-Oct-2023 02:58PM (UTC+0700)

Submission ID: 2195988082

File name: el_Effect_of_Lactobacillus_Sauerkraut_-_jurnal_Food_Research.pdf (481.46K)

Word count: 4633

Character count: 24699

Effect of *Lactobacillus plantarum* and *Leuconostoc mesenteroides* starter cultures in lower salt concentration fermentation on the sauerkraut quality^{1,*}Zubaidah, E., ¹Susanti, I., ¹Yuwono, S.S., ²Rahayu, A.P., ³Srianta, I. and ⁴Blanc, P.J.¹Department of Food Science and Technology, Faculty of Agricultural Technology, Brawijaya University, Jalan Veteran Malang, Indonesia 65145²Department of Agronomy, Faculty of Agriculture, Brawijaya University, Jalan Veteran Malang, Indonesia 65145³Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala Catholic University Surabaya, Jalan Dinoyo 42-44, Surabaya, Indonesia 60265⁴Université de Toulouse, INSA, LISBP, CNRS, UMR5504, INRA, UMR792, Ingénierie des Systèmes Biologiques et des Procédés, 135 Avenue de Rangueil, F-31077 Toulouse, France**Article history:**Received: 16 January 2020
Received in revised form: 1 February 2020
Accepted: 3 February 2020
Available Online: 10 March 2020**Keywords:**Antioxidant,
Lactobacillus plantarum,
Leuconostoc mesenteroides,
Sauerkraut**DOI:**[https://doi.org/10.26656/fr.2017.4\(4\).029](https://doi.org/10.26656/fr.2017.4(4).029)**Abstract**

This research was aimed to study the effect of *Lactobacillus plantarum* and *Leuconostoc mesenteroides* starter cultures application in lower salt concentration fermentation on sauerkraut quality. Fresh cut cabbage was fermented with different starter cultures (*L. plantarum*, *L. mesenteroides* or the combination) at different lower salt concentration (0.5% or 1%) at 28°C for 5 days. The obtained sauerkrauts were subjected to evaluation of the quality i.e. total lactic acid bacteria, pH, total acidity, total phenolic content and DPPH scavenging activity analysis. The sulfuraphane content analysis was performed by using LC-MS. The starter cultures increased total lactic acid bacteria, total acidity and decreased pH. *L. mesenteroides* resulted in the highest total phenolic content and the lowest IC₅₀ value. Sauerkraut with the addition of *L. mesenteroides* contains sulfuraphane higher 848.65 ng/g than that of control 776.47 ng/g. The results analysis of LC-MS also detected another compound, namely 2-phenethyl isothiocyanate, an antimicrobial compound. The sauerkraut is potential in functional food development with antiproliferative, anti-inflammatory, antioxidant, and anti-cancer activities.

1. Introduction

Lactic acid fermentation is expected to become an important role in the development of functional fermented vegetables. Sauerkraut, fermented cucumbers and kimchi are the most studied lactic acid fermented vegetables mainly due to their commercial importance (Swain *et al.*, 2014). Sauerkraut, means sour cabbage, is a fermentation product of cabbage through spontaneous lactic acid bacteria fermentation. In the fermentation, fresh cabbage is shredded and mixed with 2-2.5% salt to pull out water and nutrients from the cabbage, and the juice will become a substrate for the lactic acid bacteria growth (Johanningsmeier *et al.*, 2007). Several lactic acid bacteria play an important role in the fermentation process i.e. *L. mesenteroides*, *Lactobacillus cucumeris*, *L. plantarum* and *Lactobacillus pentoaceti* (Lu *et al.*, 2003; Plengvidhya *et al.*, 2007; Swain *et al.*, 2014). Addition of two types of bacteria to determine the

performance of bacteria that play an active role when making change to bioactive compounds, and to know the performance of those local bacteria when added to bacteria to cabbage that will be used by the lactic acid bacteria for their growth. Addition of a combination of bacteria to determine the acidity in sauerkraut, if the two types of bacteria added will be equally active and work in the fermentation process.

Concerning on the salt concentration in sauerkraut fermentation, consumers tend to prefer lower sodium foods. Moreover, the brine in the fermentation contains very high in nondegradable chloride ions and BOD. The ability to reduce the salt in sauerkraut fermentation would reduce the concentration of sodium chloride in the waste stream and the volume of brine formed (Johanningsmeier *et al.*, 2007). The salt has a function to draw water out of nutrients contained in cabbage that will be used by the substrate for the growth of lactic acid

*Corresponding author.

Email: elok@ub.ac.id; elzoeba@yahoo.com

bacteria. The addition of 2% salt can accelerate the fermentation process of sauerkraut but is less effective in inhibiting the growth of pathogenic microorganisms. However, the high salt concentration will kill the lactic acid bacteria. Therefore the addition of lactic acid bacteria culture can reduce the salt addition while the fermentation process still go well and increase bioactive compounds (Xiaoze *et al.*, 2019). Several researchers reported that starter cultures favored the low salt sauerkraut fermentation (Tolonen *et al.*, 2004; Wiander and Ryhanan, 2005).

On the other hand, Penas *et al.* (2010) reported that antioxidant activity increase during sauerkraut fermentation, lactic acid bacteria capable to increase the bioactive compounds i.e. phenolic and glucosinolate compounds. Phenolic compounds have the ability to increase antioxidant activity, furthermore can avoid degenerative diseases (Murray, 2009), while sulforaphane is an isothiocyanate derivative that has antiproliferative, anti-inflammatory, antioxidant, and anti-cancer activities. The ability to prevent cancer through DNA protection by modulating enzymes and inhibiting gene mutations (Romeo *et al.*, 2018). Antioxidant activity in sauerkraut can inhibit nitric oxide (NO) a factor that causes inflammation that is one of the responses in immune cells (Penas *et al.*, 2012).

This research aimed to study the effect of starter cultures application of *L. plantarum* and *L. mesenteroides* at lower salt fermentation on the sauerkraut quality.

2. Materials and methods

2.1 Materials

White cabbage (*Brassica olerace L. var*) used in this experiment was purchased from the local market. Cultures of *L. mesenteroides* FNCC 0023 and *L. plantarum* FNCC 0027 were obtained from Food and Nutrition Culture Collection, Gadjah Mada University. The cultures were maintained routinely on a MRSB medium.

2.2 Starter culture preparation

The starter cultures of *L. mesenteroides* and *L. plantarum* were prepared according to Penas *et al.*, (2012). A loopful of the culture was inoculated into 7 mL of MRSB medium and incubated at 37°C for 16 hrs. The culture suspension was harvested by centrifugation at 6000 rpm for 15 mins. The cells were washed with sterile distilled water, then put it into a 100 mL of sterile distilled water with cell density of 10⁶ CFU/mL.

2.3 Sauerkraut fermentation

The fresh cabbage was washed, cut, added with salt at different concentration (0.5% or 1%), inoculated with different starter culture (*L. plantarum*, *L. mesenteroides* or the combination) at 20% (v/w), then incubated at room temperature (28°C) for 5 days. Spontaneous fermentation with salt at 2% was used as a control. The obtained sauerkrauts were subjected to evaluation of the quality i.e. total lactic acid bacteria, pH, total acidity, total phenolic content and DPPH scavenging activity analysis. The sulforaphane content analysis was performed on the selected sauerkraut.

2.4 Sauerkraut quality evaluation

2.4.1 Total lactic acid bacteria

Total lactic acid bacteria were determined according to Penas *et al.* (2010). A total of 5 g of the sample was prepared aseptically to make a sample solution, then diluted with buffer peptone water into serial dilution. The diluted sample suspension was poured on MRS Agar and then incubated at 37°C for 48 hrs.

2.4.2 Total acidity and pH

Total acidity was measured according to Ranggana (1977) by using direct titration with NaOH solution of 0.1N and phenol-phtalein indicator. Total acidity was expressed as percentage of lactic acid. pH was measured by using pH meter (Manual pH meter Micro Bench TI 2100).

2.4.3 Total phenolic content

One gram of sample was extracted in 10 mL of methanol, centrifuged at 6000 rpm for 20 mins. 0.5 mL of supernatant was put into the test tube, added with 2.5 mL of 10% Folin Ciocalteu reagent and 2.5 mL of 7.5% Na₂CO₃, and incubated at room temperature for 90 mins. Absorbance was measured at 750 nm with spectrophotometer (Yang *et al.*, 2007). Gallic acid was used as a standard. The total phenolic content in sauerkraut was expressed as mg GAE/g.

2.4.4 DPPH scavenging activity

The extract was prepared with the same procedure of the total phenolic content analysis. The supernatant was diluted into 10, 20, 30, 40 and 50 ppm. 4 mL of sample was added into 1 mL of 0.2 mM diphenyl-1-picrylhydrazyl radical (DPPH). The mixture was incubated in the darkroom for 30 mins, then the absorbance was measured at 517 nm with a spectrophotometer (Molyneux, 2004). The DPPH scavenging activity was expressed as IC₅₀.

2.4.5 Sulforaphane content analysis

Sulforaphane was analyzed by using Liquid

Chromatography-Mass Spectrometry (LC-MS) according to Kim *et al.* (2017). Sample preparation was conducted according to Liang *et al.* (2006) with the following procedure: 5 g of the sample was extracted with methylene chloride, dissolved in acetonitrile and filtered through a 0.22 µm membrane. The extract was injected to an HPLC system (Agilent Seris 1200) connected to electrospray ionization (ESI) with API 400 Q TRAP Mass Spectrometry system. The LC operating conditions were reversed-phase C18 column in an oven set at 30°C, mobile phase of 20% acetonitrile in water then changed linearly to 60% acetonitrile with flow rate was 1 mL/min. The MS operating conditions were as follow: ESI positive ion ([M + H]⁺), ion spray voltage (5.5 kV), gas (20 psi), nebulisation gas (50 psi), heater gas (50 psi), high purity nitrogen (N₂), heater gas temperature (550°C), declustering potential (100 V), entrance potential (10 V), and spectrum range (m/z 100 -1000). Then the sulforaphane content calculation was done using a standard sulforaphane curve and expressed as µg/g.

2.5 Statistical analysis

The data were analyzed by analysis of variance (ANOVA), followed by the LSD test at $p < 0.05$.

3. Results and discussion

3.1 Lactic acid bacteria growth and activity

In sauerkraut fermentation, the lactic acid bacteria grow and transforms substrates into lactic acid and others. The lactic acid bacteria growth and activity are indicated from the differences data of total lactic acid bacteria, total acidity and pH before and after the fermentation presented in Table 1 and Table 2. The addition of 6 log CFU load starter cultures at initial fermentation increased 2 log cycles during the 5 days fermentation. This is related to the increase in total acidity and decreasing pH values during the sauerkraut fermentation. The pH decreased due to the lactic acid bacteria produce organic acids and release H⁺ ion cause an acidic atmosphere in the fermentation (Goh *et al.*, 2012).

There were significant ($p < 0.05$) starter culture differences in the total LAB at the final fermentation, with the combination starter cultures having the highest total LAB (Table 1). Combination of the cultures yielded higher growth than the single culture. The LAB growth demonstrated is thought to be due to the different contribution of both cultures, as reported by Plengvidhya *et al.* (2007) that *L. mesenteroides* dominate at the beginning of fermentation, then *L. plantarum* and *Lactobacillus brevis* continuing the fermentation. Addition of *L. mesenteroides* starter culture yielded higher growth than that of *L. plantarum*. This may indicate the different growth ability of the cultures to

Table 1. Total Lactic Acid Bacteria (LAB) total acidity before and after in sauerkraut fermentation at different starter cultures and salt concentrations

Culture	Salt (%)	Total of LAB	
		Day 0	Day 5
		10 ⁶ (CFU/mL)	10 ⁸ (CFU/mL)
<i>L. plantarum</i>	0.5	2.4±0.11	3.4±0.06 ^d
<i>L. mesenteroides</i>		2.7±0.10	2.4±0.10 ^c
<i>L. plantarum</i> + <i>L. mesenteroides</i>		2.3±0.14	5.5±0.12 ^b
<i>L. plantarum</i>	1	2.5±0.01	4.6±0.11 ^c
<i>L. mesenteroides</i>		2.3±0.14	3.1±0.09 ^d
<i>L. plantarum</i> + <i>L. mesenteroides</i>		2.6±0.23	6.6±0.10 ^a
Control (Spontaneous fermentation)	2	2.6±0.23	2.6 x 10 ⁷ ±0.23

The values are expressed as a mean ± SD (n = 3), and values in a column with the same letters are not significantly ($P > 0.05$) different.

Table 2. Total acidity and pH before and after in sauerkraut fermentation at different starter cultures and salt concentrations

Culture	Salt (%)	Total acidity (%)		pH	
		Day 0	Day 5	Day 0	Day 5
		<i>L. plantarum</i>	0.35±0.02	1.33±0.04 ^c	5.67±0.03
<i>L. mesenteroides</i>	0.5	0.37±0.03	1.31±0.01 ^c	5.75±0.12	3.79 ^a ±0.20
<i>L. plantarum</i> + <i>L. mesenteroides</i>		0.34±0.01	2.05±0.05 ^a	5.72±0.05	2.60 ^b ±0.13
<i>L. plantarum</i>		0.37±0.01	1.60±0.03 ^b	5.68±0.02	3.39 ^a ±0.06
<i>L. mesenteroides</i>	1	0.39±0.01	1.51±0.03 ^{bc}	5.68±0.02	3.46 ^a ±0.10
<i>L. plantarum</i> + <i>L. mesenteroides</i>		0.38±0.01	2.11±0.18 ^a	5.72±0.05	2.54 ^b ±0.19

The values are expressed as a mean ± SD (n = 3), and values in a column with the same letters are not significantly ($P > 0.05$) different.

maintain a pH gradient at high organic acid concentration. McDonald et al. (2009) revealed that the growth of *L. mesenteroides* stopped when a pH of 5.4 to 5.7 was reached, while the growth of *L. plantarum* stopped when pH of 4.6 to 4.8 was reached. In the sauerkraut fermentation, the initial pH in a range of 5.67 and 5.75. Consequently, the pH shift for *L. mesenteroides* growth was lower than that for *L. plantarum*. The results of the analysis of variance showed that the addition of culture and salt treatments gave a significant effect ($P < 0.05$) on the increasing of total lactic acid bacteria. Fermentation at 1% salt resulted higher total LAB than that of 0.5%. Salt can pull out the juice containing nutrients in the cabbage. Higher salt concentration, more juice pulled out, consequently more nutrients are available for the LAB growth. However, total LAB in sauerkraut control was lower although the fermentation was carried out at higher salt concentration Table 1. It may be due to the lower LAB loaded at initial fermentation in the control without starter culture. Other researchers also reported that the addition of starter culture can expedite the fermentation process and increase total lactic acid bacteria (Beganovic et al., 2011; Yang et al., 2019).

Addition of starter culture affected significantly on the total acidity and pH after fermentation at salt 0.5% and 1.0%. It reflects the LAB produce organic acid during the fermentation. Those are supported by the data of pH values, whereby the values at final fermentation were lower than those at the initial. These chemical changes are related to the LAB growth during the fermentation. Total acidity and pH of the fermentation product at the combination cultures treatment respectively higher and lower than those at single culture treatment. This agrees with several researchers reports that *L. mesenteroides* produce lactic acid and acetic acid, which caused pH decreasing. *L. plantarum* and *L. brevis* continuing the fermentation until the pH around 3 (Lu et al., 2003; Plengvidhya et al., 2007; Swain et al., 2014). The bacteria oxidize ethanol to acetaldehyde to acetic acid (Chu and Chen, 2006).

3.2 Antioxidant activity and total phenolic contents

The *in vitro* DPPH scavenging ability increased after fermentation of the sauerkraut at all the treatments, reflected from the lower values of IC_{50} at after fermentation than those at before fermentation as presented in Table 3. Lactic acid bacteria are able to activate enzymes that having the function of breaking down the phenolic complex into simple compounds (Tolonen et al., 2004). Invertase, cellulase and amylase are able to break complex bonds between phenolic and other compounds so that increase in total phenolic

content during fermentation (Essawet et al., 2015). The antioxidant activities correlate to the total phenolic contents, which naturally present in the cabbage and increased during the fermentation. Other researchers also reported that the increase of total phenol in sauerkraut goes hand in hand with an increase of antioxidant activity (Ciska et al., 2005; Martinez-Villaluenga et al., 2012; Penas et al., 2012). The bioactive compounds are able to convert free radical compounds into more stable compounds by donating hydrogen atoms and their aromatic hydroxyl (OH) groups (Dipti et al., 2003).

Both *L. plantarum* and *L. mesenteroides* are able to metabolize phenolic compounds in foods. Surprisingly, the lowest IC_{50} values, both at salt 0.5% and 1%, were found at the addition of *L. mesenteroides* culture. The highest increment of total phenolic content occurred in the addition of *L. mesenteroides* culture, consistent with the antioxidant activity.

3.3 Sulforaphane content

The sauerkraut with the highest antioxidant activity was subjected to sulforaphane content analysis by using LC-MS. The chromatogram of the sample and standard are shown in Figure 1. Sulforaphane is a glucosinolate derivative compound widely found as biologically active compound in cabbage. Sulforaphane content in the sauerkraut with- and without *L. mesenteroides* culture is presented in Table 4. Sulforaphane content in the sauerkraut with *L. mesenteroides* starter culture is higher than that of the control. This indicates that the addition of *L. mesenteroides* culture can increase myrosinase activity to break down the glucose bonds on glucoraphanin so that sulforaphane compound is active as antioxidants. Moreover, it has antiproliferative, anti-inflammatory and anti-cancer activities (Xu et al., 2005; Sayed et al., 2014). Sulforaphane has a role in apoptosis of cell proliferation, cancer evolution and stimulation of tumor necrosis factor (TNF- α), IL-1, Lipopolysaccharide (LPS) and in oxidative stress (Suganuma et al., 2011; Thakur et al., 2014; Nallasamy et al., 2014; Greaney et al., 2016). Another compound was also detected by the

Table 4. Sulforaphane content in sauerkraut

Product	Sulforaphane ($\mu\text{g/g}$)
Sauerkraut control (Without culture)	776.47 \pm 3.21 ^a
Sauerkraut (<i>L. mesenteroides</i>)	848.65 \pm 2.14 ^b

The values are expressed as a mean \pm SD (n = 3). The different letter in the same column indicate a significant difference between ($p < 0.05$).

LC-MS with m/z value of 105 and Rt 1.58 namely 2-phenethyl isothiocyanate antimicrobial compound (Abbaoui et al., 2015).

Table 3. Total phenolic content and antioxidant activities before and after in sauerkraut fermentation at different starter cultures and salt concentrations

Culture	Salt (%)	Total Phenolic Content (mg GAE/g)		Antioxidant Activities, IC ₅₀ (ppm)	
		Day 0	Day 5	Day 0	Day 5
<i>L. plantarum</i>		29.99±0.01	58.33±1.46 ^c	155.55±0.12	120.07±3.64 ^a
<i>L. mesenteroides</i>	0.5	29.89±0.03	71.50±0.20 ^a	152.45±0.31	95.55±2.37 ^c
<i>L. plantarum</i> + <i>L. mesenteroides</i>		29.99±0.18	64.10±0.96 ^b	150.44±1.21	99.62±5.68 ^{bc}
<i>L. plantarum</i>		29.00±1.17	59.80±0.20 ^c	150.73±2.00	110.13±0.74 ^{ab}
<i>L. mesenteroides</i>	1	29.50±0.18	72.99±0.88 ^a	154.01±1.67	94.85±1.66 ^c
<i>L. plantarum</i> + <i>L. mesenteroides</i>		29.68±0.13	65.48±0.69 ^b	151.52±1.60	98.84±1.36 ^c

The values are expressed as a mean ± SD (n = 3). The different letter in the same column indicate a significant difference between data on the day 0 and day 5 (p<0.05). GAE = Gallic acid equivalent, IC₅₀ (inhibition concentration at 50% scavenging).




Figure 1. Total ion chromatogram (TIC) of LC-MS analysis of: (A) 2-phenethyl isothiocyanate (control); (B) Sulforaphane (control); (C), (D) TIC (sauerkraut + *L. mesenteroides*); (E) 2-phenethyl isothiocyanate (sauerkraut + *L. mesenteroides*), and (F); Sulforaphane (sauerkraut + *L. mesenteroides*)

4. Conclusion

The starter cultures application of *L. plantarum* and *L. mesenteroides* at lower salt fermentation increased total lactic acid bacteria, total acidity and decreased pH. *L. mesenteroides* resulted in the highest total phenolic content and the lowest IC₅₀ value. Sauerkraut with the addition of *L. mesenteroides* contains sulforaphane higher than that of control.

Conflict of interest

The authors declare no conflict of interest

eISSN: 2550-2166

Acknowledgments

This work was financially supported through Professor Research Grant Brawijaya University with grant number 2571/UN10.F10/PN/2019.

References

Abbaoui, B., Lucas, C.R., Riedl, K.M., Clinton, S.K. and Mortazavi, A. (2015). Cruciferous vegetables, isothiocyanates, and prevention of bladder cancer. *Current Pharmacology Reports*, 1(4), 272–282. <https://doi.org/10.1007/s40495-015-0024-z>

© 2020 The Authors. Published by Rynnye Lyan Resources

- Beganovic, J., Pavune, A.L. Gjuracic, K., Spoljarec, M., Suskovic, J. and Kos, B. (2011). Improved sauerkraut production with probiotic strain *Lactobacillus plantarum* L4 and *Leuconostoc mesenteroides* LMG 7954. *Journal of Food Science*, 76(2), M124-M129. <https://doi.org/10.1111/j.1750-3841.2010.02030.x>
- Martinez-Villaluenga, C., Penas, E., Sidro, B., Ullate, M., Frias, J. and Vidal-Valverde, C. (2012). White cabbage fermentation improves ascorbigen content, antioxidant and Nitric Oxide production inhibitory activity in LPS-induced macrophages. *LWT-Food Science and Technology*, 46(1), 77-83. <https://doi.org/10.1016/j.lwt.2011.10.023>
- Chu, C.S. and Chen, C. (2006). Effect of origin and fermentation time on the antioxidant activity of kombucha. *Food Chemistry*, 98(3), 502-507. <https://doi.org/10.1016/j.foodchem.2005.05.080>
- Ciska, E., Magdalena, K. and Agnieszka, K. (2005). Antioxidant activity of extracts of white cabbage and sauerkraut. *Polish Journal of Food and Nutrition Sciences*, 55(4), 367-373.
- Dipti, P., Yogesh, B., Kain, A.K., Pauline, T., Anju, B., Sairam, M., Singh, B., Mongia, S.S., Kumar, G.I. and Selvamurthy, W. (2003). Lead induced oxidative stress: beneficial effects of kombucha tea. *Biomedical Environment Science*, 16(3), 276-282.
- Goh, W.N., Rosma, A., Kaur, B., Fazilah, A., Karim, A.A. and Bhat, R. (2012). Fermentation of black tea broth (kombucha): I. Effects of sucrose concentration and fermentation time on the yield of microbial cellulose. *International Food Research Journal*, 19(1), 109-117.
- Greaney, A.J., Maier, N.K., Leppla, S.H. and Moayeri, M. (2016). Sulforaphane inhibits multiple inflammasomes through an Nrf2-independent mechanism. *Journal of Leukocyte Biology*, 99(1), 189-199. <https://doi.org/10.1189/jlb.3A0415-155RR>
- Johanningsmeier, S., McFeeters, R.F., Fleming, H.F. and Thompson, R.L. (2007). Effects of *Leuconostoc mesenteroides* starter culture on fermentation of cabbage with reduced salt concentrations. *Journal of Food Science*, 72(5), M166-172. <https://doi.org/10.1111/j.1750-3841.2007.00372.x>
- Kim, H.J., Lee, M.J., Jeong, M.H. and Kim, J.E. (2017). Identification and quantification of glucosinolates in kimchi by liquid chromatography-electrospray tandem mass spectrometry. *International Journal of Analytical Chemistry*, 2017, 1-8. <https://doi.org/10.1155/2017/6753481>
- Liang, H., Yuan, Q.P., Dong, F.R. and Liu, Y.M. (2006). Determination of sulfurpahane in broccoli and cabbage by high-performance liquid chromatography. *Journal of Food Composition and Analysis*, 19(5), 473-476. <https://doi.org/10.1016/j.jfca.2005.11.005>
- Lu, Z., Breidt, F., Plengvidhya, V. and Fleming, H.P. (2003). Bacteriophage ecology in commercial sauerkraut fermentations. *Applied and Environmental Microbiology*, 69, 3192-3202. <https://doi.org/10.1128/AEM.69.6.3192-3202.2003>
- McDonald, L.C., Fleming, H.P. and Hassan, H.M. (2009). Acid Tolerance of *Leuconostoc mesenteroides* and *Lactobacillus plantarum*. *Applied Environmental Microbiology*, 56(7), 2120-2124. <https://doi.org/10.1128/AEM.56.7.2120-2124.1990>
- Molyneux, P. (2004). The use of the stable free radical diphenyl picryl hydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarinn Journal of Science and Technology*, 26(2), 211-219.
- Murray, R.K., Granner, D.K. and Rodwell, V.W. (2009). *Biokimia*. Harper. 27th ed. Jakarta: Buku Kedokteran EGC.
- Nallasamy, P., Si, H., Babu, P.V., Pan, D., Fu, Y., Brooke, E.A., Shah, H., Zhen, W., Zhu, H., Liu, D., Li, Y. and Jia, Z. (2014). Sulforaphane reduces vascular inflammation in mice and prevents TNF- α -induced monocyte adhesion to primary endothelial cells through interfering with the NF- κ B pathway. *The Nutritional Biochemistry*, 25(8), 824-833. <https://doi.org/10.1016/j.jnutbio.2014.03.011>
- Penas, E., Martinez-Villaluenga, C., Frias, J., Sanchez-Mrtinez, M.J., Perez-Corona, T.P., Madrid, Y., Camara, C. and Vidal-Valverde, C. (2012). Se improves indoleglucosinolate hydrolysis product content, Se-methylselenocysteine content, antioxidant capacity and potential anti-inflammatory properties of sauerkraut. *Food Chemistry*, 132(2), 907-914. <https://doi.org/10.1016/j.foodchem.2011.11.064>
- Penas, E., Frias, J., Sidro, B. and Vidal-Valvede, C. (2010). Chemical evaluation and sensory quality of sauerkraut obtained fermentations at different NaCl levels from *Brassica oleracea* Var. capitata Cv. Bronco grown in eastern Spain. Effect of storage. *Agricultural and Food Chemistry*, 58, 3549-3557. <https://doi.org/10.1021/jf903739a>
- Plengvidhya, V., Breidt, F.Jr., Lu, Z. and Fleming, H.P. (2007). DNA fingerprinting of lactic acid bacteria in sauerkraut fermentations. *Applied and Environmental Microbiology*, 73, 7697-7702. <https://doi.org/10.1128/AEM.01342-07>
- Ranggana, S. (1977). *Manual of Analysis of Fruit and Vegetable Products*. New Delhi, India: McGraw Hill

- Publishing Co.
- Romeo, L., Lori, R., Rollin, P., Bramanti, P. and Mazzon, E. (2018). Isothiocyanates: an overview of their antimicrobial activity against human infections. *Molecules*, 23(3), 624. <https://doi.org/10.3390/molecules23030624>
- Sayed, R.H., Khalil, W.K.B., Salem, H.A., Kenawy, S.A. and El-Sayeh, B.M. (2014). Sulforaphane increases the survival rate in rats with fulminant hepatic failure induced by d-galactosamine and lipopolysaccharide. *Nutrition Research*, 34(11), 982-989. <https://doi.org/10.1016/j.nutres.2014.10.003>
- Suganuma, H., Fahey, J.W., Bryan, K.E., Healy, Z.R. and Talalay, P. (2011). Stimulation of phagocytosis by sulforaphane. *Biochemical and Biophysical Research Communications*, 405(1), 146–151. <https://doi.org/10.1016/j.bbrc.2011.01.025>
- Swain, M.R., Anandharaj, M., Ray, R.C. and Parveen Rani, R. (2014). Fermented fruits and vegetables of Asia: a potential source of probiotics. *Biotechnology Research International*, 2014, 1-19. <https://doi.org/10.1155/2014/250424>
- Tolonen, M., Rajaniemi, S., Pihlavan, J.M., Korhonen, H. and Ryhanen, E.L. (2004). Formation of nisin, plant derived biomolecules and antimicrobial activity in starter culture fermentations of sauerkraut. *Food Microbiology*, 21(2), 391-395. [https://doi.org/10.1016/S0740-0020\(03\)00058-3](https://doi.org/10.1016/S0740-0020(03)00058-3)
- Thakur, V.S., Deb, G., Babcook, M.A. and Gupta, S. (2014). Plant phytochemicals as epigenetic modulators: role in cancer chemoprevention. *The AAPS Journal*, 16(1), 151-163. <https://doi.org/10.1208/s12248-013-9548-5>
- Xiaozhe, Y., Wenzhong, H., Aili, J., Zhilong, X., Yaru, J., Yuge G., Sarengaowa., Xiangyan, Y. (2019). Effect of salt concentration on quality of northeast sauerkraut fermented by *Leuconostoc mesenteroides* and *Lactobacillus plantarum* salt effect on northeast sauerkraut fermentation. *Food Bioscience*, 30, 1-8. <https://doi.org/10.1016/j.fbio.2019.100421>
- Xu, C., Shen, G., Chen, C., Gelinis, C. and Kong, A.T. (2005). Suppression of NF- κ B and NF- κ B-regulated gene expression by sulforaphane and PEITC through I κ B α , IKK pathway in human prostate cancer PC-3 cells. *Oncogene*, 24, 4486–4495. <https://doi.org/10.1038/sj.onc.1208656>
- Yang, J., Paulino, R., Janke-Stedronsky, S. and Abawi, F. (2007). Free radical scavenging activity and total phenol of noni (*Moriadaeitrifolia L.*) juice and powder in processing and storage. *Food Chemistry*, 102(1), 302-308. <https://doi.org/10.1016/j.foodchem.2006.05.020>
- Yang, X., Hu, W., Jiang, A., Xiu, Z., Ji, Y., Guan, Y., Sarengaowa and Yang, X. (2019). Effect of salt concentration on quality of Chinese northeast sauerkraut fermented by *Leuconostoc mesenteroides* and *Lactobacillus plantarum*. *Food Bioscience* 30, 100421. <https://doi.org/10.1016/j.fbio.2019.100421>

Effect of Lactobacillus plantarum and Leuconostoc mesenteroides

ORIGINALITY REPORT

18%

SIMILARITY INDEX

6%

INTERNET SOURCES

18%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Elok Zubaidah, Mentari S. Arum, Tri D. Widyaningsih, Aldila P. Rahayu. "Sauerkraut with the Addition of Lactobacillus casei: Effects of Salt and Sugar Concentrations on Fermentation and Antioxidant Activity", Current Nutrition & Food Science, 2020
Publication **10%**
- 2** [bmccomplementmedtherapies.biomedcentral.com](https://www.bmccomplementmedtherapies.biomedcentral.com) **2%**
Internet Source
- 3** Suzanne Johanningsmeier. "Effects of Leuconostoc mesenteroides Starter Culture on Fermentation of Cabbage with Reduced Salt Concentrations", Journal of Food Science, 6/2007
Publication **1%**
- 4** www.hindawi.com **1%**
Internet Source
- 5** Tiaan Heunis, Shelly Deane, Salome Smit, Leon M. T. Dicks. " Proteomic Profiling of the **1%**

Acid Stress Response in 423 ", Journal of Proteome Research, 2014

Publication

6

Varzakas, Theodoros, and Constantina Tzia. "Blanching", Contemporary Food Engineering, 2015.

Publication

7

"Minimally Processed Refrigerated Fruits and Vegetables", Springer Science and Business Media LLC, 2017

Publication

8

E Zubaidah, R A Ifadah, C A Afgani. "Changes in chemical characteristics of kombucha from various cultivars of snake fruit during fermentation", IOP Conference Series: Earth and Environmental Science, 2019

Publication

9

C. Martinez-Villaluenga, E. Peñas, B. Sidro, M. Ullate, J. Frias, C. Vidal-Valverde. "White cabbage fermentation improves ascorbigen content, antioxidant and nitric oxide production inhibitory activity in LPS-induced macrophages", LWT - Food Science and Technology, 2012

Publication

10

www.e3s-conferences.org

Internet Source

1 %

1 %

1 %

1 %

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On