PAPER • OPEN ACCESS

The 2nd International Conference on Natural Resources and Life Sciences (NRLS-2018)

To cite this article: 2019 IOP Conf. Ser.: Earth Environ. Sci. 293 011001

View the article online for updates and enhancements.



This content was downloaded from IP address 158.140.169.37 on 25/11/2020 at 04:29

PAPER • OPEN ACCESS

The 2nd International Conference on Natural Resources and Life Sciences (NRLS-2018)

To cite this article: 2019 IOP Conf. Ser.: Earth Environ. Sci. 293 011001

View the article online for updates and enhancements.

MITelah diperiksa kebenarannya dan sesuai dengan aslinya Declares this translation to correspond to the original Surabaya, Dekan Fakultas Taknik 152 Dean Faculty of Engineering Survadi Ismadji, IPM., ASEAN Eng. NIK. 521.93.0198 140



This content was downloaded from IP address 158.140.169.37 on 25/11/2020 at 04:29



Website http://event.ubaya.ac.id/nrls/

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution Ð of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd



PREFACE: the 2nd International Conference on Natural Resources and Life Sciences (NRLS) 2018

The 2^{nd} International Conference on Natural Resources and Life Sciences (NRLS) 2018 has been organized by the Faculty of Biotechnology of University of Surabaya, Indonesia. The theme of this conference is "*Managing Natural Resources for Better Health*". Focusing on biological aspects, the conference has facilitated chances of collaborations in research and development – as well as enlarged joint activities regarding natural resource management – among academics and professionals in their attempts to contribute further to the community through their respective fields.

The 2nd NRLS 2018 was held on August 23–24, 2018. This conference presented eight international speakers from five countries: Indonesia, Malaysia, the Netherlands, Singapore, and Thailand. Over 200 representatives of 48 institutions participated in this event, involving more than 74 abstracts submitted in the form of oral and poster presentations. After a rigorous selection process, the Scientific & Editorial Board have decided to publish 43 manuscripts in IOP Conference Series: Earth and Environmental Science (EES), an international proceedings indexed in Scopus, Scimago, Conference Proceedings Citation Index-Science (CPCI-S) of Clarivate Analytics's Web of Science, etc. From 43 selected ones above, 21 manuscripts were results of joint researches between Indonesia and various countries, e.g. Australia, England, Georgia, Germany, India, Japan, Latvia, Lithuania, Malaysia, the Netherlands, the Republic of Korea, Spain, and Sweden. Those manuscripts cover various biological themes, i.e. Food Biotechnology, Agricultural Biotechnology, Medical Biotechnology & Forensics, and Environmental Biotechnology & Renewable Energy. Each of the 43 manuscripts in IOP Conference Series-EES have been reviewed by at least two experts using double-blind system. The published manuscripts have passed all necessary improvement requirements (according to the IOP Proceedings standard), reviewer's comments, SI (Système International d'Unités), and similarity tests (with the highest threshold of 25 %) as well as editing procedure by professional editors from seven countries (Georgia, India, Indonesia, Latvia, Lithuania, Malaysia, and Sweden).

Our appreciation goes to the reviewers, editors and members of the Scientific & Editorial Board for their big efforts in reviewing and improving the manuscripts. For the generous supports in succeeding the NRLS-2018, we extend our gratitute toward the University of Surabaya's management and supporting units, our co-hosts the Faculty of Pharmacy and the Faculty of Medicine, and our sponsors VISION TEKNIK, SCIENCEWERKE, INDOLAB UTAMA, and MEGAH SEJAHTERA SCIENTIFIC.

Last but not least, we thank you all presenters and attendees for the active contribution to share scientific ideas, inspire new researches, and exchange new contacts for closer co-operations. We hope you have had enjoyable time with us and are currently encouraged to collaborate further in order to explore natural resources and life sciences in various aspects of living. We look forward to welcoming you and your team in the 3rd NRLS-2020!

Surabaya May 02, 2019 Republic of Indonesia - National Education Day



Johan Sukweenadhi, Ph.D. Executive Chief of the 2nd NRLS-2018

Editor in Chief: Roy Hendroko Setyobudi (Malang, IDN)

Board of Editor: Mariana Wahjudi (Surabaya, IDN), Maizirwan Mel (Kuala Lumpur, MYS), Olga Anne (Klaipėda, LTU), Peeyush Soni (Kharagpur, IND), Tsitsino Turkadze (Kutaisi, GEO), Yahya Jani (Kalmar, SWD), and Zane Vincēviča-Gaile (Riga, LVA).

PAPER • OPEN ACCESS

Scientific & Editorial Boards the 2nd NRLS-2018

To cite this article: 2019 IOP Conf. Ser.: Earth Environ. Sci. 293 011004

View the article online for updates and enhancements.



This content was downloaded from IP address 158.140.169.37 on 25/11/2020 at 04:28



SCIENTIFIC & EDITORIAL BOARDS the 2nd NRLS-2018

- Andoniana Rakoto Malala, Centre de Formation et d'Application du Machinisme Agricole (CFAMA), Antsirabe, MDG
- Christina Avanti, Faculty of Pharmacy, University of Surabaya, IDN •
- Fauna Herawati, Faculty of Pharmacy, University of Surabaya, IDN
- Finna Setiawan, Faculty of Pharmacy, University of Surabaya, IDN •
- Ida Bagus Made Artadana, Faculty of Biotechnology, University of Surabaya, IDN •
- Johan Sukweenadhi, Faculty of Biotechnology, University of Surabaya, IDN
- Juris Burlakovs, Faculty of Health and Life Sciences, Linnaeus University, SWD •
- Kartini, Faculty of Pharmacy, University of Surabaya, IDN •
- Lieke Riadi, Faculty of Chemical Engineering, University of Surabaya, IDN •
- Lim Kok Kuan, Nippon Biodiesel Fuel, Odawara-city, Kanagawa, JPN •
- Maizirwan Mel, Department of Biotechnology Engineering, Faculty of Engineering . International Islamic University Malaysia. MYS
- Mariana Wahjudi, Faculty of Biotechnology, University of Surabaya, IDN
- Maria Goretti Marianti Purwanto, Fac.of Biotechnology, University of Surabaya, IDN •
- Oeke Yunita, Faculty of Biotechnology, University of Surabaya, IDN
- Olga Anne, Faculty of Marine Technology and Natural Sciences of Klaipeda University, LTU
- Peeyush Soni, Department of Agricultural and Food Engineering, Indian Institute of • Technology Kharagpur, IND
- Popy Hartatie Hardjo, Faculty of Biotechnology, University of Surabaya, IDN
- Praptiningsih Gamawati Adinurani, Department of Agrotechnology, Faculty of Agriculture, Merdeka University of Madiun, IDN
- Rangga Kala Mahaswa, Graduate Program, Faculty of Philosophy, Universitas • Gadjah Mada, Yogyakarta, IDN
- Rika Yulia, Faculty of Pharmacy, University of Surabaya, IDN
- Risma Ikawaty, Faculty of Medical, University of Surabaya, IDN
- Roy Hendroko Setyobudi, Postgraduate of Agriculture Science, Waste Laboratory University of Muhammadyah Malang, IDN
- Sulistyo Emantoko, Faculty of Biotechnology, University of Surabaya, IDN
- Tjie Kok, Faculty of Biotechnology, University of Surabaya, IDN
- Tsitsino Turkadze, Department of Chemical and Environmental Technologies, Akaki Tsereteli State University, Kutaisi, GEO
- Yahya Jani, Departement of Biology and Environmental Science, Linnaeus • University, Kalmar, SWD
- Yalun Arifin, Faculty of Food Business and Technology, Prasetiya Mulya University, • Jakarta, IDN
- Zahrah Nurfadhilah, Faculty of Biology, Universitas Gadjah Mada, Yogyakarta, IDN
- Zane Vincēviča-Gaile, Department of Environmental Science, University of Latvia, Riga – LVA

 $(\mathbf{\hat{H}})$

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

Table of contents

Volume 293

2019

♦ Previous issue
 Next issue ▶

The 2nd International Conference on Natural Resources and Life Sciences (NRLS-2018) 23–24 August 2018, Ibis Styles Hotel, Surabaya, Indonesia

Accepted papers received: 17 May 2019 Published online: 01 July 2019

Open all abstracts

Preface

OPEN ACCESS			011001
The 2nd Internat	ional Conference or	n Natural Resources and Life Sciences (NRLS-2018)	
	View article	PDF	
OPEN ACCESS			011002
Conference Phot	os		
	View article	PDF	
OPEN ACCESS			011003
Conference Spor	isors		
	View article	🔁 PDF	
OPEN ACCESS			011004
Scientific & Edit	corial Boards the 2nd	⁴ NRLS-2018	
	View article	🔁 PDF	
OPEN ACCESS			011005
Peer review state	ement		
	View article	🔁 PDF	

 \odot

Papers

ODEN ACCESS			
Android based ri	ce pest detection sy	stem using learning vector quantization method	012001
A Budiman, P Utor	no and S Rahayu		
	View article	PDF	
OPEN ACCESS			012002
Community beha	vior and single-use	plastic bottle consumption	
A Khoironi, S Ang	goro and S Sudarno		
	View article	🔁 PDF	
OPEN ACCESS			012003
Genetic variabili	ty of soybean (Glyc	eine max L. Merrill) genotypes for pod shattering resistance	
A Krisnawati and N	A M Adie		
	View article	🔁 PDF	
OPEN ACCESS	a from nolm oil mil	1 offluent (DOME): Challenges and encerturities in Indensia	012004
A D · · · K · · ·		in endent (POME): Chanenges and opportunities in indonesia	
A Kajani, Kusnadi,	A Santosa, A Saepud	in, S Gobikrishnan and D Andriani	
	View article	PDF	
OPEN ACCESS			012005
Early detection o protein analysis	f somaclonal variat	ion in oil palm callus culture through cytological and SDS-PAGE	
A Sahara, Reflini, G	C Utomo and T Liwan	g	
+ Open abstract	View article	PDF	
OPEN ACCESS Genetic diversity and microsatellit	r of Indonesian soyt e markers	bean (<i>Glycine max</i> L. Merrill) germplasm based on morphological	012006
A Sulistyo, F C Ind	riani, M J Mejaya, A	N Sugiharto and J Agranoff	
+ Open abstract	View article	PDF	
OPEN ACCESS			012007
Fermentation qua sugarcane molass	ality of <i>Pennisetum</i> ses in tropic	purpureum cv. Mott ensiled with Lactobacillus plantarum and	
A Wahyudi, L Heno	draningsih, Sutawi, R	H Setyobudi and M Mel	
	View article	🔁 PDF	

The evaluation of estimated breeding value and the most probable producing ability for the basis selection of Ettawa crossbred goat (*Capra hircus* sp.) at Malang, East Java, Indonesia

A Winaya, Suyatno	, P Coy and N Fauzi		
	View article	🔁 PDF	
OPEN ACCESS			012009
Menthol from the	e stem and leaf <i>in-vi</i>	itro Mentha piperita Linn.	
B K Wijaya, P H H	ardjo and S Emantoko		
	View article	🔁 PDF	
OPEN ACCESS			012010
A review: Bioma	iss-based fuel pellet	usage in biomass gasifier-an optimization on non-wood material	
D Andriani, T D At	tmaja, M Arifin, A Raj	ani and Kusnadi	
	View article	🔁 PDF	
OPEN ACCESS Effectiveness of seeds nila plants	the use of organic w <i>(Indigofera</i> sp.)	vaste as fertilizer and physical scarification of seeds on growth of	012011
D Roeswitawati, M	Huda, D Indratmi and	l M Mel	
	View article	PDF	
OPEN ACCESS Physiology and g condition	genotyping of adapti	ive and sensitive oil palm progenies under unwatered stress	012012
D Yono, E Purwant	ti, A Sahara, Y A Nugr	roho, Z A Tanjung, R Aditama, C Dewi, A E Sihotang, C Utomo and T Liwang	
+ Open abstract	View article	PDF	
OPEN ACCESS Effect of phospho gallus domesticu	oric acid pretreatme s L.) bones	ent on characterization of gelatin from broiler chicken (Gallus	012013
D Yuliani, A Maun	atin, A Jannah and H H	H Fauziyyah	
	View article	🔁 PDF	
OPEN ACCESS Selection for dro	ught-resistant rice (Oryza sativa L.) using polyethylene glycol	012014
E D Purbajanti, F K	Kusmiyati, E Fuskhah,	R Rosyida, P G Adinurani and Z Vincēviča-Gaile	
+ Open abstract	View article	PDF	
OPEN ACCESS			012015

Effectiveness the source of nitrogen from NO₃ and NH₄ for *Panicum maximum* Jacq. growth in saline soil

E D Purbajanti, P G Adinurani,	T Turkadze, Z Vincēviča-	Gaile and R H Setyobudi
--------------------------------	--------------------------	-------------------------

5 ,	,	, <u>,</u>	
	View article	🔁 PDF	
OPEN ACCESS Anticancer poten cell line	tial from ethanol ex	stract of Zanthoxylum acanthopodium DC. seed to against MCF-7	012016
E V Arsita, D E Sar	agih and K Aldrin		
	View article	🔁 PDF	
OPEN ACCESS			012017
Long-term Proton	n Pump Inhibitors in	nduces recurrent Urinary Tract Infections: A case study	
F Cokro and S T Ar	rang		
	View article	🔁 PDF	
OPEN ACCESS Molecular dockir against VEGFR-2	ng, drug-likeness, an 2	nd ADMET study of 1-benzyl-3-benzoylurea and its analogs	012018
F Suhud, D H Tjahj	ono, T A Yuniarta, G	S Putra and J Setiawan	
	View article	PDF	
OPEN ACCESS Microcontroller a made from cassav	application for pH a va (Manihot esculer	nd temperature control system in liquid sugar liquefaction process <i>nta</i> Crantz.)	012019
H Santosa, Yuliati, I	I J Mulyana, D S Sirai	it and S D Novitasari	
	Tiew article	PDF	
OPEN ACCESS The effect of ferm <i>polyrhiuzus</i> [F.A.	nentation process o C. Weber] Britton o	n physical and chemical characteristics of pitaya (<i>Hylocereus</i> & Rose) stem flour	012020
H Soedjatmiko, R C	Chrisnasari and P H Ha	ardjo	
	View article	🔁 PDF	
OPEN ACCESS The performance drought stress at	of three local rice (early seedling stage	(<i>Oryza sativa</i> L.) cultivar from East Kalimantan - Indonesia under	012021
I B M Artadana, I T	Dewi and J Sukween	adhi	
+ Open abstract	View article	PDF	
OPEN ACCESS Coffee reduced th	ne production of ne	utrophil superoxide radical in vitro	012022

I D A Susilawati, A Safaatin and J Burlakovs

	View article	🔁 PDF
 Open abstract 	View article	🔁 PDF

Maftuchah, I Z Fahr	ni, A Zainudin, A Ikh	wan, Djumali and L K Kuan	
OPEN ACCESS The drought stres	s tolerance of phys	ic nut (Jatropha curcas Linn.) genotypes	012029
+ Open abstract	View article	PDF	
M Wahjudi, W D Ku	urniawan, M T Gulton	n, I B M Artadana and N I M Puad	
OPEN ACCESS Inhibition of <i>Burk</i> isolates	kholderia cenocepa	cia H111 quorum sensing system by environmental bacterial	012028
	View article	PDF	
M M Adie, A Krisna	awati and D Suryati		
OPEN ACCESS Characterization of Indonesia	of adaptive and pro	ductive soybean (G. max L.) genotypes in dry land of Kalimantan,	012027
+ Open abstract	View article	PDF	
L Zalizar, I D Rahay	yu, Sujono and Y A N	or	
OPEN ACCESS Potency of <i>Fibran</i>	urea tinctora Lour.	extract as anti-bacterial agents towards pathogenic bacteria	012026
	View article	🔁 PDF	
K Siregar, A H Tam	bunan, Sholihati, S S	Wirawan and T Araki	
OPEN ACCESS Comparison of en biodiesel products assessment	ergy production, no ion from oil palm (2	et energy balance, net energy ratio, and renewable index for Elaeis guineensis Jacq.) and jatropha (Jatropha curcas L.) based on l	012025 ife cycle
	View article	🔁 PDF	
J Sukweenadhi, J Y	Choi, Y J Kim, L Kal	iraj, S Abid, J C Ahn and D C Yang	
OPEN ACCESS Callus induction a Dendropanax mod	and <i>in vitro</i> mass cu <i>rbifera</i> Lev.	lture of adventitious roots from leaf segment explants of	012024
← Open abstract	E View article	► PDF	
I D Rahayu, W Wid	odo, I Prihartini, A W	inaya, L Zalizar, 1 Untari and M Mel	
performances and	meat nutritional co	omposition of broiler chickens	
The potential of e	xtract of Zingiber z	<i>terumbet</i> (L.) Smith as a feed additive to improve the production	012023
OPEN ACCESS			012022

	View article	PDF	
OPEN ACCESS Characterization of	of thermostable chit	inase from <i>Bacillus licheniformis</i> B2	012030
N Jayanthi, M G M	Purwanto, R Chrisnasa	ari, T Pantjajani, A Wahjudi and M Sugiarto	
	View article	PDF	
OPEN ACCESS Antibacterial activ epidermidis (Evan	vity roll on deodora ns 1916) <i>in vitro</i>	nt with Pluchea indica (L.) leaf extract against Staphylococcus	012031
O Komala, I Y Wier	ndarlina and N Rizqiya	ina	
	Tiew article	PDF	
OPEN ACCESS Production potens planting pattern as	tial of sweet corn (Z nd phosphorus sour	<i>Lea mays</i> Linn. var. Saccharata Sturt) 'Bonanza' to different ces	012032
P G Adinurani, S Ka	inayu, L S Budi, S Pan		
	View article	PDF	
OPEN ACCESS Genome-wide ide <i>Ganoderma bonin</i>	ntification of oil pa	lm (Elaeis guineensis Jacq.) chitinases and their response to	012033
R Aditama, R Tryon	o, Z A Tanjung, C Uto	omo and T Liwang	
	View article	🄁 PDF	
OPEN ACCESS Effect of new NPI	K fertilizer on lowla	and rice (Oryza sativa L.) growth	012034
K Budiono, P G Adi			
	View article	PDF	
OPEN ACCESS Prospect of Fe not	n-heme on coffee fl	our made from solid coffee waste: Mini review	012035
R H Setyobudi, L Za	alizar, S K Wahono, W	Widodo, A Wahyudi, M Mel, B Prabowo, Y Jani, Y A Nugroho, T Liwang and	
A Zaebudin			
+ Open abstract	View article	PDF	
OPEN ACCESS			012036
Biomass to metha	nol plant based on g	gasification of palm empty fruit bunch	
R Heryadi, A S Uyu	n, E Yandri, S M Nur,	K Abdullah and O Anne	
+ Open abstract	View article	🔁 PDF	

OPEN ACCESS			012037
Review article: N	Ayopia - Geneticall	y inherited or environmental influences	
S E E Tjoa and S E	D Putra		
+ Open abstract	View article	🔁 PDF	
OPEN ACCESS			012038
Rapid reversed-p content in food p	bhase high performa products	nce liquid chromatography assay of Tert-butylhydroquinone	
S Sutanto and M G	M Purwanto		
	View article	🔁 PDF	
OPEN ACCESS Developmental e (<i>Bemisia tabaci</i> (effect of cashew nut Genn.)	shell extract against nymphal instar of Silver leaf Whitefly	012039
W R Andayanie and	d N Ermawati		
+ Open abstract	View article	🔁 PDF	
Identification of West Java - Indo Wuryantoro, I R Pu	a local variety of 'u nesia based on tube Ispitawati, R I Fitriyan	wi' (<i>Dioscorea alata</i> Linn.) in four agro-climate regions of East- r character ii and P Soni	
	View article	PDF	
OPEN ACCESS Genome-wide SN restriction site as	NP-discovery and an associated DNA sequ	nalysis of genetic diversity in oil palm using double digest	012041
Y A Nugroho, Z A	Tanjung, D Yono, A S	S Mulyana, H M Simbolon, A S Ardi, Y Y Yong, C Utomo and T Liwang	
	View article	🔁 PDF	
OPEN ACCESS Application of bl	lood donor routine o	letector using K-Nearest neighbors	012042
Y Nurdiansyah, P I	Pandunata, N D Praset	yo, A Trihartono, F G Putrianti and F Wijayanto	
	View article	🔁 PDF	
OPEN ACCESS Identification of	the bacterium isolat	te from Mackerel Fish (Rastrelliger sp.) using 16S rRNA Gene	012043
Y S Ismail, Febrian	ni, C Yulvizar and R Ra	amadhani	
	Tiew article	🔁 PDF	

JOURNAL LINKS

Information for organizers	
Information for authors	
Search for published proceedings	
Contact us	

Reprint services from Curran Associates

PAPER • OPEN ACCESS

Microcontroller application for pH and temperature control system in liquid sugar liquefaction process made from cassava *(Manihot esculenta* Crantz.)

To cite this article: H Santosa et al 2019 IOP Conf. Ser.: Earth Environ. Sci. 293 012019

View the article online for updates and enhancements.

Microcontroller application for pH and temperature control system in liquid sugar liquefaction process made from cassava (Manihot esculenta Crantz.)

H Santosa^{1,*}, Yuliati², I J Mulyana¹, D S Sirait² and S D Novitasari¹

¹Industrial Engineering Department, Widya Mandala Surabaya Catholic University Jl. Kalijudan 37, Surabaya, Jawa Timur, 60114, Indonesia ²Electrical Engineering Department, Widya Mandala Surabaya Catholic University Jl. Kalijudan 37, Surabaya, Jawa Timur, 60114, Indonesia

*Corresponding author: hadi-s@ukwms.ac.id, yuliati@ukwms.ac.id

Abtract. Sugar is a major source of sweetening but the increase of sugar demand is not in line with sugar cane production capacity in Indonesia. Therefore it is necessary to look for alternative other sweeteners as sugar cane substitution. The process of making glucose syrup made from cassava, sweet potato corn, sago is relatively simple and can be done on a small or medium scale industry. But in fact in Indonesia, glucose syrup producers are almost all big industries. This is due to the lack of development of appropriate technology for the manufacture of liquid sugar. Based on preliminary experiments, there is a significant influence of the combination of α -amylase enzyme factor and liquefaction time factor on reducing sugar content. This is by controlling the processing time, temperature (70 to 80) °C by stirring constantly and pH level 5.5. These three parameters are controlled and are being implemented with the design of an industrial liquid sugar processing apparatus. This system is designed with stainless steel material, and arduino microcontroller. Consumption of system power when the standby condition is 4.5 W. When the system is working at 311 W by activating the stirrer motor, gas valve and lighter.

Keywords: Appropriate technology, arduino, enzyme, glucose.

1. Introduction

Sugar as one of the nine basic human needs is a major source of sweetening and is widely used throughout the community. Sugar demand in Indonesia always increases along with the increase of population and industry growth in Indonesia. Ironically, the increase of sugar demand is not in line with sugar cane production capacity in Indonesia. so that import becomes one choice. In 2012, the need for white crystal sugar is 5.13×10^6 t, of which 2.60×10^6 t is household and the remaining 2.53×10^6 t is the industry need. While the amount of production is only 2.5×10^6 t. Dependence on imports will continue to be consistent with population growth, and increased public revenues and industry growth. According to the Indonesian Refined Sugar Association (AGRI), in 2015 the domestic production is only around 2.47×10^6 t and national sugar needs of 6×10^6 t or equivalent to 6.38×10^6 t of raw sugar. As a result, there is a deficit of around 3.53×10^6 t which must be covered by imports [1].

Therefore it is necessary to look for alternative other sweeteners as sugar cane substitution. Sugar made from starch like cassava, sweet potato corn, sago can be an alternative [2]. The process of making glucose syrup is relatively simple and can be done on a small or medium scale industry. But in

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

The 2nd International Conference on Natural Resources and Life Sciences	(NRLS)	IOP Publishing
IOP Conf. Series: Earth and Environmental Science 293 (2019) 012019	doi:10.1088/1755-12	315/293/1/012019

fact in Indonesia, glucose syrup producers are almost all big industries. This is due to the lack of development of appropriate technology for the manufacture of liquid sugar is simple and cheap. In this research will be done the design of equipment production process of liquid sugar made from raw cassava that can be developed by small industry. The device is designed with stainless steel material, and uses arduino uno microcontroller as its controlling center, the working of this tool is the pH meter probe and the Resistance Temperature Detector (RTD) produces a measured resistance value and is accepted by the signal conditioning circuit. If the arduino uno detects and yes changes the measured voltage value and does not match the acid-base and temperature limits then the arduino will run the acid-base pump and the heating regulator.

2. Materials and methods

2.1. Material research

Liquid sugar is known to the public by the name glucose syrup or fructose syrup. Glucose syrup is a condensed solution including monosaccharide groups obtained from starch by means of a complete hydrolysis by using an acid or enzyme catalyst, further purified and thickened. Diagram process flow the making of liquid sugar can be seen in figure 1.



Figure 1. Liquid sugar making process [3].

The process of making liquid sugar is divided into several stages:

(i) Liquefaction stage: Liquefaction process is the process of change of starch from condensed to dilute. The mixture of starch and water (star suspension) heated to boiling will thicken (gelatinization). The ratio of water and flour is 3:1 stirred until well mixed and then included a number of α -amylose enzymes of 1 mL kg⁻¹ of starch. A pH setting of pH 6.2 to pH 6.4 was performed with the addition of one table-spoon of lime. Liquefaction process can be stopped if the solution is completely liquid and clear brown.

(ii) The stage of saccharification: The process of saccharification is the process of dextrin to sugar changes. The starch has been broken down into a later descooled cooling from 105 °C to 60 °C with the addition of an enzyme amyloglucosidase of 1 mL kg⁻¹ of starch to break down the descrank chain

The 2nd International Conference on Natural Resources and Life Sciences	(NRLS)	IOP Publishing
IOP Conf. Series: Earth and Environmental Science 293 (2019) 012019	doi:10.1088/1755-13	15/293/1/012019

into glucose. The saccharification process takes a maximum of 72 h. The saccharification process is completed when the viscosity value 30 Brix to 35 Brix has been achieved.

(iii) Phase of purification: The purification process is done by mixing the glucose liquid with activated charcoal. Activated charcoal has a very strong adhesion or absorption capability that can bind, coagulate and precipitate inorganic or organic components to free the syrup from unwanted impurities. Purification is done by mixing the liquid glucose with activated charcoal with temperature setting 80 $^{\circ}$ C.

(iv) Filtering stage: Filtering is useful for separating the activated charcoal and components attached to the syrup liquid. This filtration is expected to retain the particles of impurities that have been previously coagulated by the activated char so that the resulting liquid is clear yellow.

(v) Evaporation stage: Evaporation is performed on reactors previously used for liquefaction and saccharification processes. The process is carried out at 70 °C. Evaporation aims to concentrate glucose from (30 to 35) brix to (43 to 80) brix to obtain clear colored yellowish sugar. The type of cassava used cultivated the same is white cassava. Use of the α -amylase enzyme is purchased from the same manufacturer and the residual α -amylase enzyme already used is stored in the refrigerator to avoid damaging the α -amylase enzyme.

2.2 Methods of research

The research methods include the following steps:

(i) Designing of liquid sugar processing and manufacture of liquid processing tool based on milrocontroller which includes mechanical and electronic design both hardware and software. The block diagram of the control system of the system can be shown in figure 2.



Figure 2. Block diagram.

The power supply from the 220 VAC PLN line is connected to the power supply circuit to supply the agitating motor and generate 5 VDC and 12 VDC voltages to supply the driver, microcontroller and sensor circuit. Valve type used is a DC valve. The source of this power supply is not directly connected to the valve and the water pump but through the driver circuit first. This driver circuit will get an input signal from a microcontroller to set the load or valve closing time.

In making this liquid sugar using a voltage source from the power supply to supply the voltage in the microcontroller circuit, the driver circuit and solenoid valve, microcontroller will process the value readings to the pH and temperature sensors by performing some calculations [4, 5]. The relay is used to drive the solenoid valve to open the gas line of the LPG gas tank for heating.

The 2nd International Conference on Natural Resources and Life Sciences	(NRLS)	IOP Publishing
IOP Conf. Series: Earth and Environmental Science 293 (2019) 012019	doi:10.1088/1755-1	315/293/1/012019

Both sensors are inserted in the liquid sugar solution to measure their pH. The pH probe produces an electrical voltage according to the measured pH. And the temperature sensor produces a binary value according to the measured liquid sugar temperature, the result of the sensor reading value being sent to the LCD. The next process of programming to run the driver circuit in accordance with the value obtained to be able to set the desired temperature. The work flow chart of the system automation system liquefaction and saccharification process in liquid sugar making from cassava is shown in figure 3:

(i) Measurement and testing of tools that have been made to know the performance of the tool as a whole. In the manufacture of liquid sugar using two processes of liquefaction and saccharification. Liquefaction the cassava (starch) solution is heated at 95 °C to 105 °C and maintained for 1 h, whereas at pH it is maintained at acidity of pH 6.2 to pH 6.4 to maintain pH with the addition of lime. Liquefaction process can be stopped if the solution is completely liquid and clear brown. After 1 h of warming, it is continued in the process of saccharification, after cooling to 60 °C, enzyme amiloglukosidase added 1 mL kg⁻¹ of starch. The pH is adjusted to pH 4 to pH 4.6 by adding HCl solution into the starch solution. The saccharification process takes up to 76 h.

(ii) Conducting several experiments to collect data and analyze it. After doing data analysis, drawing conclusions from the overall experimental results.



Figure 3. Flow chart diagrams.

3. Results and discussion

The results obtained in this study can be explained as follows:

3.1. Construction tank

The process of liquefaction and saccharification process of liquid sugar consists of two parts, namely stirrer and heater and panel box control system. The stirrer and heating apparatus used for the framework is made of iron and tank and stirrer stainless steel stirrer with tank size of 80 L with the diameter of 47 cm and height 47 cm shown in figure 4.



Figure 4. Mixing and heating equipment.

The panel box in the design of liquid sugar processing contains several components such as power supply, microcontroller, LCD, and driver circuit to run solenoid valve, lighter, peristaltic pump and push button start or stop. The panel box has a dimension $30 \text{ cm} \times 40 \text{ cm} \times 10 \text{ cm}$ can be shown in figure 5.



Figure 5. Panel box.

The circuit of electronics as the overall controller of the system is composed of a series of modules PT 100, Series of pH Meter Modules.

3.1.1. The PT100 Module circuit. MAX31865 module is a device of Pt100 to get the desired temperature value. Use amplifiers designed to read low resistance, and have amplifiers that can adjust and compensate for output output. Schematic Pt100 module can be shown in figure 6. Temperature measurement is done by inserting the PT100 sensor into a pan containing water that has been heated and then lowered in temperature to obtain a number of samples from 30 °C to 120 °C. The average percentage error between the measurement thermometer and PT 100 sensor is 0.86 %.



Figure 6. Schematic module Pt100.

3.1.2. Network module pH meter [6]. The output of the pH sensor is a voltage value in which the value of this voltage must be measured with an amplifier circuit in the form of OP-Amp whose task is to amplify the voltage so that the voltage can enter the ADC work area. Figure 7 is an image of the pH sensor Logo module.



Figure 7. Schematic module logo pH sensor.

The 2nd International Conference on Natural Resources and Life Sciences	(NRLS)	IOP Publishing
IOP Conf. Series: Earth and Environmental Science 293 (2019) 012019	doi:10.1088/1755-13	315/293/1/012019

Reinforcement of RPS (Signal Conditioning Circuit) is three times that the output voltage does not exceed the input of the maximum ADC voltage allowed. The trick is to set the potentiometer R5 so that the gain does not exceed the specified ADC limit. In conducting the preparation of liquid heating sugar required at certain desired temperatures and pH, Arduino Uno microcontroller first initialises the port. PT100 and Probe pH meter will read the temperature and pH values and value read will be displayed on the LCD. If the start or stop button is pressed (0) then the Solenoid Valve will live to open the LPG gas tap, then the lighter will live for 3 s to spark fire, and the agitator to stir the processed liquid sugar. When the tool does the heating and stirring process, the PT100 Sensor will keep reading the temperature change. If the temperature value has reached the limit then the Solenoid Valve with a comparative buffer solution and a fixed temperature which is the room temperature around 24 °C to 26 °C to be able to determine the level of accuracy of the tool designed. The average percentage error between the measurement of pH meter ATC and pH meter analog is 1.09 %.

3.2. System testing based on ATmega328 AVR series microcontroller [7]

System testing aims to find out the automation works well and to know the length of time the process occurs. Measurements are made by mixing starch flour with water at a rate of 5 Kg in volume 15 L. The power consumption is 4.5 W when the system standby, and needs 311 W by activating the stirrer motor, gas valve and lighter. The collection of temperature and pH, power and currents data is carried out every 5 min or there is a change in the material being tested. From the data obtained, the testing time is 134 min. The results and measurements system based on microcontroller performance can be shown in Table 1.

Timer	Temperature (°C)	pН	Stove	Watt (W)	Current (A)	Information
-	-	-	-	285	3.01	Mixed starch
-	32	7.05	ON	266	2.97	Heating
-	54	7.19	ON	261	2.94	Heating
-	84	7.11	ON	304	3.12	Gelatinization
1	100	6.99	OFF	292	3.04	Enzym on
3	96	6.89	OFF	277	3.00	The material melts
5	96	6.85	ON	277	2.99	Light brown liquid
7	98	6.78	ON	275	2.98	Dark brown liquid
10	101	6.77	OFF	260	2.93	Dark brown liquid
15	98	6.70	ON	269	2.95	
-	-	-	-	-	-	Time break
15	79	6.98	ON	280	2.99	
20	83	6.93	ON	272	2.94	
25	89	6.85	ON	266	2.94	
30	93	6.85	ON	252	2.89	
35	96	6.83	ON	261	2.94	
40	92	6.83	ON	263	2.94	Small fire
45	90	6.81	ON	267	2.94	Small fire
50	92	6.78	ON	262	2.93	Small fire
55	95	6.77	ON	258	2.90	Small fire
59	97	6.76	ON	260	2.94	Small fire
-	-	-	-	4.5	-	Process done

Table 1. Data of temperature dan pH based on microcontroller performance.

3.3. Data retrieval and data analysis

In the process of making liquid sugar by using enzymatic methods, there are two processes that use the enzyme that is the process of liquefaction and saccharification process [3, 8–10]. Liquefaction process uses α -amylase enzyme while saccharification process using glucoamylase enzyme. The volume of

The 2nd International Conference on Natural Resources and Life Sciences	(NRLS)	IOP Publishing
IOP Conf. Series: Earth and Environmental Science 293 (2019) 012019	doi:10.1088/1755-13	15/293/1/012019

cassava solution used is 1 L. The solution is heated by hot plate while continuous stirring using stirring rod. This stirring aims to keep the solution homogeneous. The heating temperature of the solution is about (70 to 80) °C up formed gel on the solution of cassava starch. This process is called gelatinization process. The gelatinization process takes about 20 min.

The gel-formed solution then measured its pH using a universal pH paper. The pH of the solution is adjusted to a pH of 5.5 using vinegar. After reaching pH 5.5, the gel-ready solution is ready to enter the next process that is the liquefaction process. The gel-formed solution is then added α -amylase enzyme according to the predetermined level while remaining heated by hot plate with continuous stirring until ± 100 °C. In this process, the temperature of the solution is kept constant until the process is complete. Factors influential in the process of liquefaction which made free variable is the amount of enzyme α -amylase and time of the process of liquefaction. The level of the independent variables can be shown in table 2. The number of α -amylase enzymes in ml units and the liquefaction process time in minutes.

 Table 2. Free variable level full factorial design method.

Independent variabel			
Number of enzymes α-amylase (X1), mL	0.8	1.2	1.6
Processing time (X2), min	30	45	60

The combination of the factors and the level of the liquidity process and the result of the measurement of the reduction sugar content from the resulting liquidity process can be shown in table 3. The result of the experiments of the liquefaction process is then measured by reducing the reducing sugar by using the spectrophotometer in the laboratory. The experiment was done by replicating two times.

Number of enzymes α-amylase (mL)	Processing time (X2), min	Replication 1	Replication 2	Mean
0.8	30	33.3233	43.499	38.4112
0.8	45	52.68	63.166	57.923
0.8	60	74.3467	80.87	77.6084
1.2	30	167.467	158.568	163.017
1.2	45	102.433	107.925	105.179
1.2	60	78.1067	61.626	69.8664
1.6	30	28.4733	31.553	30.0132
1.6	45	50.3833	53.2112	51.7973
1.6	60	307.8	301.75	304.775

Table 3. Factor and experiment level combinations, measurement results and mean calculations.

In combination of 1.6 mL of α -amylase enzyme and 60 min of liquefaction process, the average value of reducing sugar content was 304.775 g L⁻¹. From table 2, there is a significant effect of the combination between the enzyme α -amylase factor and the time factor of the liquefaction process on the reduction sugar content.

4. Conclusion

Based on preliminary experiments, there is a significant influence of the combination of α -amylase enzyme factor factor and liquefaction time factor on reducing sugar content. This is by regulating and maintaining the processing time, temperature (70 to 80) °C by stirring constantly and maintained pH level 5.5. These three parameters are regulated and are being implemented with the design of an industrial liquid sugar processing apparatus. From the liquefaction process carried out by the tool it takes around 134 min with a capacity of 20 L. Consumption of system power when the standby condition is 4.5 W. When the system is working at 311 W by activating the stirrer motor, gas valve and lighter.

References

- [1] Aron H H B S 2016 RI Masih Impor Gula Mentah dari Tiga Negara Ini [RI Still Imports of Raw Sugar from Three Countries] [Online] from <u>https://finance.detik.com/industri/d-3248169/ri-masih-impor-gula-mentah-dari-tiga-negara-ini</u> [Accessed December 1, 2018]
- [2] Simanjuntak W, Satria H and Utami N 2014 Production of reducing sugar from cassava solid waste by simultaneous ultrasonication and acid hydrolysis *Indones. J. Chem.* 14(3) 233–38 <u>https://jurnal.ugm.ac.id/ijc/article/view/21233</u>
- [3] Mulyana I J, Santosa L M H and Yuliati 2016 Rancang bangun mesin pembuatan sari tapioka pada pemroses gula cair (sirup glukosa) dari singkong (*Manihot utillissima*) untuk industri kecil menengah [Designing the making machine for tapioca juice in liquid sugar processors (glucose syrup) from cassava (*Manihot utillissima*) for small and medium industries] *Jurnal Teknik Industri* 22(11) 826–33

https://journal.uii.ac.id/jurnal-teknoin/article/view/8937

 [4] Beran J A 2009 Laboratory Manual for Principles of General Chemistry (New Jersey: John Willey & Sons)

https://www.amazon.com/Lab-Manual-Principles-General-Chemistry/dp/0470567929

[5] Laird B B and Chang, R 2009 University Chemistry (Boston: The McGraw-Hill Companies, Inc.)

https://www.worldcat.org/title/university-chemistry/oclc/269444553

- [6] Scholz F 2010 Electroanalytical Methods Guide to Experiments and Applications 2nd ed (Berlin: Springer-Verlag Berlin Heidelberg) https://www.springer.com/gp/book/9783642029141
- [7] Microchip 2018 megaAVR®Data Sheet (Arizona: Microchip Technology Inc.) http://ww1.microchip.com/downloads/en/DeviceDoc/ATmega48A-PA-88A-PA-168A-PA-328-P-DS-DS40002061A.pdf
- [8] Samaranayake M D W, De Silva A B G C J, Fernando W R D, Gunawardhane K V T and Herath H M T 2017 Optimization of liquefaction and saccharification times for laboratory scale production of glucose syrup from cassava starch and scaling up process of optimized conditions at pilot scale *Res. J. Chem. Sci.* 7(7) 16–25 http://www.isca.in/rjcs/Archives/v7/i7/4.% 20ISCA-RJCS-2017-048.php
- [9] Edama N A, Sulaiman A and Rahim S N A 2014 Enzymatic saccharification of Tapioca processing wastes into biosugars through immobilization technology *BRJ* 1 2–6 <u>https://www.biofueljournal.com/article_4745_9805b1af2b58b470230bd9cbe62217ca.pdf</u>
- [10] Suripto, Ma'arif S and Arkeman Y 2013 Pengembangan gula cair berbahan baku ubi kayu sebagai alternatif gula kristal dengan pendekatan sistem inovasi [Development of liquid sugar made from raw cassava as an alternative to crystal sugar with the innovation system approach] *Jurnal Teknik Industri* 3(2) 147–56 [in Bahasa Indonesia] https://www.trijurnal.lemlit.trisakti.ac.id/tekin/article/view/1575