

HYPER-COAGULATION AND THE USED OF ANTICOAGULANT FOR PATIENT WITH COVID-19

by Bernadette Dian Novita Dewi

Submission date: 29-Nov-2021 09:50PM (UTC+0700)

Submission ID: 1715254223

File name: 11-Hyper-coagulation_and_dr.Vita.pdf (865.94K)

Word count: 2307

Character count: 13306

HYPER-COAGULATION AND THE USED OF ANTICOAGULANT FOR PATIENT WITH COVID-19

Alyssa Claudia V.Gunawan ¹⁾, Bernadette Dian Novita ²⁾

ABSTRACT

Coronavirus SARS-CoV-2 (COVID-19) infection has become a global infection with high mortality and morbidity rate. The pathophysiology of COVID-19 has not yet clear but associated with increasing proinflammatory cytokines and chemokines that leads to cytokine storm. Cytokine storm leads to increasing coagulation that defined by increasing D-dimer, Prothrombin Time (PT), and activated partial thromboplastin time (aPTT). The use of anticoagulant in patient with COVID-19 decrease the mortality rate and become one of the recommended therapy.

Keyword: COVID-19, Hyper-coagulation, Anticoagulant

ABSTRAK

Virus korona SARS-CoV-2 (COVID-19) telah menjadi infeksi global dengan angka mortalitas dan morbiditas yang tinggi. Patofisiologi dari COVID-19 ini masih belum jelas dan dikaitkan dengan peningkatan sitokin dan kemokin inflamasi yang menyebabkan badai sitokin. Badai sitokin ini menyebabkan peningkatan koagulasi dan ditandai dengan peningkatan D-dimer, *Prothrombin Time* (PT), dan *activated partial thromboplastin time* (aPTT). Pemberian antikoagulan pada pasien COVID-19 terbukti dapat mengurangi angka mortalitas sehingga menjadi salah satu terapi yang dianjurkan.

Kata Kunci: COVID-19, Hyper-coagulation, Anticoagulan

1) *Alumnus Faculty of Medicine Widya Mandala Catholic University Surabaya*

2) *Department of Pharmacology and Therapy, Faculty of Medicine Widya Mandala Catholic University Surabaya Corresponding : Bernadette Dian Novita Email : novita@ukwms.ac.id*

INTRODUCTION

Coronaviruses (CoVs) are enveloped RNA viruses with club-like spikes, exceptionally large RNA genome, and has different replication strategy compare to other viruses as its characters.

Coronaviruses affects to mammals and poultry including humans and cause many types of sign and symptoms. Types of coronaviruses that have caused endemic to humans are *Severe Acute Respiratory Syndrome coronavirus* (SARS-CoV) and

Middle Eastern Respiratory Syndrome Coronavirus (MERS-CoV).¹

Coronaviruses SARS-CoV-2 is the caused of COVID-19, which has become a global infection with high mortality and morbidity rates. In Indonesia, the first finding happened on March 2, 2020 and increase continuously while spreading

throughout Indonesia with mortality rate around 6,7% of all positive cases.²

COVID-19 could be associated with higher mortality rate in patient with comorbidity as hypertension, diabetes, coronary heart disease, tumour, chronic obstructive pulmonary disease, chronic kidney disease, tuberculosis, asthma, and hepatitis B.³⁻⁵

Table 1 : Reported outcomes among COVID-19 patients of all ages, by hospitalization status, underlying health condition, and risk factor for severe outcome from respiratory infection — United States, February 12–March 28, 2020.⁵

Underlying health condition/Risk factor for severe outcomes from respiratory infection (no., % with condition)	No. (%)			
	Not hospitalized	Hospitalized, non-ICU	ICU admission	Hospitalization status unknown
Total with case report form (N = 74,439)	12,217	5,285	1,069	55,868
Missing or unknown status for all conditions (67,277)	7,074	4,248	612	55,343
Total with completed information (7,162)	5,143	1,037	457	525
One or more conditions (2,692, 37.6%)	1,388 (27)	732 (71)	358 (78)	214 (41)
Diabetes mellitus (784, 10.9%)	331 (6)	251 (24)	148 (32)	54 (10)
Chronic lung disease* (656, 9.2%)	363 (7)	152 (15)	94 (21)	47 (9)
Cardiovascular disease (647, 9.0%)	239 (5)	242 (23)	132 (29)	34 (6)
Immunocompromised condition (264, 3.7%)	141 (3)	63 (6)	41 (9)	19 (4)
Chronic renal disease (213, 3.0%)	51 (1)	95 (9)	56 (12)	11 (2)
Pregnancy (143, 2.0%)	72 (1)	31 (3)	4 (1)	36 (7)
Neurologic disorder, neurodevelopmental, intellectual disability (52, 0.7%)†	17 (0.3)	25 (2)	7 (2)	3 (1)
Chronic liver disease (41, 0.6%)	24 (1)	9 (1)	7 (2)	1 (0.2)
Other chronic disease (1,182, 16.5%) [‡]	583 (11)	359 (35)	170 (37)	70 (13)
Former smoker (165, 2.3%)	80 (2)	45 (4)	33 (7)	7 (1)
Current smoker (96, 1.3%)	61 (1)	22 (2)	5 (1)	8 (2)
None of the above conditions [§] (4,470, 62.4%)	3,755 (73)	305 (29)	99 (22)	311 (59)

US CDC shown that some underlying disease are more at risk to Intensive Care Unit (ICU) admission as diabetes mellitus, cardiovascular disease, chronic lung disease, and chronic kidney disease.³

Patient with COVID-19 frequently show symptoms as fever, cough, sore throat, myalgia, diarrhoea, and severe pneumonia with tachypnoea (≥ 30 breaths per min), oxygen saturation $\leq 93\%$ at rest, or PaO₂/FiO₂ ratio < 300 mm Hg) with

complication that can be associated with higher mortality rate including pneumonia, septic, and Acute Respiratory Distress Syndrome (ARDS).⁴⁻⁹

Pathophysiology of COVID-19

Pathophysiology of COVID-19 related ARDS has similarities to other pneumonia. There is overproduction of early response proinflammatory cytokines including tumour necrosis factor (TNF), interleukin (IL)-6, and IL-1 β that resulted in cytokine storm. Cytokine storm related to

the risk of vascular hyperpermeability, multiorgan failure, and eventually death.¹⁰

Cytokine Storm Mechanism in COVID-19

In-vitro cells study shown that at early stage of COVID-19 there is delayed release of cytokines and chemokines in respiratory epithelial cells, dendritic cells, and macrophages. Later, the cells secrete low levels of antiviral factors interferons (IFNs) and high levels of proinflammatory cytokines (IL-6, IL-1 β , and tumour necrosis factor (TNF) and chemokines (C-C motif chemokine ligand (CCL)-2, CCL-3, and CCL-5). Coronavirus infected airway epithelial cells, THP-1 cells (monocyte cell), human peripheral blood monocyte-derived macrophages, and dendritic cells that resulted in delayed but elevated levels of proinflammatory cytokines and chemokines.¹¹⁻¹³

Patient with severe symptoms had significantly higher serum cytokines and chemokines levels than patient with mild to moderate symptom. This phenomenon are prompt by substantial number of neutrophils and monocytes in lung tissues and peripheral blood and lead to lung pathology.^{14,15}

Patient with COVID-19 shown increasing levels cytokines as IL-1 β , IFN- γ , IP-10, and monocyte chemoattractant protein 1 (MCP-1).⁴ This cytokines trigger

the reaction of T-helper type 1 (TH1) cells.¹⁶ There is also increasing numbers of cytokines as IL-4 and IL-10 which inhibit the inflammatory response that secreted from TH2 cells. The serum levels of cytokines IL-2 and IL-6 in patient with COVID-19 influence the severity of the disease with higher levels mean more critically ill patients.⁷ Some study also shown elevated serum levels of granulocyte colony-stimulating factor, IP-10, MCP-1, macrophage inflammatory protein-1A, and TNF- α in patient with COVID-19 that treated in ICU compared to the one in general unit. The increasing cytokines are positively correlated with disease severity.⁴

More than 70% of COVID-19 related pneumonia patients required mechanical ventilation and around 70% suffered ARDS.¹⁷ In patient with ARDS there are significantly increasing serum levels of cytokine that positively correlated with development and progression of the disease.¹⁸ Increasing cytokine that cause cytokine storm also play a role that determined multiple organs failure.¹⁹

Several therapeutic approaches have been used to deal with cytokine storms in COVID-19 including IFN- λ , corticosteroids, Intravenous Immunoglobulin (IVIG), IL-1 antagonists, IL-6 antagonists, TNF blockers, IFN- $\alpha\beta$, chloroquine, and so on.¹³

Hyper-coagulation in COVID-19

Infection triggers a responses that lead to coagulation. Coagulation itself limits the invasiveness of the pathogen.²⁰ Coagulation pathway leads to overproduction of proinflammatory cytokines. Thrombin promotes clot formation by activating platelets and converting fibrinogen to fibrin. Thrombin also stimulates inflammation through proteinase-activated receptors (PARs). Thrombin production is managed by negative feedback loops and anticoagulants such as antithrombin III, tissue factor pathway inhibitor, and the protein C system.²¹ These three anticoagulant mechanism impaired during the inflammation then imbalance of thrombin production and consumption occurs.¹⁰

The imbalance between procoagulant and anticoagulant induced the development of micro-thrombosis, disseminated intravascular coagulation (DIC), and multiorgan failure.²² In patient with COVID-19, coagulopathy defined as increasing prothrombin time (PT) > 3 seconds, increasing activated partial

thromboplastin time (aPTT) > 5 seconds, and increasing D-dimer > 1.0 mcg/mL as main indicator. Patient COVID-19 with severe pneumonia and associated with increasing mortality shown an elevated D-dimer, increased PT, and elevation in IL-6.²³ Increased IL-6 levels itself correlated with increased fibrinogen levels which increase fibrin and further clot promotion.²⁴

Anticoagulant in Patient with COVID-19

Anticoagulant such as heparin decrease the mortality rate. Patient COVID-19 patient in Wuhan with Sepsis Induces Coagulopathy (SIC) score ≥ 4 or increased D-dimer more than 6 fold considered as more severe.^{22,25} Low molecular weight heparin (LMWH) therapy for at least 7 days show decrease mortality by 20% from 52,4% to 32,8%.²⁵ In New York, the anticoagulant therapy affect to the mortality of patient COVID-19 with 22.5% with median survival of 21 days compared with mortality rate of 22.8% with median survival of 14 days in patient without anticoagulant therapy.²⁶

Table 2. ISTH SIC scoring system¹⁰

Item	Score	Range
Platelet count ($\times 10^9/L$)	1	100-150
	2	<100
PT-INR	1	1.2-1.4
	2	>1.4
SOFA score	1	1
	2	≥ 2
Total score for SIC	≥ 4	

ARDS considered one of many complication of COVID-19 with highest mortality rate. This lung coagulopathy are established by localized tissue factor-mediated thrombin generation and decreased bronchoalveolar plasminogen activator-mediated fibrinolysis.^{28,29} Treatment using LMWH within 7 days onset of ARDS reduce the mortality up to 35% and the risk of 28 days mortality by around 37%.³⁰

Treatment with systemic anticoagulant also shown significantly increased PT, aPTT, lactate dehydrogenase, ferritin, C reactive protein, and D-dimer compared to those who did not receive the treatment. The mortality of patient also associated with the duration of the anticoagulant treatment²⁶. Low molecular heparin considered as a potential and effective anticoagulant in COVID-19 patient with hyper-coagulation.

However, Heparin administration itself come with complication as bleeding events. Major bleeding define as 1) decreasing Haemoglobin (Hb) < 7 g/dL and required blood cell transfusion, 2) given minimal two units red blood cell transfusion within 2 days or 3) diagnose for internal bleeding including intracranial haemorrhage, hematemesis, melena, peptic ulcer with haemorrhage, colon, rectal, or anal haemorrhage, haematuria, ocular

haemorrhage, and acute haemorrhagic gastritis. Among patient with anticoagulant therapy 3% individuals had bleeding event compared with 1,9% patient with bleeding event without anticoagulant therapy.²⁶

REFERENCES

1. Fehr AR, Perlman S. Coronaviruses: An overview of their replication and pathogenesis. *Molecular Biology* vol 1282. 2015. DOI: 10.1007/978-1-4939-2438-7_1
2. Peta Sebaran [Internet]. Gugus Tugas Percepatan Penangan COVID-19; 2020. Available from <https://covid19.go.id/peta-sebaran>
3. Li X, Xu S, Yu M, et al. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *J Allergy Clin Immunology*; April 2020.
4. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497-506
5. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019 – United States, February 12 – March 28. CDC COVID-19 Response

- Team, 2020. *MMWR* April 2020; vol. 69; no. 13.
6. Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020; published online Feb 28. DOI:10.1056/NEJMoa2002032.
 7. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020; 395: 507–13.
 8. Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2020; 41: 145–51
 9. Verity R, Okell L, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet*. Published online March 2020. DOI: 10.1016/S1473-3099(20)30243-7.
 10. Jose RJ. COVID-19 Cytokine Storm: the interplay between inflammation and coagulation. *Lancet*; published online April 2020. DOI: 10.1016/S2213-2600(20)30216-2
 11. Law HKW, Cheung CY, Ng HY, Sia SF, Chan YO, Luk W, et al. Chemokine up-regulation in SARS-coronavirus-infected, monocyte-derived human dendritic cells. *Blood*. 2005;106(7):2366–74 PubMed PMID: 15860669. Epub 2005/04/28.
 12. Cheung CY, Poon LLM, Ng IHY, Luk W, Sia S-F, Wu MHS, et al. Cytokine responses in severe acute respiratory syndrome coronavirus-infected macrophages in vitro: possible relevance to pathogenesis. *J Virol*. 2005;79(12):7819–26 PubMed PMID: 15919935.
 13. Ye Q, Wang B, et al. The pathogenesis and treatment of the ‘Cytokine Storm’ in COVID-19. *Journal of Infection*. Published online March 2020. DOI: 10.1016/j.jinf.2020.03.037
 14. Kim ES, Choe PG, Park WB, Oh HS, Kim EJ, Nam EY, et al. Clinical progression and cytokine profiles of middle east respiratory syndrome coronavirus infection. *J Korean Med Sci*. 2016;31(11):1717–25 PubMed PMID: 27709848.
 15. Ng DL, Al Hosani F, Keating MK, Gerber SI, Jones TL, Metcalfe MG, et al. Clinicopathologic, immunohistochemical, and ultrastructural findings of a fatal

- case of middle east respiratory syndrome coronavirus infection in the United Arab Emirates, April 2014. *Am J Pathol.* 2016;186(3):652–8 PubMed PMID: 26857507. Epub 2016/02/05.
16. Marchingo JM, Sinclair LV, Howden AJM, Cantrell DA. Quantitative analysis of how Myc controls T cell proteomes and metabolic pathways during T cell activation. *eLife.* 2020;9:e53725 2020/02/05
 17. Yang X, Yu Y, Xu J, Shu H, Ja Xia, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respirat Med.* 2020 S2213-600(20)30079-5. PubMed PMID: 32105632.
 18. Parsons PE, Eisner MD, Thompson BT, Matthay MA, Ancukiewicz M, Bernard GR, et al. Lower tidal volume ventilation and plasma cytokine markers of inflammation in patients with acute lung injury. *Critical Care Med.* 2005;33(1):1–232 PubMed PMID: 15644641.
 19. Wang H, Ma S. The cytokine storm and factors determining the sequence and severity of organ dysfunction in multiple organ dysfunction syndrome. *Am J Emerg Med.* 2008;26(6):711–15 PubMed PMID: 18606328.
 20. Esmon CT, Xu J, Lupu F. Innate Immunity and Coagulation. *J Thromb Haemost.* 2011; 9(Suppl 1): 182–188.
 21. José RJ, Williams AE, Chambers RC. Proteinaseactivated receptors in fibroproliferative lung disease. *Thorax.* 2014; 69: 190–92
 22. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost.* 2020; 18: 844–47.
 23. Zhou F, Yu T, et al. Clinical course and risk factors for mortality of adult inpatient with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet.* 2020; 395: 1054-62.
 24. Ranucci M, Balotta A, et al. The procoagulant pattern of patient with COVID-19 acute respiratory distress syndrome. *J Thromb Haemost.* 2020.
 25. Tang N, Bai H, Chen X, et al. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J*

- Thromb Haemost. 2020; DOI: 10.1111/jth.14817.
26. Paranjpe I, Fuster V, et al. Association of Treatment Dose Anticoagulant with In-Hospital Survival Among Hospitalized Patient with COVID-19. JACC. 2020; DOI: 10.1016/j.jacc.2020.05.001.
 27. Iba T, Nisio M, et al. New criteria for sepsis-induced coagulopathy (SIC) following the revised sepsis definition: a retrospective analysis of a nationwide survey. BMJ Open. 2017; 7:e017046. DOI:10.1136/bmjopen-2017-017046.
 28. Ozolina A, Sarkele M, Sabelnikovs O, et al. Activation of coagulation and fibrinolysis in acute respiratory distress syndrome: a prospective pilot study. Front Med. 2016;3:64.

HYPER-COAGULATION AND THE USED OF ANTICOAGULANT FOR PATIENT WITH COVID-19

ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

16%

PUBLICATIONS

13%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Institute of Technology, Nirma University Student Paper	4%
2	Anshika Chauhan, Rajandeep Kaur, Prantar Chakraborti, Arnab Pal. "Silent Hypoxemia" Leads to Vicious Cycle of Infection, Coagulopathy and Cytokine Storm in COVID-19: Can Prophylactic Oxygen Therapy Prevent It?", Indian Journal of Clinical Biochemistry, 2021 Publication	2%
3	www.theijoem.com Internet Source	2%
4	Submitted to University of Keele Student Paper	2%
5	link.springer.com Internet Source	2%
6	Submitted to University of Wolverhampton Student Paper	1%

7	www.ccjm.org Internet Source	1 %
8	www.ncbi.nlm.nih.gov Internet Source	1 %
9	onlinelibrary.wiley.com Internet Source	1 %
10	www.researchsquare.com Internet Source	1 %
11	Shahab Qureshi, Drishti D. Kampani, Tara Ali Hassan Al-Qutbi, Aalya Mohamed, Mubarak Alfaresi. "COVID-19 Observations from Hospitalized Ward Patients in the Northern Emirates: A Practice Only Preached", Cold Spring Harbor Laboratory, 2021 Publication	1 %
12	ima-contentfiles.s3.amazonaws.com Internet Source	1 %
13	www.tandfonline.com Internet Source	1 %
14	www.ashp.org Internet Source	1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On