

# Protective factor evaluation of purslane (*Portulaca grandiflora*) magenta flower variety herbs extract cream formula

*by* Antonius Budiawan

---

**Submission date:** 10-Mar-2025 12:35PM (UTC+0700)

**Submission ID:** 2610389597

**File name:** uation\_P\_grandiflora\_Extract\_Cream\_Formula\_Kirana\_et\_al\_2023.pdf (499.32K)

**Word count:** 4270

**Character count:** 22691



1

## Protective Factor Evaluation of Purslane (*Portulaca grandiflora*) Magenta Flower Variety Herbs Extract Cream Formula

Bida Cincin Kirana\*, Erlien Dwi Cahyani, Antonius Budiawan

Department of Pharmacy Diploma III, Vocational Faculty, Universitas Katolik Widya Mandala Surabaya, Indonesia

\*Corresponding author: [bida.cincin.kirana@ukwms.ac.id](mailto:bida.cincin.kirana@ukwms.ac.id)

Submitted: 26 June 2023

Revised: 10 December 2023

Accepted: 19 December 2023

### Abstract

**Background:** Indonesia is an equatorial country that is rich in sunlight all year. UV light is divided into three wavelength groups: UV-A (320-400nm), UV-B (280-320nm), and UV-C (100-290nm). Intracellular chromophores in skin cell membranes such as riboflavin, porphyrin, nicotinamide, and enzymes will absorb the UV-A light. The UV-B light penetrates the dermis layer and causes DNA structure changes, which lead to wrinkles and a rising risk of skin cancer. Premature skin aging and skin cancer can be prevented with sunscreen preparation containing compounds that can protect the skin from UV radiation. Flavonoid is one of the purslane (*Portulaca grandiflora*) active metabolites that have the potency to be developed as sunscreen. **Objective:** This research aimed to determine the ability of purslane (*Portulaca grandiflora*) magenta flower variety herbs extract cream as a sunscreen as indicated by the %Te, %Tp, and Sun Protective Factor value. **Methods:** This research was an experimental study with various purslane magenta flower variety herbs extract cream formulas that were tested for their %Te, %Tp, and SPF value with a UV-Vis spectrophotometer. **Results:** The sunscreen cream preparation with 2.5% of purslane (*Portulaca grandiflora*) magenta flower variety herb extract had %Te, %Tp, and SPF values of  $25.86 \pm 2.41\%$ ,  $36.05 \pm 2.82\%$ , and  $3.97 \pm 0.35$  respectively. At the same time, preparations with 5% concentration of extract had %Te, %Tp, and SPF values of  $8.23 \pm 0.86\%$ ,  $16.65 \pm 0.92\%$ , and  $8.03 \pm 0.38$ , respectively. **Conclusion:** The sunscreen activity of all extract concentration creams was significantly different compared to the negative control (cream base) in all parameters. Flavonoids are the compounds responsible for the sunscreen activity of purslane extract.

**Keywords:** sun protective factor, purslane, *Portulaca grandiflora*, magenta, cream

2

### How to cite this article:

Kirana, B. C., Cahyani, E. D. & Budiawan, A. (2023). Protective Factor Evaluation of Purslane (*Portulaca grandiflora*) Magenta Flower Variety Herbs Extract Cream Formula. *Jurnal Farmasi dan Ilmu Kefarmasian Indonesia*, 10(3), 379-385. <http://doi.org/10.20473/jfiki.v10i22023.379-385>

## INTRODUCTION

One of the energy sources needed by living creatures is sunlight because of its involvement in every stage of the living process. For example, it is vitamin D generation. Otherwise, sunlight overexposure also has a negative effect.

Sunlight radiation consists of infrared light (wavelength > 760 nm), visible light (400-760 nm), and UV (ultraviolet) light consisting of UV-A (320-400 nm), UV-B (290-320 nm), as well as UV-C (200-290 nm) (Limpiangkarn & Limpiangkarn, 2010). UV-A and UV-B rays are radiation from sunlight that reaches the earth's surfaces and has an impact on the skin (Wang *et al.*, 2008). The UV-A light that reaches the skin surface will be absorbed by intracellular chromophores in cell membranes such as riboflavin, porphyrin, nicotinamide, and enzymes. This causes oxidative stress, whereas Reactive Oxygen Species (ROS) production overwhelms the natural skin antioxidant mechanism, leading to a decrease in collagen production and wrinkle appearance (Graghani *et al.*, 2014). The UV-B light that reaches the skin will penetrate the dermis layer and cause DNA structure changes, which lead to wrinkles and rising skin cancer risk (Matsuda *et al.*, 2013).

Indonesia is a country in the equatorial area with an abundance of sunlight, which leads to high premature skin aging and skin cancer risk. Premature skin aging and skin cancer can be prevented with sunscreen preparation containing compounds that can protect the skin from UV radiation. The ability to protect skin from sunlight exposure is shown by erythema transmission percentage (%Te), pigmentation transmission percentage (%Tp), and Sun Protective Factor (SPF). Commonly, plants rich in flavonoids have a high SPF value because of their chromophore chemical structure capable of absorbing UV light radiation energy (Saewan & Jimtaisong, 2013). Besides that, flavonoids also have antioxidant activity to prevent oxidative stress so that premature skin aging and cancer risk can be prevented (Chen *et al.*, 2012).

Purslane is a plant that has been studied for its health benefits because of its various secondary metabolites. Flavonoid is one of the purslane active metabolites that have the potency to develop as sunscreen. Purslane *Portulaca grandiflora* magenta flower variety has higher levels of flavonoids compared to other varieties, making it suitable to be developed as a sunscreen (Budiawan *et al.*, 2023).

Cream is a topical preparation with a semi-solid emulsion system. This preparation is easy to use and gives a comfortable feeling in its application. In

addition, with the right cream consistency, the extracts in sunscreen can stick long enough and provide sufficient time for the flavonoid compounds to be absorbed by the skin, providing a maximum sun protection effect.

## MATERIALS AND METHODS

### Materials

Fresh purslane (*Portulaca grandiflora*) magenta flower variety herb was obtained from Madiun regency, East Java, Indonesia. Ethanol 96% was used in the extraction process. Ceto stearyl alcohol, stearic acid, cetyl alcohol, methylparaben, propylparaben, tween 80, and aqua destilata were used to make cream base preparation. Magnesium powder, HCl 2N, C<sub>2</sub>H<sub>5</sub>OH, NH<sub>3</sub>, CHCl<sub>3</sub>, Dragendorff, Mayer, Bouchardat, FeCl<sub>3</sub>, H<sub>2</sub>SO<sub>4</sub> concentrate, and methanol pro analysis were used as reagents.

### Tools

Tools used in this experiment were Spectrophotometer UV-Vis (JaSCO V-730), rotary evaporator, water bath, analytical balance, and glassware (pyrex).

### Method

#### Extract preparation

Two hundred grams of dried purslane (*Portulaca grandiflora*) magenta flower variety herb was extracted with maceration method using 500 ml ethanol 96% for five days. The dregs were re-macerated using the same solvent for another five days. The first and second filtrates were then mixed and thickened with a rotary evaporator at 40°C until one-third of the volume was remaining. The thickened filtrates were then dried in the oven at 50°C for 24 hours until thick extract was obtained.

#### Phytochemical screening

##### 1. Flavonoids Identification

Half a gram of purslane (*Portulaca grandiflora*) magenta flower variety extract was dissolved into water and transferred into a test tube. Magnesium metal and five drops of HCl 2N were added into the tube, and then the mixture was heated for 5-10 minutes. After filtration, the filtrate waited until it cooled down and was added with amil alcohol and then shaken hard. The reaction was positive if a red color formed in the amil alcohol layer (Harbone, 1987; Hanani, 2017).

##### 2. Alkaloid Identification

Half a gram of purslane (*Portulaca grandiflora*) magenta flower variety extract was basified with 1 mL of ammonia, then chloroform was added and crushed vigorously. The chloroform liquid was filtered, the

filtrate was placed in a test tube, 2 N HCl was added, the mixture was shaken, and then left to separate. In a separate test tube: Filtrate 1: As much as one drop of Dragendorff reagent solution is dropped into the filtrate, the presence of alkaloids is indicated by the formation of precipitate or turbidity that is coloured brown. Filtrate 2: As much as one drop of Mayer's reagent solution is dripped into the filtrate, the presence of alkaloids is indicated by the formation of a white precipitate or turbidity. Filtrate 3: As a blank or negative control (Harbone, 1987; Hanani, 2017).

### 3. Saponin Identification

Half a gram of purslane (*Portulaca grandiflora*) magenta flower variety extract was put in a test tube, added hot water and cooled, then shaken for 10 seconds. A stable foam will form in less than 10 minutes, 1-10 cm high, and with the addition of 1 drop of HCl the 2N foam was persistence which indicated the presence of saponins (Harbone, 1987; Hanani, 2017).

### 4. Tannin Identification

Half a gram of purslane (*Portulaca grandiflora*) magenta flower variety extract was put into a test tube and reacted with FeCl<sub>3</sub> 1% solution. The extract contains tannins if a green-black or dark-blue color was formed (Harbone, 1987; Hanani, 2017).

### 5. Terpenoid Identification

Half a gram of purslane (*Portulaca grandiflora*) magenta flower variety extract was put into a test tube and added with chloroform and H<sub>2</sub>SO<sub>4</sub> concentrated. The extract would contain terpenoids if a brown colour was formed (Harbone, 1987; Hanani, 2017).

### Cream preparation

The preparation of this research cream was based on the Arisca formula (2018) with minor modifications.

Each material was measured and the oil phase (ceto stearyl alcohol, cetyl alcohol, stearic acid, and propylparaben) was mixed. The water phase (tween 80 and methylparaben) was mixed. Both phases were heated at 80°C with a water bath until dissolved. Cream preparation was done by adding the hot water

phase to the hot oil phase and the mixture stirred at 12500 rpm until a cold cream base formed. The purslane (*Portulaca grandiflora*) magenta flower variety herb extract was added to the cream base and stirred at 20 rpm until a homogeneous cream was formed.

### Sunscreen activity test

The sunscreen activity test was carried out by determining the SPF value in vitro using the UV-Vis spectrophotometry method. The purslane magenta flower variety herbs extract cream was dissolved into methanol for analysis to obtain a 1000 ppm concentration of test solution. After that, the test solution transmission was read at 272.5–372.5 nm wavelength (every 5 nm interval). The amount of erythema flux that was transmitted by the sunscreen agent (E<sub>e</sub>) is calculated by the formula:  $E_e = \Sigma T.Fe$  while the pigmentation flux is calculated by the formula:  $E_p = \Sigma T.Fe$ . %T<sub>e</sub> and %T<sub>p</sub> value calculated by formula %T<sub>e</sub> =  $E_e / \Sigma E_e$  and %T<sub>p</sub> =  $E_p / \Sigma E_p$ , where T = Transmission value, Fe = a constant of flux erythema, Fp = a constant of flux pigmentation (Cumpelick, 1927). SPF value measurement was done by reading the test solution absorbance at 290-320 nm wavelength with 5 nm interval (Mansur *et al.*, 1986; Mishra *et al.*, 2012) (Table 2). SPF was obtained using the formula:

$$SPF = CF \times \sum_{290}^{320} EE \times I \times Abs$$

Where: CF: correction factor, EE: erythema effect spectrum, I: light intensity spectrum, Abs: sample absorbance.

Table 2. Value of EE x I

λ (nm)	EE x I
290	0.015
295	0.0817
300	0.2874
305	0.03278
310	0.1864
315	0.0839
320	0.018

Table 1. Purslane magenta flower variety herb extract cream formula

Ingredients	Negative Control Formula (%)	2,5% Extract Formula (%)	5% Extract Formula (%)
Ceto stearyl alcohol	7	7	7
Stearic acid	7	7	7
Cetyl alcohol	6	6	6
Nipagin	0.15	0.15	0.15
Nipasol	0.05	0.05	0.05
Purslane extract	-	2.5	5
Tween 80	0.5	0.5	0.5
Aqua destilata ad	100	100	100

Table 5. %Te value of cream preparation

Replication	Negative Control (Cream Base)	Value of %Te (%) 2.5% Extract	5% Extract
I	97,56	27,36	8,97
II	90,28	23,08	8,44
III	93,06	27,14	7,29
Mean	93,64 ± 3,67	25,86 ± 2,41	8,23 ± 0,86

#### Data analysis

The %Te, %T<sub>13</sub> and SPF values were analyzed statistically using a one-way ANOVA analytical method with  $\alpha = 0.05$  and followed by a post hoc test.

#### RESULTS AND DISCUSSION

Fresh purslane (*Portulaca grandiflora*) magenta flower variety herb dried and extracted using maceration method until 5.51 gram thick extract was obtained with 10.64% yield (Table 3).

The next step was qualitative phytochemical identification of the obtained extract. The phytochemical screening result is explained in Table 4.

Table 3. Purslane (*P. grandiflora*) magenta flower variety herb extract yield

Simplicia	Powder weight (g)	Extract weight (g)	Yield (%)
Purslane magenta flower variety herb	51.78	5.51	10.64

Table 4. Purslane (*P. grandiflora*) magenta flower variety herb extract phytochemical screening

Test	Reagent	Result
Flavonoid	Mg powder + HCl 2N + C <sub>2</sub> H <sub>5</sub> OH	+
Alkaloid	NH <sub>3</sub> +CHCl <sub>3</sub> +HCl 2N+ Dragendorff/ Mayer/Bouchardat	+
Saponin	Foam test	+
Tannin	FeCl <sub>3</sub>	+
Terpenoid	H <sub>2</sub> SO <sub>4</sub> concentrate + CHCl <sub>3</sub>	+

The purslane extract was used as an active ingredient in cream preparation with 2.5% and 5% concentration variations. Cream preparation was chosen because it has benefits such as being easy to use, comfortable, and easy to wash with water. The cream also has various functions as a drug carrier, skin emollient, and protection from different interferences, including sunlight.

Sunscreen preparation could contain active ingredients in an inorganic compound (reflect UV radiation) and an organic compound (absorb UV radiation). The purpose of sunscreen application is not only to protect the skin from UV rays exposure, which can trigger negative effects, but also it's expected to inhibit ROS formation, which triggers gene mutations, premature aging, and carcinogenic effects in the long term. Therefore, active antioxidant compounds that can help increase the physical activity of sunscreen are needed in sunscreen preparations. Purslane (*Portulaca grandiflora*) magenta flower variety herb extract contains various secondary metabolites and has antioxidant activity (Addor *et al.*, 2022).

The cream sunscreen activity is determined from the percentage of erythema transmission (% Te), percentage of penetration transmission (% Tp), and the Sun Protective Factor (SPF) value (Table 5).

The percent erythema transmission value (%Te) describes the amount of UV rays exposure from the sun that hits the skin after using sunscreen, which causes erythema (redness) on the skin (Chen *et al.*, 2012). The lower the % Te value, the better the sunscreen protection to prevent erythema. Based on Table 5, the highest %

Te was in the negative control (preparation basis), which was 93.64 ± 3.67%, followed by preparations containing 2.5% extract, which was 25.86 ± 2.41%, and the lowest was the preparation with 5% extract, which was 8.23 ± 0.86%. Based on the statistical test results, it showed that there was a significant difference in the % Te value both in the negative control preparations with 2.5% and 5% extract. This indicates that the extract concentration determines the erythema transmission value of sunscreen preparations.

The percentage value of pigmentation transmission (% Tp) describes the amount of exposure to UV rays from the sun that hits the skin after using sunscreen, which causes pigmentation of the skin. As with % Te, the lower % Tp value indicates better protection of sunscreen against pigmentation on the skin.

Table 6. %Tp value of cream preparation

Replication	Negative Control (Cream Base)	Value of %Tp (%) 2.5% Ekstrak	5% Extract
I	98.29	36.69	17.46
II	91.49	32.96	16.83
III	95.66	38.49	15.65
Mean	95.15 ± 3.43	36.05 ± 2.82	16.65 ± 0.92

Table 7. SPF value of cream preparation

Replication	Negative Control (Cream Base)	Value of SPF 2.5% Ekstrak	5% Extract
I	0.18	3.84	7.83
II	0.16	3.71	8.47
III	0.19	4.37	7.79
Mean	0.18 ± 0.02	3.97 ± 0.35	8.03 ± 0.38

Based on Table 6, shows that the highest % Tp was in the negative control (preparation basis), which was  $95.15 \pm 3.43$ , followed by preparations containing 2.5% extract, which was  $36.05 \pm 2.82$ , and the lowest is a preparation with 5% extract, namely  $16.65 \pm 0.92$ . Based on the results of statistical tests, there was a significant difference in % Tp value in the negative control, preparations with 2.5%, and 5% extract with a  $p > 0.05$ . This shows that the extract concentration in preparation determines the pigmentation transmission value of the sunscreen preparation.

The SPF value states how many times the skin's natural resistance is multiplied so that it is safe in the sun without experiencing sunburn<sup>16</sup>. Based on the test result, the SPF value also shows an increasing trend with increasing extract concentration in the preparation. Based on Table 7, it can be seen that the lowest SPF value was in the negative control (preparation base), which was  $0.18 \pm 0.02$ , followed by preparations containing 2.5% extract, which was  $3.97 \pm 0.35$ , and those having the highest was preparation with 5% extract, namely  $8.03 \pm 0.38$ . Based on the results of statistical tests, it showed that there was a significant difference in the SPF value in the negative control, preparations with extracts of 2.5%, and 5% with  $p > 0.05$ . This shows that the extract concentration determines the SPF value of the sunscreen preparation.

Based on these results, a cream preparation containing 5% purslane herb extract had a higher % Te and % Tp value compared to preparations containing 2.5% purslane herb extract. At the % Tp value, both preparations showed the maximum protection, which was in the sunblock category, while based on % Te, the protection was still in the tanning and suntan categories (Kasitowati *et al.*, 2021). To obtain better protection

based on the % Te value, a higher extract content is needed in the preparation.





Figure 1. Absorbance spectra at 290-400 nm wavelength of cream preparation

The potential<sup>13</sup> for sunscreen activity can be expressed through the value of the Sun Protective Factor (SPF). The SPF value is determined from the absorption results of the preparation at UV wavelengths between 290 – 400 nm (Figure 1). Based on the research results, it was obtained that the SPF values of cream preparations containing 2.5% and 5% purslane herb extract were  $3.97 \pm 0.35$  and  $8.03 \pm 0.38$ , respectively. Both of these SPF values are less than 12, which is in the minimal protection category. This SPF value is not optimal for providing protection because, with an SPF value of less than 15, protection is only given for 1.5 hours (Buso *et al.*<sup>18</sup>, 2017).

Statistical tests were carried out to determine whether there were differences in sunscreen activity in each preparation. Based on statistical tests using one-way ANOVA, a significance value of  $<0.05$  was obtained for both the %Te, %Tp, and SPF values. This means that the purslane herb extract concentration in preparation determines the sunscreen activity. The

higher the extract concentration in the preparation, the better the sunscreen protection will be.



**Figure 2.** Structure of flavonoid A: rutin, B: quercetin (Ganeshpurkar *et al.*, 2017; El-Saber Batiha *et al.*, 2020)

Flavonoids are compounds contained in purslane herb extracts that are thought to have sunscreen activity (Cahyani *et al.*, 2022). Flavonoids can absorb energy at UV light wavelengths because they have covalent bonds in their structure. Flavonoids can absorb energy at a wavelength of 240-290 nm, and flavonoids that have conjugated covalent bonds can absorb at a wavelength of up to 550 nm (He *et al.*, 2021). The Magenta flower variety purslane herb contains the and most significant amount of flavonoids compared to other variants. Therefore, this study used extracts from the purslane herb with the magenta flower variant. Flavonoids contained in purslane (*Portulaca grandiflora*) herb are rutin, quercetin, and isoquercetin compounds (Husein *et al.*, 2021). Flavonoids have at least two aromatic rings with a basic carbon skeleton consisting of 15 C atoms forming C6-C3-C6 (Figure 2) (Julianto, 2019).

## CONCLUSION

The sunscreen cream preparation with 2.5% of purslane (*Portulaca grandiflora*) magenta flower variety herb extract had %Te, %Tp, and SPF values of  $25.86 \pm 2.41\%$ ,  $36.05 \pm 2.82\%$ , and  $3.97 \pm 0.35$  respectively. In comparison, preparations with 5% concentration of extract had %Te, %Tp, and SPF values of  $8.23 \pm 0.86\%$ ,  $16.65 \pm 0.92\%$ , and  $8.03 \pm 0.38$ , respectively. The sunscreen activity of all extract concentration creams was significantly different compared to the negative control (cream base) in all parameters. Flavonoids are the compounds responsible for the sunscreen activity of purslane (*Portulaca grandiflora*) magenta flower variety herb extract.

## AUTHOR CONTRIBUTIONS

Conceptualization, B. C. K., E. D. C., A. B.; Methodology, B. C. K., E. D. C.; Software, E. D. C., A. B.; Validation, B. C. K., E. D. C.; Formal Analysis, B.

C. K., A. B.; Investigation, E. D. C., A. B.; Resources, B. C. K.; Data Curation, B. C. K.; Writing - Original Draft, B. C. K., E. D. C., A. B.; Writing - Review & Editing, A. B.; Visualization, E. D. C.; Supervision, B. C. K.; Project Administration, B. C. K.; Funding Acquisition, B. C. K.

## FUNDING STATEMENT

This research did not receive any specific grant from funding agencies in the public, commercial, or not for profit sectors.

## CONFLICT OF INTEREST

The authors declared no conflict of interest.

## REFERENCES

- Addor, F. A. S. anna, Barcaui, C. B., Gomes, E. E., Lupi, O., Marçon, C. R., & Miot, H. A. (2022). Sunscreen lotions in the dermatological prescription: review of concepts and controversies. *Anais Brasileiros de Dermatologia*; 97(2); 204–222. <https://doi.org/10.1016/j.abd.2021.05.012>
- Arisca, H. (2018). Uji Aktivitas Antioksidan Ekstrak Angkak dan Formulasinya dalam Sediaan Krim yang Diuji Daya Penetrasi dan Keamanan pada Kelinci. *Skripsi*. Universitas Setia Budi Surakarta.
- Budiawan, A., Purwanto, A., Puradewa, L., Cahyani, E. D., & Purwaningsih, C. E. (2023). Wound healing activity and flavonoid contents of purslane (*Portulaca grandiflora*) of various varieties. *RSC Advances*; 13; 9871–9877. <https://doi.org/10.1039/D3RA00868A>
- Buso, P., Radice, M., Baldisserotto, A., Manfredini, S., & Vertuani, S. (2017). Guideline for The Development of Herbal-Based Sunscreen. IntechOpen; doi: 10.5772/intechopen.72712
- Cahyani, E. D., Budiawan, A., & Puradewa, L. (2022). Sunscreen activity of soursop seeds extract. *Strada Journal of Pharmacy*; 4(1); 23-26
- Chen, L., Hu, J. Y., & Wang, S. Q. (2012). The role of antioxidants in photoprotection: A critical review. *Journal of the American Academy of Dermatology*; 67(5); 1013–1024. <https://doi.org/10.1016/j.jaad.2012.02.009>
- Cumpelick, B. (1927). Analytical Procedures and Evaluation of Sunscreens. *Journal of the Society of Cosmetics Chemists*; 23(6); 333–345.
- El-Saber Batiha, G., Beshbishy, A. M., Ikram, M., Mulla, Z. S., Abd El-Hack, M. E., Taha, A. E., Algammal, A. M., & Ali Elewa, Y. H. (2020). The

- pharmacological activity, biochemical properties, and pharmacokinetics of the major natural polyphenolic flavonoid: Quercetin. *Foods*; 9(3). <https://doi.org/10.3390/foods9030374>
- Ganeshpurkar, A., & Saluja, A. K. (2017). The Pharmacological Potential of Rutin. *Saudi Pharmaceutical Journal*; 25(2); 149–164. <https://doi.org/10.1016/j.jsps.2016.04.025>
- Gragnani, A., Cornick, S. Mac, Chominski, V., Ribeiro de Noronha, S. M., Alves Corrêa de Noronha, S. A., & Ferreira, L. M. (2014). Review of Major Theories of Skin Aging. *Advances in Aging Research*; 03(04); 265–284. <https://doi.org/10.4236/aar.2014.34036>
- Hanani, E. (2017). *Analisis Fitokimia*. Jakarta: Buku Kedokteran EGC.
- Harbone, J. (1987). *Metode Fitokimia Edisi Kedua. Padmawinata K. Soediro I. penerjemah. Terjemahan dari Phytochemical Methods*. Bandung: ITB.
- He, H., Li, A., Li, S., Tang, J., Li, L., & Xiong, L. (2021). Natural components in sunscreens: Topical formulations with sun protection factor (SPF). *Biomedicine and Pharmacotherapy*; 134(December 2020). <https://doi.org/10.1016/j.biopha.2020.111161>
- Husein, S. G., Sundalian, M., & Husna, N. (2021). Review: Component Analysis of Purslanes Chemicals Compound (*Portulaca oleraceae* L. and *Portulaca grandiflora* Hook.). *Jurnal Sains dan Kesehatan*; 3(2); 317–327.
- Julianto, T. S. (2019). *Fitokimia: Tinjauan Metabolit Sekunder dan Skrining Fitokimia*. Yogyakarta: Universitas Islam Indonesia.
- Kasitowati, R., Wahyudi, A., Asmara, R., Aliviyantri, D., Iranawati, F., Panjaitan, M. A. P., Pratiwi, D. C., & Arsad, S. (2021). Identification photoprotective activity of marine seaweed: *Eucheuma sp.* *International Conference on Biotechnology and Food Sciences*; 679; 1–5. <https://doi.org/10.1088/1755-1315/679/1/012014>
- Limpiangkarn, W., & Limpiangkarn, W. (2010). Special article: Photo-aging: A literature review. *Journal of the Medical Association of Thailand*; 93(6); 753–757.
- Mansur, J. S., Breder, M. N., Mansur, M. C., & Azulay, R. D. (1986). Determination of Sun Protective Factor by Spectrophotometry. *An Bras Dermatol*; 61; 121–124.
- Matsuda, M., Hoshino, T., Yamakawa, N., Tahara, K., Adachi, H., Sobue, G., Maji, D., Ihn, H., & Mizushima, T. (2013). Suppression of UV-induced wrinkle formation by induction of HSP70 expression in mice. *Journal of Investigative Dermatology*; 133(4); 919–928. <https://doi.org/10.1038/jid.2012.383>
- Mishra, A. K., Mishra, A., & Chattopadhyay, P. (2012). Assessment of in vitro sun protection factor of *Calendula officinalis* L. (*asteraceae*) essential oil formulation. *Journal of Young Pharmacists*; 4(1); 17–21. <https://doi.org/10.4103/0975-1483.93575>
- Saewan, N., & Jimtaisong, A. (2013). Photoprotection of natural flavonoids. *Journal of Applied Pharmaceutical Science*; 3(9); 129–141. <https://doi.org/10.7324/JAPS.2013.3923>
- Wang, S.Q., Stanfield, M.S., & Osterwalder, U. (2008). In Vitro Assessment of UV A Protection by Populer Sunscreen Available in the United States. *Journal of America Dermatology*; 59; 934–42

# Protective Factor Evaluation P grandiflora Extract Cream

## Formula\_Kirana et al\_2023

### ORIGINALITY REPORT

18%

SIMILARITY INDEX

11%

INTERNET SOURCES

16%

PUBLICATIONS

4%

STUDENT PAPERS

### PRIMARY SOURCES

1

[jurnal.jomparnd.com](http://jurnal.jomparnd.com)

Internet Source

3%

2

Bianca Levie Tania, Rini Dwiastuti, Agatha Budi Susiana Lestari, Dewi Setyaningsih.

"Sunscreen Cream Formulation of Noni Leaf Extract (Morinda citrifolia L.) with Emulsifier Combination of Tween 80 and Lecithin", JURNAL FARMASI DAN ILMU KEFARMASIAN INDONESIA, 2022

Publication

2%

3

Antonius Budiawan, Agus Purwanto, Levi Puradewa, Erlien Dwi Cahyani, Christiana Endang Purwaningsih. " Wound healing activity and flavonoid contents of purslane ( ) of various varieties ", RSC Advances, 2023

Publication

2%

4

Submitted to Sriwijaya University

Student Paper

2%

5

Mujaddid Idulhaq, Fajar Baskoro Gardjito, Ibnu Yudistiro. "DISTAL TIBIAL RECONSTRUCTION WITH INTRAMEDULLARY NAIL COMBINED WITH T-PLATE FOLLOWING EN-BLOC-EXCISION OF GIANT CELL TUMOR OF DISTAL TIBIA: CASE-REPORT", (JOINTS) Journal Orthopaedi and Traumatology Surabaya, 2021

Publication

1%

6	J Rohmah, F E Wulandari, C S Rini. " In Vitro: Sunscreen Activity of Red Lettuce Extract ( var. L.) ", IOP Conference Series: Earth and Environmental Science, 2020 Publication	1 %
7	Syamsu Nur, Nursamsiar Nursamsiar, Muhammad Aswad, Aprilia Ester Eunike Tumigolung, Risfah Yulianti, Asril Burhan. "Screening Bioactivity of Kersen Fruits (Muntingia calabura L.) as a Sunscreens Candidate", Jurnal Farmasi Galenika (Galenika Journal of Pharmacy) (e-Journal), 2021 Publication	1 %
8	<a href="http://jurnal.ugm.ac.id">jurnal.ugm.ac.id</a> Internet Source	1 %
9	<a href="http://nexusacademicpublishers.com">nexusacademicpublishers.com</a> Internet Source	1 %
10	Sofia Fatmawati, Fitria Nugrahaeni, Fith Khaira Nursal, Ana Fitriana. "Sunscreen Factor Formulation and Test of Gel Preparations of 70% Ethanol Extract on Arabica Coffee Leaf (Coffea arabica L.)", IOP Conference Series: Earth and Environmental Science, 2022 Publication	1 %
11	Endang Sri Sunarsih, Ebta Narasukma Anggraeny, Patricia Sanggita Listyoputri Wibowo, Novi Elisa. "Phytochemical Screening and Antioxidant Activity of Strawberry Juice (Fragaria ananassa Duchesse) Against Urem Level, Creatinin, and Enzyme Catalase Activity In Isoniazid-Induced Wistar Male Rats", STRADA Jurnal Ilmiah Kesehatan, 2020 Publication	1 %
12	Rahma Syafira, Santi Perawati, Medi Andriani. "Pengaruh Pemberian Ekstrak Buah	1 %

Semangkuk (*Scaphium affine* (Mast.) Pierre)  
terhadap Jumlah Eritrosit dan Leukosit pada  
Mencit Putih Jantan (*Mus musculus*)",  
PHARMACY: Jurnal Farmasi Indonesia  
(Pharmaceutical Journal of Indonesia), 2022

Publication

13

Qonitah Fardiyah, Taslim Ersam, Suyanta,  
Agus Slamet, Suprpto, Fredy Kurniawan.

"New potential and characterization of  
*Andrographis paniculata* L. Ness plant  
extracts as photoprotective agent", *Arabian  
Journal of Chemistry*, 2020

Publication

1 %

14

Mary M. Skrypek, Bruce C. Bostrom, Anne E.  
Bendel. "Medical Cannabis Certification in a  
Large Pediatric Oncology Center", *Children*,  
2019

Publication

1 %

15

Maulita Indrisari, Sartini Sartini, Upik A.  
Miskad, Khairuddin Djawad, Karlina Amir  
Tahir, Nurkhairi Nurkhairi, Lukman Muslimin.  
"Photoprotective and Inhibitory Activity of  
Tyrosinase in Extract and Fractions of  
*Terminalia catappa* L.", *Open Access  
Macedonian Journal of Medical Sciences*, 2021

Publication

1 %

16

Submitted to Miami University of Ohio

Student Paper

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On

# Protective Factor Evaluation P grandiflora Extract Cream

## Formula\_Kirana et al\_2023

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7