

Exploring the Mood-Boosting Potential of Bananas: A Comprehensive Review

by Lannie Hadisoewignyo

Submission date: 17-Jul-2025 04:26PM (UTC+0700)

Submission ID: 2716285454

File name: he_Mood-Boosting_Potential_of_Bananas_A_Comprehensive_Review.pdf (862.76K)

Word count: 10455

Character count: 58459

Exploring the Mood-Boosting Potential of Bananas: A Comprehensive Review

Ivonne Soeliono^{1,2}, Ika Puspitasari^{3,4*}, Andayana Puspitasari Gani⁵ and Lannie Hadisoewignyo⁶

1. Doctoral Program, Faculty of Pharmacy, Gadjah Mada University, Jalan Sekip Utara, Yogyakarta, Special Region of Yogyakarta, Indonesia.
2. Department of Clinical and Community Pharmacy, Faculty of Pharmacy, Widya Mandala Surabaya Catholic University, Jalan Raya Kalisari Selatan No. 1, Pakuwon City, Surabaya, East Java, Indonesia.
3. Department of Pharmacology, Faculty of Pharmacy, Gadjah Mada University, Jalan Sekip Utara, Yogyakarta, Special Region of Yogyakarta, Indonesia.
4. Academic Hospital of Gadjah Mada University, Jalan Kabupaten (Lingkar Utara), Special Region of Yogyakarta, Yogyakarta, Indonesia
5. Department of Pharmaceutical Biology, Faculty of Pharmacy, Gadjah Mada University, Faculty of Pharmacy, Jalan Sekip Utara, Yogyakarta, Special Region of Yogyakarta, Indonesia.
6. Department of Pharmaceutical Science and Technology, Faculty of Pharmacy, Widya Mandala Surabaya Catholic University, Jalan Raya Kalisari Selatan No. 1, Pakuwon City, Surabaya, East Java, Indonesia

Article Info

Submitted: 20-07-2023

Revised: 30-09-2024

Accepted: 08-10-2024

*Corresponding author
Ika Puspitasari

Email:
ika_tunggul@ugm.ac.id

ABSTRACT

Food, like bananas, can be used as depression therapy. Bananas are known for their high tryptophan content. Tryptophan is the precursor of the neurotransmitter serotonin, which is closely related to the pathophysiology of anxiety and depression. This comprehensive review examines the available research on the link between banana consumption and mood. Screening of original articles or reviews was done on the Google Scholar database, Science Direct, MedLine (PubMed), Scielo, and Proquest with the keywords "banana," "Musa," "depression," and "anxiety" published no later than March 31, 2023. From 8,469 articles found, 18 studies were selected. From the experimental trial articles obtained (9 studies), most reported positive antidepressant and anxiolytic effects, but not all were significant. The results of the three clinical trials were also inconsistent. Bananas have anxiolytic and antidepressant properties, but the scientific literature on their efficacy is limited. Therefore, more evidence is needed to justify its use as a primary or adjunctive treatment for depression and anxiety.

Keywords: anxiety, banana, depression, mood disorders

INTRODUCTION

Mood disorders, such as major depression disorder, are commonly treated with selective serotonin reuptake inhibitors (SSRIs), although many adverse effects are found (Edinoff et al., 2021). Currently, SSRIs have been the first line of treatment for anxiety disorders (National Institute for Health and Care Excellence (NICE), 2011). However, SSRIs have various limitations, such as the time lag until the onset of pharmacological effects (a few hours before therapeutic effects, within two weeks for sub-chronic therapy) (Blier, 2001). Hence, the patient's mood may deteriorate (Asher et al., 2017). Patients treated with SSRIs were also anxious about drug dependency and stigma (Dome et al., 2019). Sometimes,

conventional treatment also does not give the expected effect.

A more comprehensive understanding of the pathophysiology of depression has encouraged using compounds with anti-inflammatory, antioxidant, and neuroprotective effects as an alternative therapy or supportive therapy for depression (Dobrek, 2023; Jembrek et al., 2023; Moragrega & Ríos, 2021; Tayab et al., 2022; Winiarska-mieczan et al., 2023). Explanations of depression theories other than from the perspective of monoamine deficiency have been supported by evidence showing an increase in inflammatory biomarkers and oxidative stress and a decrease in the capacity of antioxidants in the body, such as vitamins A and C, glutathione

peroxidase, superoxide dismutase, and catalase (Winiarska-mieczan et al., 2023). The presence of inflammation is related to decreased levels of tryptophan in the body because inflammation causes hyperactivation of the enzyme that converts tryptophan into N-formylkynurenine (indoleamine-2,3-dioxygenase), which is then metabolized into kynurenine (Paul et al., 2022). It has also been proven that tryptophan concentrations are inversely proportional to pro-inflammatory cytokine levels (Dowlati et al., 2010).

Natural products are used to treat depression because they are believed to have lower toxicity, better compatibility, better cultural acceptability, and are more affordable and accessible to get (Al-Harrasi et al., 2023; Dwyer et al., 2011; Peng et al., 2022). One of the natural products that can be used to treat depression is food. Foods can affect human mood significantly. Healthy foods such as vegetables and fruit can improve mood and mental health (AlAmmar et al., 2020; Gibson et al., 2020; Grases et al., 2019; Martínez-González & Sánchez-Villegas, 2016). Since the monoamine theory is believed to contribute to the pathophysiology of depression and anxiety disorders, therapy to increase serotonin levels in patients with both illnesses makes sense.

Previous studies have shown that eating a diet high in tryptophan can help treat depression (Chojnacki et al., 2023; Lindseth et al., 2015; Parker & Brotchie, 2011). Tryptophan is an essential amino acid obtained from food. However, only a small amount is stored in the body, which is critical for human physiological functions (Friedman, 2018; Haleem, 2023; Modoux et al., 2021). Tryptophan is a precursor in the production of serotonin. This neurotransmitter plays an essential role in the pathophysiology of depression and anxiety (Davidson et al., 2022; Kałużna-Czaplińska et al., 2019; Roth et al., 2021; Torrente et al., 2012; Waløen et al., 2017). The daily need for tryptophan is around 4-5 mg/kg of adult body weight (Institute of Medicine, 2005; Joint WHO/FAO/UNU Expert Consultation, 2007). This need can be met by consuming tryptophan-rich foods such as meats, milk, wheat bread, chocolate, tuna, cheese, peanuts, oats, dried prunes, bananas, and apples (Ramakrishna et al., 2011; Richard et al., 2009).

Even though the amount of tryptophan in bananas may not be as high as in other types of food, the ratio of tryptophan to other amino acids is relatively high compared to other foods mentioned previously (Richard et al., 2009). The high ratio of

tryptophan to other amino acids is important because, as the only precursor of serotonin, up to 95% of tryptophan distributed throughout the body is bound to albumin (Hood et al., 2005).

Free tryptophan can penetrate the BBB when approaching the blood-brain barrier (BBB) (Baier et al., 2022). Tryptophan competes with other large neutral amino acids such as histidine, isoleucine, methionine, phenylalanine, threonine, tyrosine, and valine for the cerebrovascular transport system (Lucini et al., 1996). However, tryptophan has a higher affinity for that transport system because of its sizeable hydrophobic side chain (Smith, 1988). Therefore, the bioavailability of tryptophan is expressed in the ratio of tryptophan to other amino acids that compete for the same transporter (Lucini et al., 1996). Higher concentration of tryptophan available for serotonin synthesis can be achieved by increasing the amount of tryptophan or decreasing the amount of other competing amino acids.

Banana (*Musa* sp.) is an herbaceous fruit plant from Southeast Asia that has spread to Africa, South America, and Central America (De Langhe et al., 2009; Elayabalan et al., 2017; Perrier et al., 2011). Bananas are a popular fruit cultivated and consumed in Indonesia because they are cheap but nutritious. Moreover, based on the 2022 gross production value, Indonesia was ranked number one in global banana production (FAO, 2022).

Banana has high tryptophan content among other fruits (26.15 ± 0.37 µg/g) (Islam et al., 2016). A previous research reported a higher content of tryptophan (1000 µg/g) in banana peel powder (Soeliono et al., 2023). Another study showed increased serotonin in human platelets after banana administration (Xiao et al., 1998). Providing unripe banana pulp powder for 36 days can significantly increase serotonin levels in the brains of Swiss albino mice (Erigbali et al., 2017). Increased serotonin expression also occurred after administration of bananas in the form of banana flakes (5%; 3 weeks) and banana peel extract (16 mg/kg; 3 days) (Fatchurohmah et al., 2019; Meliala et al., 2020).

However, the antianxiety and antidepressant effects of bananas are still debatable. Understanding the potential mood-enhancing properties of bananas can provide valuable insight into nutritional strategies to improve mental well-being. Therefore, this comprehensive review examines the available research on the relationship between banana consumption and mood. By integrating and

assessing available data, this review aims to show how bananas may alter mood and emotional well-being.

METHODS

This study investigates the depression or anxiety effect of bananas. The following databases were screened: Google Scholar, Science Direct, MedLine (PubMed), Scielo, and Proquest. The main keywords searched include "banana," "Musa," "depression," and "anxiety." Each keyword was searched in the article title. The articles collected were imported to Mendeley software (Elsevier, USA) to remove duplicates. The inclusion criteria are all studies investigating the antianxiety and antidepressants of bananas, published no later than March 31, 2023; written in English or Indonesian; available in full text; and original article or review. The abstracts of all articles were screened, and the selected articles' full text was retrieved. All data were tabulated and analyzed in Microsoft Excel 2021 (Microsoft Corporation, USA).

Table I. Search strategy

Search strategy			
PubMed (MEDLINE)			
#1	"Musa"[Mesh]	AND	"Anti-Anxiety Agents"[Mesh] → 5 results
#2	"Musa"[Mesh]	AND	"Antidepressive Agents"[Mesh] → 4 results
Google Scholar			
#1	(Musa OR Banana OR pisang) AND (anxiety OR depression OR cemas OR depresi) → 8.440 results		
Science Direct			
Title, abstract, keywords: (Banana OR musa) AND (anxiety OR depression) → 17 results			
Scielo			
#1	(banana OR musa OR plátano) AND (depression OR depresión) → 2 results		
#2	(banana OR musa OR plátano) AND (anxiety OR ansiedad) → 1 result		

RESULT AND DISCUSSION

Characteristics of the articles

Eight thousand four hundred sixty-nine articles were screened through all databases (Table I). After reading the title and abstract, 8.451 studies were excluded. With careful analysis, 18 studies were selected (nine experimental studies (Table II), three clinical studies (Table III), and six review articles. Most of the articles retrieved

were experimental studies (47%) from India (Figure 1).

It is well-known that the location of plant growth affects the pharmacological effects (Li et al., 2010). The pharmacological effects of plants are highly dependent on the content and production of secondary metabolites. Secondary metabolites are not directly involved in growth and reproduction processes. Factors that influence the production of secondary metabolites include harvest time, season, soil type, nutrient supply, altitude, geographical location, maturity level, genotype or cultivar, biotic and abiotic stress (such as temperature, drought, salinity, and light intensity), as well as fruit maturity and post-harvest handling and processing (Bachheti et al., 2021; Vu et al., 2018).

The ripeness level of banana peel affects the type of secondary metabolites that can be extracted. Previous research on *Musa paradisiaca* L. proved that the ethyl acetate extract of unripe banana peel does not contain tannin and alkaloids; meanwhile, the results were positive in the ripe banana peel. Likewise, in the ethanol extract, the results of the tannin content test were positive and saponin negative; in the unripe peel, the test results were the opposite (Widoyanti et al., 2023). Some studies show higher antioxidant compound content in green bananas than in ripe bananas, but some studies report the opposite (Fatemeh et al., 2012; Sundaram et al., 2011; Vu et al., 2019). When the green color of the banana peel changed to yellow, the chlorophyll content degraded by around 90%; meanwhile, the carotenoid increased by 50%, and the flavonoid increased by 27% (Vu et al., 2019).

Unfortunately, although the level of banana ripeness determines the content of secondary metabolites that impact pharmacological effects, the preclinical and clinical studies cited in this review did not detail the level of ripeness of the banana samples used. Soltani et al. (2011) mentioned that banana ripeness levels are generally expressed in 7 stages based on color comparison with a graphic scale. In stage 1, the banana peel is entirely green. In stage 2, the banana peel is green with a hint of yellow. In stage 3, the banana peel is greener than yellow. In stage 4, the banana peel is more yellow than green. In stage 5, the banana skin is yellow with a hint of green. In stage 6, the entire banana skin is yellow. Finally, in stage 7, the whole banana skin is yellow with brown spots.

Table II. Preclinical studies of bananas as an antidepressant and antianxiety

References	Banana Species	Developmental Stage	Dosage	Laboratory studies	Result
Parle and Malik (2010)	<i>Musa paradisiaca</i> from Hissar	Not mentioned	3 grams/day of banana paste (5,10, 20%) p.o. for 15 days	• FST and TST • MAO-A, MAO-B, and MDA levels in the brain	<ul style="list-style-type: none"> • There was a significant reduction in the duration of immobility in both tests ($p<0.01$), and the effect is comparable with fluoxetine and imipramine. • Combining BFP with prazosin, baclofen, and p-CFA significantly decreased the effect of BFP ($p<0.01$ and $p<0.05$). • BFP significantly reduced brain MAO-A, MAO-B, and MDA ($p<0.01$) → antioxidant effect.
Hallikeri <i>et al.</i> (2012)	<i>Musa paradisiaca</i>	Unripe	95% alcohol extract 500 mg/kg p.o. for five days	Anti-punishment and anti-frustration (Skinner Box)	The banana peel extract increased anti-punishment and anti-frustration activity compared to the control.
Samad <i>et al.</i> (2017)	<i>Musa sapientum</i> from Punjab	Not mentioned	Banana pulp paste 600 mg/kg p.o. and 70% acetone peel extract 400 mg/kg p.o.	MWM, LDA, EPM, FST	<ul style="list-style-type: none"> • Anxiety test: increasing time spent in open arm in EPM (peel > pulp); increasing time spent in lightbox in LDA (peel > pulp). • Depression test: decreasing immobility time in FST (peel ~ pulp). • Memory test: decreasing time to reach the platform in short- and long-term memory (peel ~ pulp).
Reddy <i>et al.</i> (2017)	<i>Musa sapientum</i>	Not mentioned	Aqueous extract of stem 25, 50, 100 mg/kg p.o.	EPM, OFT	The dose of 100 mg/kg was most consistent in showing the difference in the EPM test parameter with control, and it had a comparable effect with diazepam on locomotor activity.
Salako (2019)	<i>Musa sapientum</i> Linn. from Papa Alanto	Not mentioned	Aqueous extract of leaves 50, 100, 200 mg/kg p.o.	FST, TST, EPM, hole board (mice)	<ul style="list-style-type: none"> • There was a reduction in immobility duration with the highest reduction in the 200 mg/kg group (79.6% in FST, 66.9% in TST) • There was no difference in open-arm exploration in EPM • There was no difference in head dips of the hole board test • Combination with sulpiride/prazosin/ metergoline → ↑ immobility • Combination with yohimbine → ↓ immobility

References	Banana Species	Developmental Stage	Dosage	Laboratory studies	Result
Meliola (2020)	<i>Musa balbisiana</i> <i>Colla</i> (kepok banana) from Sleman	Not mentioned	Peel flakes diet 5%, 10%, 20% TST (immobility time for 6 minutes)	Rats	There was a significant difference in diabetic + 20% flakes with diabetic control. The effect was higher with a higher concentration.
Samad et al. (2020)	<i>Musa sapientum</i>	Not mentioned	Fruit paste 600 mg/kg/ OFT day for two weeks		<ul style="list-style-type: none"> The exploratory activity was increased along with the dose. The time spent in a lightbox and open arm, as well as immobility time and memory, were increased in the unstressed and stressed groups. Administration of BPF inhibited noise stress-induced behavioral deficits (anxiolytic and antidepressant effects) and improved cognitive abilities. The exploratory activity was increased along with the BPE dose. Time spent in a lightbox and open arms increases in unstressed, stressed animals treated with BPE. Immobility time was decreased in unstressed, stressed animals treated with BPE. It had decreased in time to reach the platform in short- and long-term memory in groups treated with BPE.
Samad et al. (2021)	<i>Musa sapientum</i>	Not mentioned	70% acetone extract of banana peel (BPE) 400 mg/kg p.o. 14 days	<ul style="list-style-type: none"> Open field activity Home cage activity LDA EPM FST MWM MDA SOD, CAT, GPx 	<ul style="list-style-type: none"> MDA level was decreased in the treated group. BPE increased the activity of SOD, CAT, and GPx. The rats fed a control meal supplemented with 30% banana peel floss (BPF) had a substantially reduced escape delay and spent less time in the target quadrant than the other groups. They decreased in immobility time. The cortisol level did not decrease with BPF administration. Meanwhile, the serotonin level was increased in the control meal supplemented with 30% and 60% BPF.
Meliola et al. (2021)	<i>Musa balbisiana</i> <i>Colla</i>	Not mentioned	Banana peel floss (BPF) as diet supplementation (15%, 30%, or 60%) <i>ad libitum</i>	<ul style="list-style-type: none"> CMS + MWM CMS + TST Cortisol + serotonin 	

BPF: banana fruit paste; BPE: banana peel floss; CAT: catalase; CMS: chronic mild stress, EPM: elevated plus maze, FST: forced swim test, GPx: glutathione peroxidase; LDA: light dark activity; MAO: monoamine oxidase, MDA: malondialdehyde, MWM: Morris water maze, OFT: open field test, p-CPA: para chlorophenyl alanine, TST: tail suspension test, SOD: superoxide dismutase.

Table III. Clinical studies of bananas as an antidepressant and antianxiety

Reference	Randomized	Population	Intervention	Control	Outcome
Satyarini <i>et al.</i> (2020)	No	60 subjects Female, schizophrenic anxiety Moderate-severe With antipsychotic drugs Without anxiolytics, chronic illness, mental retardation	<ul style="list-style-type: none"> Banana type: <i>Musa paradisiaca</i> sp. (ambon banana) \pm 130 gram/fruit Treatment groups (12 people/group): <ul style="list-style-type: none"> - New patient two bananas per day - New patient three bananas per day - Old patient two bananas per day - Old patient three bananas per day Duration: 14 days (New patient: first hospitalized with a medical diagnosis of schizophrenia and had never received antipsychotic treatment. Old patient: had been hospitalized with a medical diagnosis of schizophrenia and had received antipsychotic treatment.) 	Without banana	Hamilton Anxiety Rating Scale (HARS) scores were decreased in all groups after 7- and 14-day interventions. HARS scores were highest in the control group. There was no difference between 2 and 3 bananas/day.
Ji <i>et al.</i> (2020)	Not mentioned	24,673 participants (13,327 males, 42.3 \pm 12.0 years; females, 40.6 \pm 11.7 years) This cohort includes nearly all occupations and retired persons over the age of 18 who are living in residential communities of Tianjin	No intervention No asking the mean frequency of banana consumption in the last month: Group 1: rarely (reference); Group 2: <1 time/week; Group 3: 1-3 times/week; Group 4: \geq 4 times/week.	Rarely eat a banana	The Chinese version of the Self-Rating Depression Scale (SDS) Prevalence of depression: male 16.1%, female 18.4% Male: Group 1: OR 0.86 Group 2: OR 0.76 Group 3: OR 0.97 Female: Group 1: OR 1.11 Group 2: OR 0.99 Group 3: OR 1.22
Kobayashi <i>et al.</i> (2022)	Yes	20 participants	13 subjects: 2 bananas daily for two weeks	No banana intake	50% of people who ate bananas had a decrease in intestinal environmental markers, stress markers, and autonomic nervous system activity markers \rightarrow banana intake for two weeks improved the intestinal environment that also increased the parasympathetic activity, and provided stress release




Figure 1. Countries of studies




Figure 2. Banana species (A) and banana part (B) used in studies

Most articles use forced swim, tail suspension, elevated plus maze, and open field tests (Table II). The most studied species were *Musa paradisiaca* and *Musa sapientum* (Figure 2A). Four studies reported antidepressant effects, three

The properties of banana

This study reviews the antidepressant and antianxiety effects of bananas. Bananas contain carbohydrates, protein, vitamins, and minerals (Netshiheni et al., 2020; Sidhu & Zafar, 2018). Raw and mature bananas contain bioactive compounds such as phenolic compounds, steroids, biogenic amine, carotenoids, and phytosterols that have a positive effect on health, especially as antioxidants (Afzal et al., 2022; EA et al., 2019; Lopes et al., 2020; Mengstu et al., 2021; Netshiheni et al., 2020; Prasad

reported antianxiety effects, and five reported both effects. Both antianxiety and antidepressants were reported positive, but not all were significant. Most studies used banana fruit or peel (Figure 2B).

& Das, 2024; Qamar & Shaikh, 2018; Singh et al., 2016).

The phenolic compounds in bananas are mainly flavonoids, which have antioxidant effects based on their ability to scavenge oxygen free radicals and inhibit oxidative enzymes that cause the emergence of reactive oxygen species (Dong et al., 2016). The flavonoid content of bananas ranged from 0.40-7.45 quercetin equivalent per 100-gram dry weight (C. V. Borges et al., 2014). The antioxidant effect rises as the banana ripens (Prasad & Das, 2024).

Moreover, banana fruit and peel are rich in carotenoids and vitamins A and C, contributing to antioxidant activity (Pereira & Maraschin, 2015; Toh et al., 2016). Carotenoids (provitamin A) are precursors of vitamin A that the mammalian body cannot synthesize (Hammond & Renzi, 2013). The banana's yellow color is a good predictor of its carotenoid content (Englberger et al., 2006). The banana's carotenoid content varies with cultivars and ripening stage; some reported total carotenoids as high as $6405.03 \pm 217.06 \mu\text{g}/100 \text{ g d.w}$ (C. V. Borges et al., 2019; Chandra et al., 2020).

Furthermore, Bananas contain phytosterols, such as fatty acids, sterols, and steryl esters, which are known to reduce cholesterol levels in the blood and can increase cardiovascular (CVD) risk (Sheng et al., 2017; Vilela et al., 2014). FDA has stated that for lowering CVD risk, the amount of plant sterol and stanol esters are minimally 1.3 and 3.4 g per day ("Food Labeling: Health Claims; Plant Sterol/Stanol Esters and Coronary Heart Disease. Food and Drug Administration, HHS. Interim Final Rule," 2000). The lipophilic extract of banana contains 11.1-28.0% sterol, some of which are campesterol, stigmasterol, and β -sitosterol (Mondal et al., 2021; Vilela et al., 2014).

Banana fruit and peel also contain bioactive amines such as serotonin and dopamine, contributing to antioxidant activity (González-Montelongo et al., 2010; Pereira & Maraschin, 2015a). Moreover, banana contains tryptophan, a protein that will be converted into serotonin inside the body; thus, eating a banana will relax, improve mood, and increase happiness (Kumar et al., 2012; Rajesh, 2017).

Bananas are high in nutrients, and various parts of the banana plant have shown efficacy as a medicinal plant. Banana flowers cure dysentery, ulcers, and bronchitis (Kumar et al., 2012). Banana sap contains astringent chemical compounds and treats hysteria, epilepsy, leprosy, fever, dysentery and diarrhea, bleeding, hemorrhoids, and insect bites (Kumar et al., 2012). Banana leaves have been developed as dressing for burn patients and skin graft donor areas (Gore & Akolekar, 2003b, 2003a; Guenova et al., 2013). Banana roots have been proven to have androgenic properties that support their traditional use as an aphrodisiac treatment (Yakubu et al., 2013). Traditionally, banana peel ingredients are used to treat burns, anemia, diarrhea, ulcers, inflammation, diabetes, coughs, snakebites, and menorrhagia (Pereira & Maraschin, 2015). Nowadays, banana peel is also used as a teeth-whitening agent, anticancer, antimicrobial,

antidiabetic, anti-ulcerative, analgesic, mosquito bite, and skin wound treatment (Bhavani et al., 2023; Kemiseti et al., 2022). In addition, banana peels show activity to prevent or treat depression (Kemisetti et al., 2022). Based on Iranian and Indian traditional medicine, bananas are prescribed for depression (Pereira & Maraschin, 2015b). The chemical content of banana peels and fruits is primarily carotenoids, phenolic compounds, and biogenic amines (serotonin, norepinephrine, and dopamine) (Pereira & Maraschin, 2015b; Vu et al., 2018; Waalkes et al., 1958).

Tryptophan's role in banana

Tryptophan, as a supplement, has been used in the UK as a monotherapy or adjuvant for depressant patients resistant to treatment (Joint Formulary Committee, 2021). The tryptophan content made bananas a potential treatment for depression. Tryptophan is an essential amino acid required in serotonin production. Serotonin can help a person relax and improve overall mood and happiness (Jyothirmayi & Rao, 2015; Lal et al., 2017; Singh et al., 2016). A survey by MINDS, an Indian NGO, showed that people with depression felt much better after eating a banana (Kumar et al., 2012; Rajesh, 2017).

Tryptophan, an essential amino acid and a precursor to serotonin, has been proven to elevate the mood of healthy individuals (Kikuchi et al., 2021). Tryptophan in the body is converted into serotonin. In humans, acute dietary depletion of tryptophan is related to increased depressive symptoms or increased severity of depression (K. A. Smith et al., 1997). It has been found that patients with depression have a lower tryptophan/neutral amino acid ratio (Manosso et al., 2013).

Tryptophan is transported from blood to the brain by a transport carrier that is not specific for tryptophan but also for other amino acids (Ruddick et al., 2006). After tryptophan has entered the extracellular or cerebrospinal fluid, tryptophan can be uptake into cells and incorporated into several metabolism pathways, one of which is serotonin synthesis (Ruddick et al., 2006). Serotonin is a bioactive compound and one of the neurotransmitters involved in many physiological functions such as cardiovascular, pain, feeding, and reproduction and behavioral functions such as cognition, impulsivity, aggression, and mood (Vadodaria et al., 2018).

Once in the brain, tryptophan is hydroxylated to 5-hydroxytryptophan (5-HTP) by

tryptophan 5-hydroxylase, a step that requires vitamin C as a cofactor. 5-HTP is then decarboxylated to 5-hydroxytryptamine, which requires pyridoxine as a cofactor. Serotonin is then metabolized by monoamine oxidase to 5-hydroxy indole acetic acid. Biotransformation of tryptophan to serotonin also occurs in the gut, platelets, and mast cells (Knapp et al., 2003).

The meta-analysis of 108 studies reported improved depressive symptoms with L-tryptophan or 5-hydroxytryptophan supplementation (Shaw, Turner & Del Mar, 2002). It has been mentioned before that bananas have a high content of tryptophan. Therefore, the consumption of bananas is thought to have the same effect.

Serotonin compound in banana

In addition to containing tryptophan, bananas contain serotonin. Serotonin is a physiologically active amine and neurotransmitter that regulates mammals' mood, sleep, and anxiety (Paredes et al., 2016). In plants, it is synthesized from tryptophan. Tryptophan is first catalyzed into tryptamine and then into serotonin (Ramakrishna et al., 2011). It is known that serotonin, but not tryptophan and 5-HTP, does not cross the blood-brain barrier. However, a previous study reported that the level of mice's brain serotonin was increased after 30 days of banana-feeding (Erigbali et al., 2017). The author hypothesized that the increase of serotonin in the brain was because the precursor, tryptophan, crossed the blood-brain.

Previously, serotonin could be detected in other high quantities in banana plants (*Musa* spp.). *Musa acuminata* fresh fruit contains 9.48 ± 0.09 µg/g serotonin (second highest after kiwi), 26.15 ± 0.37 µg/g tryptophan (second highest after watermelon), and 0.959 ± 0.28 tryptamine (the highest) µg/g fresh fruit (Islam et al., 2016). Previous studies have also shown increased platelet serotonin after administering bananas (Xiao et al., 1998). Another review regarding the serotonin content in *Musa sapientum* peel and pulp reported a higher serotonin content in banana peel, and the serotonin is lower in the ripening stage (Ramakrishna et al., 2011). It was mentioned that serotonin is involved in growth regulation, flowering, xylem sap exudation, ion permeability, and plant morphogenesis (Paredes et al., 2016). Biogenic polyamines and amines also contribute to the antioxidant activity of bananas, where their antioxidant abilities correlate with the number of amine groups. Biogenic amines are aliphatic

organic that play a role in animal, plant, and microorganisms' metabolic and physiological functions. In plants, amines such as serotonin can be protective compounds against insects and fungi. Dopamine and serotonin have antioxidant and anti-inflammatory effects (Lopes et al., 2020).

Antioxidant activity of banana

Banana has antioxidant properties that can counter the stress effect in our body so that sympathetic neuron activation will decrease (Aboul-Enein et al., 2016; Alhabysi et al., 2014; Ehigior et al., 2018; Fidrianny et al., 2014, 2018; Hallikeri et al., 2010; Jami'ah et al., 2018; Lopes et al., 2020; Manzoor & Ahmad, 2021; Marikkar et al., 2016; Mentari et al., 2019; Nifinluri et al., 2019; Pane, 2013; Phuaklee et al., 2012; Pratiwi & Krisbianto, 2019; Rebello et al., 2014; Samad et al., 2018; Schmidt et al., 2015; Sukmawati et al., 2015). Oxidative stress, a damaging effect of free radicals, increases in people with depression. The increased oxidative stress markers are evidence of this (Black et al., 2015). A previous study has mentioned that banana pulp consumption in mice can increase antioxidant enzyme activities while decreasing brain lipid peroxidation (Samad et al., 2020). The content of phenolic compounds in banana peels varies between 4.95 – 47 gallic acid equivalents/g dry matter (GAE/g d.m.), which is 1.5-3 times higher than that found in the fruit flesh. The activity of capturing and reducing the number of free radicals is much higher than in other fruit peels (Sukmawati et al., 2015).

Phenolic compounds are secondary metabolites in high amounts of bananas and provide many benefits, such as preventing cardiovascular disease, cancer, diabetes, and obesity. Phenolic compounds in bananas show antioxidant effects by limiting the production of reactive oxygen species (ROS), scavenging ROS, and activating antioxidant enzymes (Zou et al., 2022).

There are more than 40 phenolic compounds in banana peels that have been identified and can be divided into four groups, namely hydroxycinnamic acid (predominantly ferulic acid), flavonols (predominantly rutin and its conjugates), flavan-3-ol (the most dominant group of phenolic compounds), and catecholamines (Vu et al., 2018). Methanol and acetone extract from the bark of *Musa paradisiaca* L. showed quite a high polyphenol content (17.89 ± 0.16 ; 15.44 ± 0.19 mg/g d.w.) as well as vigorous antioxidant activity (IC_{50} 56.22 ± 8.61 ; IC_{50} 55.45 g/mL) (Aboul-Enein et al., 2016).

Musa acuminata Colla AAA showed high activity in scavenging 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azino-bis(3-ethylbenzothiazoline)-6-sulphonic acid (ABTS) free radicals and in inhibiting lipid peroxidation (González-Montelongo et al., 2010). *Musa* AAA flour also showed high total phenolic content (about 29 mg/g GAE) with very high antioxidant activity (FRAP 14 M/g TE, ABTS 242 M/g, ORAC 436 M/g) (Jami'ah et al., 2018).

Research on the fruit and skin of eight banana cultivars in Malaysia showed difference in total phenolic content and antioxidant activity. The highest total phenolic content (76.37 ± 1.79 mg GAE/g d.w.) was obtained from the fresh fruit of the Raja cultivar, which was freeze-dried. The maximum activity of DPPH (19.39 ± 0.15 mg TE/g d.w.) was achieved by the chloroform extract of dried mas banana peel. Most of the chloroform extracts of dried fruit showed the highest ferric-reducing antioxidant potential (FRAP), especially in the awak cultivar (22.57 ± 0.13 mg TE/g d.w.).

Another study showed a high content of phenolic compounds in milk banana flowers (80.13 ± 4.64 mg GAE/g extract) and high antioxidant activity (ABTS+ 24.73 ± 0.04 , DPPH 25.10 ± 0.15 mol of TE /g extract) and cavendish bananas and salalah bananas (Alhabsyi et al., 2014; Amri & Hossain, 2018; Manzoor & Ahmad, 2021).

Ethanol extract from *Musa sapientum* banana fruit and peel showed vigorous antioxidant activity (IC₅₀ 19.10; 44.07 g/ml) (Dahham et al., 2015). In another study, *Musa sapientum* had the highest phenolic content in the decoction extract of unripe fresh banana peel (117.68 ± 0.19 mg GAE/g) and the best antioxidant activity in 50% ethanol extract of dried raw banana peel (EC₅₀ 7.33 ± 0.55 g/ml) (Phuaklee et al., 2012).

In Indonesia, there have been several studies on the topic of banana antioxidant potential, including tongka langit banana flour (*Musa troglodytarum* L.), which shows a relatively high antioxidant activity (DPPH IC₅₀ 159.41 mg/mL), goroho banana peel extract (*Musa acuminata* L.) (ethanol extract 75.71%, methanol extract 74.29%, acetone extract 73.37%), and agung banana peel extract (DPPH IC₅₀ 684.69 ± 120.73 mg/mL) (Fidrianny et al., 2018; Mentari et al., 2019; Pratiwi & Krisbianto, 2019; Soeliono et al., 2023). Research comparing the antioxidant activity of the skin of various banana cultivars in Indonesia, including raja bulu banana, muli banana, and Ambon banana, using n-hexane, ethyl acetate, or ethanol as solvent reported that the highest DPPH scavenging

capacity was from muli banana ethanol extract (IC₅₀ 4.39 ppm). Meanwhile, in an antioxidant test using ABTS, ethyl acetate extract of ambon lumut banana has the highest capacity (IC₅₀ 1.91 ppm) (Fidrianny et al., 2014).

Based on the evidence presented, bananas, particularly their peels, are rich sources of antioxidants. These compounds, primarily phenolic compounds, are crucial in neutralizing harmful free radicals, thereby mitigating oxidative stress. They can contribute to various health benefits, including antidepressants and antianxiety.

Anti-inflammation activity of banana

Evidence has shown an association between inflammation and depressive symptoms. There was an increasing concentration of pro-inflammatory cytokines and cell-mediated immune activation in the depression pathophysiology. Those inflammatory states are also influenced by lifestyle, such as a diet rich in tryptophan, including bananas (Strasser et al., 2016). Various types of bananas also show effectiveness as an anti-inflammatory, which may be due to the content of polyphenols, alkaloids, and terpenes. Its antioxidant activity and radical scavenging can produce an anti-inflammatory effect, modulation of cellular activity of inflammatory cells, the activity of pro-inflammatory enzymes, production of other pro-inflammatory molecules, and expression of pro-inflammatory genes (Bellik et al., 2013). That is because of an increase in reactive oxygen species (ROS) in inflammatory conditions. During the inflammatory process, mast cells and leukocytes are recruited to the site of cell damage, causing an oxidative burst in which ROS production increases (Reuter et al., 2010). The anti-inflammatory effect test on *M. paradisiaca* stem methanol extract (200; 400 mg/kg, p.o.) showed a significant anti-inflammatory effect (Biswas et al., 2012). Likewise, the anti-inflammatory test results of *M. paradisiaca* bark extract (100, 200, 400 mg/kg) showed significant results, especially at a 400 mg/kg dose compared to positive controls (Pane, 2013). In another study, the ethanol extract and water fraction at a 200 mg/kg dose showed significant inhibition compared to the diclofenac sodium (Rebello et al., 2014). Ethanol extract of kepok banana peel at doses of 75 mg/kg BW, 150 mg/kg BW, and 300 mg/kg BW, as well as ethanol extract of Ambon banana leaf at doses of 500, 750, and 1000 mg/kg BW, were also shown to have anti-inflammatory activity (Ehigiator et al., 2018; Marikkar et al., 2016).

Preclinical and Clinical Studies of Banana as Antianxiety and Antidepressant

In preclinical studies (Table II), bananas showed antianxiety and antidepressant properties. Several studies report positive antianxiety effects. The antianxiety effects are indicated by an increase in time in the open arm on the EPM test and an increase in time on the lightbox in the LDA test (Reddy et al., 2017; Samad et al., 2017, 2021). In addition to the significant EPM or LDA, OFT test results show an increase in exploratory activity after giving bananas (Samad et al., 2020). However, one study reported the adverse antianxiety effects of bananas (Salako et al., 2019).

The results of the antidepressant test were shown by the results of the FST and TST tests in units of immobility time, which showed a significant decrease (Meliala et al., 2021; Parle & Malik, 2010; Salako et al., 2019; Samad et al., 2017, 2018, 2021). A memory test was also carried out with the MWM test, which showed an increase in short and long-term memory (Meliala et al., 2021; Samad et al., 2017, 2021). Although all the review articles in this study supported bananas as antianxiety and antidepressants, the clinical study did not. Some studies reported an increase in memory function and cognitive abilities. The effects were increased in a dose-dependent manner.

Nevertheless, the clinical studies' results were inconsistent (Table III). One clinical study reported decreased anxiety scores, but no difference was found in the intervention of two or three bananas daily (Setyarini & Wasita, 2020). The other study was inconsistent because the reduction in depression prevalence seems not to increase with the number of bananas eaten daily (Ji et al., 2020). However, the current clinical study showed that some people who eat bananas for two weeks have an improved intestinal environment and increased parasympathetic activity (Kobayashi et al., 2022).

Future perspectives for bananas as an antidepressant

Bananas contain both tryptophan and serotonin and can show antioxidant and anti-inflammatory activity. The results of preclinical trials and clinical trials of bananas as antianxiety and antidepressants are still limited in number. Further testing is needed to determine the mechanism of action underlying this effect. These findings can help to better understand the potential of bananas as antianxiety and antidepressants and

the development of bananas as a natural remedy for patients with anxiety or depression.

CONCLUSION

The use of bananas as antianxiety and antidepressant treatment still seems far from expectations. Although sufficient scientific reasons exist to enforce the use of bananas for treatment, the number of preclinical and clinical studies was rare. More clinical studies are needed to support using bananas as a mood booster.

ACKNOWLEDGMENTS

The authors acknowledge the support of Widya Mandala Surabaya Catholic University, which awarded the first author a doctoral scholarship.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

- Aboul-Enein, A. M., Salama, Z. A., Gaafar, A. A., Aly, H. F., Bou-Ellella, F. A., & Ahmed, H. A. (2016). Identification of phenolic compounds from banana peel (*Musa paradisiaca* L.) as antioxidant and antimicrobial agents. *Journal of Chemical and Pharmaceutical Research*, 8(4), 46–55.
- Afzal, M. F., Khalid, W., Akram, S., Khalid, M. A., Zubair, M., Kauser, S., Abdelsamea Mohamedahmed, K., Aziz, A., & Anusha Siddiqui, S. (2022). Bioactive profile and functional food applications of banana in food sectors and health: a review. *International Journal of Food Properties*, 25(1), 2286–2300. <https://doi.org/10.1080/10942912.2022.2130940>
- Al-Harrasi, A., Bhatia, S., Behl, T., Aldawsari, M. F., Kaushik, D., & Chigurupati, S. (2023). Recent advances in natural products science. In *Recent Advances in Natural Products Science* (1st ed.). CRC Press.
- AlAmmar, W. A., Albeesh, F. H., & Khattab, R. Y. (2020). Food and mood: the corresponsive effect. *Current Nutrition Reports*, 9(3), 296–308. <https://doi.org/10.1007/s13668-020-00331-3>
- Alhabsyi, D. F., Suryanto, E., & Wewengkang, D. S. (2014). Aktivitas antioksidan dan tabir surya pada ekstrak kulit buah pisang goroho (*Musa acuminata* L.) [Antioxidant and

- sunscreen activity of goroho banana peel extract (*Musa acuminata* L.]. *Jurnal Ilmiah Farmasi Pharmacol*, 3(2), 107–114.
- Amri, F. S. Al, & Hossain, M. A. (2018). Comparison of total phenols, flavonoids and antioxidant potential of local and imported ripe bananas. *Egyptian Journal of Basic and Applied Sciences*, 5(4), 245–251. <https://doi.org/10.1016/j.ejbas.2018.09.002>
- Asher, G. N., Gerkin, J., & Gaynes, B. N. (2017). Complementary therapies for mental health disorders. *Medical Clinics of North America*, 101(5), 847–864. <https://doi.org/10.1016/j.mcna.2017.04.004>
- Bachheti, A., Deepti, Bachheti, R., & Husen, A. (2021). Medicinal plants and their pharmaceutical properties under adverse environmental conditions. In A. Husen (Ed.), *Harsh Environment and Plant Resilience: Molecular and Functional Aspects* (pp. 458–459). Springer. <https://doi.org/10.1007/978-3-030-65912-7>
- Baier, M. P., Salman, M. G., & Awwad, H. O. (2022). Connections between micronutrients, cerebral monoamines and related brain disorders. In W. Mohamed & T. Yamashita (Eds.), *Role of Micronutrients in Brain Health* (p. 194). Springer Nature Singapore.
- Bellik, Y., Boukraâ, L., Alzahrani, H. A., Bakhotmah, B. A., Abdellah, F., Hammoudi, S. M., & Iguer-Ouada, M. (2013). Molecular mechanism underlying anti-inflammatory and anti-allergic activities of phytochemicals: an update. *Molecules*, 18(1), 322–353. <https://doi.org/10.3390/molecules18010322>
- Bhavani, M., Sonia, M., Deepika, S., & Awuchi, C. G. (2023). Bioactive, antioxidant, industrial, and nutraceutical applications of banana peel. *International Journal of Food Properties*, 26(1), 1277–1289. <https://doi.org/10.1080/10942912.2023.2209701>
- Biswas, C., Basak, D., Chakroverty, R., Banerjee, A., Dey, S., & Mazumder, K. (2012). Effect of methanol extract of *Musa paradisiaca* (linn) stem juice on chemically induced acute inflammation. *International Journal of Pharmacy and Pharmaceutical Sciences*, 4(S5), 148–150.
- Black, C. N., Bot, M., Scheffer, P. G., Cuijpers, P., & Penninx, B. W. J. H. (2015). Is depression associated with increased oxidative stress? A systematic review and meta-analysis. *Psychoneuroendocrinology*, 51, 164–175. <https://doi.org/10.1016/j.PSYNEUEN.2014.09.025>
- Blier, P. (2001). Pharmacology of rapid-onset antidepressant treatment strategies. *J Clin Psychiatry*, 62(suppl 15), 12–17.
- Borges, C. V., Amorim, V. B. D. O., Ramlov, F., Ledo, C. A. D. S., Donato, M., Maraschin, M., & Amorim, E. P. (2014). Characterisation of metabolic profile of banana genotypes, aiming at biofortified *Musa* spp. cultivars. *Food Chemistry*, 145, 496–504. <https://doi.org/10.1016/j.foodchem.2013.08.041>
- Borges, C. V., Minatel, I. O., Amorim, E. P., Belin, M. A. F., Gomez-Gomez, H. A., Correa, C. R., & Lima, G. P. P. (2019). Ripening and cooking processes influence the carotenoid content in bananas and plantains (*Musa* spp.). *Food Research International*, 124, 129–136. <https://doi.org/10.1016/j.foodres.2018.08.022>
- Chandra, R. D., Siswanti, C. A., Prihastyanti, M. N. U., Heriyanto, Limantara, L., & Brotosudarmo, T. H. P. (2020). Evaluating provitamin a carotenoids and polar metabolite compositions during the ripening stages of the agung semeru banana (*Musa paradisiaca* L. aab). *International Journal of Food Science*, 2020. <https://doi.org/10.1155/2020/8503923>
- Chojnacki, C., Gąsiorowska, A., Popławski, T., Konrad, P., Chojnacki, M., Fila, M., & Blasiak, J. (2023). Beneficial effect of increased tryptophan intake on its metabolism and mental state of the elderly. *Nutrients*, 15(847), 1–13.
- Dahham, S. S., Mohamad, T. A., Tabana, Y. M., & Majid, A. (2015). Antioxidant activities and anticancer screening of extracts from banana fruit (*Musa sapientum*). *Academic J Cancer Res*, 8(2), 28–34. <https://doi.org/10.5829/idosi.ajcr.2015.8.2.95162>
- Davidson, M., Rashidi, N., Nurgali, K., & Apostolopoulos, V. (2022). The role of tryptophan metabolites in neuropsychiatric disorders. *International Journal of Molecular Sciences*, 23, 9968. Multidisciplinary Digital Publishing Institute. <https://doi.org/10.3390/ijms23179968>

- De Langhe, E., Vrydaghs, L., De Maret, P., Perrier, X., & Denham, T. (2009). Why bananas matter: an introduction to the history of banana domestication. *Ethnobotany Research and Applications*, 7. <https://doi.org/10.17348/era.7.0.165-177>
- Institute of Medicine. (2005). Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. (2005). In *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients)*. National Academies Press. <https://doi.org/10.17226/10490>
- Dobrek, L. (2023). Depression and its phytopharmacotherapy - a narrative review. *International Journal of Molecular Sciences*, 24, 4772.
- Dome, P., Tombor, L., Lazary, J., Gonda, X., & Rihmer, Z. (2019). Natural health products, dietary minerals and over-the-counter medications as add-on therapies to antidepressants in the treatment of major depressive disorder: a review. *Brain Research Bulletin*, 146(December 2018), 51–78. <https://doi.org/10.1016/j.brainresbull.2018.12.015>
- Dong, C., Hu, H., Hu, Y., & Xie, J. (2016). Metabolism of flavonoids in novel banana germplasm during fruit development. *Frontiers in Plant Science*, 7(AUG2016), 1–10. <https://doi.org/10.3389/fpls.2016.01291>
- Dowlati, Y., Herrmann, N., Swardfager, W., Liu, H., Sham, L., Reim, E. K., & Lanctôt, K. L. (2010). A meta-analysis of cytokines in major depression. *Biological Psychiatry*, 67(5), 446–457. <https://doi.org/10.1016/j.biopsych.2009.09.033>
- Dwyer, A. V., Whitten, D. L., & Hawrelak, J. A. (2011). Herbal medicines, other than St. John's wort, in the treatment of depression: a systematic review. *Alternative Medicine Review*, 16(1), 40–49.
- EA, B., N, S., I, M., & A, R. (2019). Bioactive compounds in peanuts and banana. *Biochemistry & Analytical Biochemistry*, 08(02), 1–4. <https://doi.org/10.35248/2161-1009.19.8.382>
- Edinoff, A. N., Akuly, H. A., Hanna, T. A., Ochoa, C. O., Patti, S. J., Ghaffar, Y. A., Kaye, A. D., Viswanath, O., Urits, I., Boyer, A. G., Cornett, E. M., & Kaye, A. M. (2021). Selective serotonin reuptake inhibitors and adverse effects: a narrative review. *Neurology International*, 13, 387–401.
- Ehigiatior, B. E., Offonry, N. C., Adikwu, E., & Inemesit, B. O. (2018). Safety, anti-inflammatory and analgesic assessments of methanolic extract of *Musa paradisiaca* peel in Sprague Dawley rats. *African Journal of Pharmacology and Therapeutics*, 7(2), 46–52.
- Elayabalan, S., Subramaniam, S., V., & Ashokkumar, K. (2017). An overview on phytochemical composition of banana (*Musa spp.*). *Indian Journal of Natural Sciences*, 7(42), 12408–12419. www.tnsroindia.org.in
- Englberger, L., Wills, R. B. H., Blades, B., Dufficy, L., Daniells, J. W., & Coyne, T. (2006). Carotenoid content and flesh color of selected banana cultivars growing in Australia. *Food and Nutrition Bulletin*, 27(4), 281–291. <https://doi.org/10.1177/156482650602700401>
- Erigbali, P., Kiridi, K., & Ogregade, I. (2017). Effects of diets containing unripe plantain diet on brain serotonin in mice. *African Journal of Biomedical Research*, 20(3), 321–323. <https://doi.org/10.4314/ajbr.v20i3>
- FAO. (2022). *Value of Agricultural Production*. FAOSTAT. <https://www.fao.org/faostat/en/#data>
- Fatchurohmah, W., Meliala, A., & Sulistyoningih, R. C. (2019). Effect of banana peel extract on serotonin immunoreactivity and stool consistency in colon of healthy male Wistar rat. *AIP Conference Proceedings*, 2094(1). <https://doi.org/10.1063/1.5097491>
- Fatemeh, S. R., Saifullah, R., Abbas, F. M. A., & Azhar, M. E. (2012). Total phenolics, flavonoids and antioxidant activity of banana pulp and peel flours: influence of variety and stage of ripeness. *International Food Research Journal*, 19(3), 1041–1046.
- Fidrianny, I., Anggraeni, N. A. S., & Insanu, M. (2018). Antioxidant properties of peels extracts from three varieties of banana (*Musa sp.*) grown in West Java-Indonesia. *International Food Research Journal*, 25(1), 57–64.
- Fidrianny, I., Rizki R, K., & Insanu, M. (2014). *In vitro* antioxidant activities from various extracts of banana peels using ABTS, DPPH assays and correlation with phenolic, flavonoid, carotenoid content. *International Journal of*

- Research in Pharmaceutical Sciences, 6(8), 299–303.
- Food labeling: health claims; plant sterol/stanol esters and coronary heart disease. Food and Drug Administration, HHS. Interim final rule. (2000). *Federal Register*, 65(175), 54686–54739.
- Friedman, M. (2018). Analysis, nutrition, and health benefits of tryptophan. *International Journal of Tryptophan Research*, 11, 1–12. <https://doi.org/10.1177/1178646918802282>
- Gibson, D., Mariska, S., Ingeborg, B., Marjolein, A. B., Erik, V., & Penninx, B. W. J. H. (2020). Association of food groups with depression and anxiety disorders. *European Journal of Nutrition*, 59(2), 767–778. <https://doi.org/10.1007/s00394-019-01943-4>
- González-Montelongo, R., Gloria Lobo, M., & González, M. (2010). Antioxidant activity in banana peel extracts: Testing extraction conditions and related bioactive compounds. *Food Chemistry*, 119(3), 1030–1039. <https://doi.org/10.1016/j.foodchem.2009.08.012>
- Gore, M. A., & Akolekar, D. (2003a). Banana leaf dressing for skin graft donor areas. *Burns*, 29(5), 483–486. [https://doi.org/10.1016/S0305-4179\(03\)00049-4](https://doi.org/10.1016/S0305-4179(03)00049-4)
- Gore, M. A., & Akolekar, D. (2003b). Evaluation of banana leaf dressing for partial thickness burn wounds. *Burns*, 29(5), 487–492. [https://doi.org/10.1016/S0305-4179\(03\)00050-0](https://doi.org/10.1016/S0305-4179(03)00050-0)
- Grases, G., Colom, M. A., Sanchis, P., & Grases, F. (2019). Possible relation between consumption of different food groups and depression. *BMC Psychology*, 7(14), 1–6.
- Guenova, E., Hoetzenecker, W., Kisuze, G., Teske, A., Heeg, P., Voykov, B., Hoetzenecker, K., Schippert, W., & Moehrl, M. (2013). Banana leaves as an alternative wound dressing. *Dermatologic Surgery*, 39(2), 290–297. <https://doi.org/10.1111/dsu.12067>
- Haleem, D. J. (2023). Nutritional importance of tryptophan for improving treatment in depression and diabetes. *Nutrition Reviews*, 81(1), 133–149. <https://doi.org/10.1093/nutrit/nuac042>
- Hallikeri, C. S., Bhoomannavar, V. S., Patil, V. P., Shivkumar, S. I., Alagawadi, K. R., & Hatapakki, B. C. (2010). Anti-inflammatory activity of unripe fruits of *Musa paradisiaca* Linn. *Advances in Pharmacology and Toxicology*, 11(1), 55–58.
- Hammond, B. R., & Renzi, L. M. (2013). Carotenoids. *Advances in Nutrition*, 4(4), 474–476. <https://doi.org/10.3945/an.113.004028>
- Hood, S. D., Bell, C. J., & Nutt, D. J. (2005). Acute tryptophan depletion. Part I: rationale and methodology. *Australian and New Zealand Journal of Psychiatry*, 39, 558–564.
- Islam, J., Shirakawa, H., Nguyen, T. K., Aso, H., & Komai, M. (2016). Simultaneous analysis of serotonin, tryptophan and tryptamine levels in common fresh fruits and vegetables in Japan using fluorescence HPLC. *Food Bioscience*, 13, 56–59. <https://doi.org/10.1016/j.fbio.2015.12.006>
- Jami'ah, S. R., Haya, M., Pusmarani, J., & Nurhikma, E. (2018). Uji aktivitas antioksidan ekstrak metanol kulit pisang raja (*Musa paradisiaca sapientum*) dengan Metode DPPH (2,2-difenil-1-pikrilhidrazil). [Antioxidant activity test of methanol extract of banana peel (*Musa paradisiaca sapientum*) using the DPPH (2,2-diphenyl-1-picrylhydrazyl) method]. *Jurnal Mandala Pharmacoon Indonesia*, 4(1), 33–38. <https://doi.org/10.35311/jmpl.v4i1.22>
- Jembrek, M. J., Oršolić, N., Karlović, D., & Peitl, V. (2023). Flavonols in action: targeting oxidative stress and neuroinflammation in major depressive disorder. *International Journal of Molecular Sciences*, 24, 6888.
- Ji, T., Li, X., Meng, G., Gu, Y., Zhang, Q., Liu, L., Wu, H., Yao, Z., Zhang, S., Wang, Y., Zhang, T., Wang, X., Cao, X., Li, H., Liu, Y., Wang, X., Wang, X., Sun, S., Zhou, M., ... Niu, K. (2020). The association between banana consumption and the depressive symptoms in Chinese general adult population: A cross-sectional study. *Journal of Affective Disorders*, 264(December 2019), 1–6. <https://doi.org/10.1016/j.jad.2019.12.008>
- Joint Formulary Committee. (2021). BNF 81 (British National Formulary) March 2021. In *Bnf* (Issue 81).
- Joint WHO/FAO/UNU Expert Consultation. (2007). Protein and amino acid requirements in human nutrition. In *World Health Organ Tech Rep Ser* (Vol. 935, p. 143).
- Jyothirmayi, N., & Rao, M. (2015). Banana medicinal uses. *Jour of Med Sc & Tech*, 4(2)(November), 152–160.

- Kałużna-Czaplińska, J., Gątarek, P., Chirumbolo, S., Chartrand, M., & Björklund, G. (2019). How important is tryptophan in human health? *Critical Reviews in Food Science and Nutrition*, 59, 72–88. <https://doi.org/10.1080/10408398.2017.1357534>
- Kemisetti, D., Das, R., & Bhattacharya, B. (2022). A Comprehensive review on *Musa paradisiaca* taxonomical, morphological classification and its pharmacological activities. *Journal of Pharmaceutical Negative Results*, 13(February 2023), 737–749. <https://doi.org/10.47750/pnr.2022.13.S10.079>
- Kikuchi, A. M., Tanabe, A., & Iwahori, Y. (2021). A systematic review of the effect of L-tryptophan supplementation on mood and emotional functioning. *Journal of Dietary Supplements*, 18(3), 316–333. <https://doi.org/10.1080/19390211.2020.1746725>
- Knapp, B. M., Cupp, M. J., & Tracy, T. S. (2003). L-Tryptophan. In M. J. Cupp & T. S. Tracy (Eds.), *Dietary Supplements: Toxicology and Clinical Pharmacology* (pp. 341–385). Humana Press. https://doi.org/10.1007/978-1-59259-303-3_23
- Kobayashi, A., Hu, A., Yamaguchi, T., & Tabuchi, M. (2022). Banana intake relieves the stress of daily life in healthy adult volunteers: an open, randomized, parallel-group comparative study. *Journal of Food and Nutrition Research*, 10(12), 841–849. <https://doi.org/10.12691/jfnr-10-12-2>
- Kumar, S. K. P., Bhowmik, D., Duraivel, S., & Umadevi, M. (2012). Traditional and medicinal uses of banana. *Journal of Pharmacognosy and Phytochemistry*, 1(3), 51–63.
- Lal, N., Sahu, N., Shiurkar, G., & Jayswal, D. K. (2017). Banana: awesome fruit crop for society (Review). *The Pharma Innovation Journal*, 6(7), 223–228.
- Li, P., Yi, L., & Liu, H.-J. (2010). Collection and identification of raw herbal materials. In W. H. W. Liu (Ed.), *Traditional Herbal Medicine Research Methods: Identification, Analysis, Bioassay, and Pharmaceutical and Clinical Studies* (p. 27). John Wiley & Sons, Inc. <https://doi.org/10.1002/9780470921340>
- Lindseth, G., Helland, B., & Caspers, J. (2015). Archives of psychiatric nursing the effects of dietary tryptophan on affective disorders. *Archives of Psychiatric Nursing*, 29(2), 102–107. <https://doi.org/10.1016/j.apnu.2014.11.008>
- Lopes, S., Borges, C. V., Sousa Cardoso, S. M., Almeida Pereira da Rocha, M. F., & Maraschin, M. (2020). Banana (*Musa spp.*) as a source of bioactive compounds for health promotion. In *Handbook of Banana Production, Postharvest Science, Processing Technology, and Nutrition* (pp. 227–244). Wiley. <https://doi.org/10.1002/9781119528265.ch12>
- Lucini, V., Lucca, A., Catalano, M., & Smeraldi, E. (1996). Predictive value of tryptophan/large neutral amino acids ratio to antidepressant respons. *Journal of Affective Disorders*, 36(1996), 129–133. [https://doi.org/10.1016/0165-0327\(95\)00071-2](https://doi.org/10.1016/0165-0327(95)00071-2)
- Manosso, L. M., Moretti, M., & Rodrigues, A. L. S. (2013). Nutritional strategies for dealing with depression. *Food and Function*, 4(12), 1776–1793. <https://doi.org/10.1039/C3FO60246j>
- Manzoor, A., & Ahmad, S. (2021). Banana peel: characteristics and consideration of its extract for use in meat products preservation: a review. *ACS Food Science & Technology*, 1(9), 1492–1506. <https://doi.org/10.1021/ACSF00DSCITECH.1C00235>
- Marikkar, J. M. N., Tan, S. J., Salleh, A., Azrina, A., & Shukri, M. A. M. (2016). Evaluation of banana (*Musa sp.*) flowers of selected varieties for their antioxidative and anti-hyperglycemic potentials. *International Food Research Journal*, 23(5), 1988–1995.
- Martínez-González, M. A., & Sánchez-Villegas, A. (2016). Food patterns and the prevention of depression. *Proceedings of the Nutrition Society*, 75, 139–146. <https://doi.org/10.1017/S0029665116000045>
- Meliala, A., Andwi, Y., Sumiwi, A. R. I., Narwidina, P., Sulisty, S. R. I. L., & Setyaningsih, W. (2020). Banana peel flakes alleviate blood glucose and stress in a dose-dependent manner. *International Journal of Pharmacy and Pharmaceutical Sciences*, 12(8), 75–81.
- Meliala, A., Narwidina, P., Supriyanto, I., Sumarno, Y. T., & Pratama, Y. Y. (2021). The role of banana (*Musa balbisiana* Colla) peel floss as

- functional food matrix to alleviate chronic stress. *Macedonian Journal of Medical Sciences*, 10(A), 1608–1616. <https://doi.org/10.3889/oamjms.2022.8432>
- Mengstu, A., Bachheti, A., Abate, L., Bachheti, R. K., & Husen, A. (2021). Health-promoting benefits, value-added products, and other uses of banana. In *Non-Timber Forest Products: Food, Healthcare and Industrial Applications* (pp. 339–364). https://doi.org/10.1007/978-3-030-73077-2_15
- Mentari, A. B., Kholisoh, S. N., Hidayat, T. N., & Hasbullah, U. H. A. (2019). Pembuatan kopi dari kulit pisang kepok (*Musa paradisiaca* Linn) dan aktivitas antioksidannya. [Making coffee from kepok banana peel (*Musa paradisiaca* Linn) and its antioxidant activity] *Jurnal Ilmu Pangan Dan Hasil Pertanian*, 3(1), 94. <https://doi.org/10.26877/jiphp.v3i1.3872>
- Modoux, M., Rolhion, N., Mani, S., & Sokol, H. (2021). Tryptophan metabolism as a pharmacological target. *Trends in Pharmacological Sciences*, 42(1), 60–73. <https://doi.org/10.1016/j.tips.2020.11.006>
- Mondal, A., Banerjee, S., Bose, S., Das, P. P., Sandberg, E. N., Atanasov, A. G., & Bishayee, A. (2021). Cancer preventive and therapeutic potential of banana and its bioactive constituents: a systematic, comprehensive, and mechanistic review. In *Frontiers in Oncology* (Vol. 11). <https://doi.org/10.3389/fonc.2021.697143>
- Moragrega, I., & Ríos, J. L. (2021). Medicinal plants in the treatment of depression: evidence from preclinical studies. *Planta Medica*, 87(09), 656–685. <https://doi.org/10.1055/a-1338-1011>
- National Institute for Health and Care Excellence (NICE). (2011). Generalised anxiety disorder and panic disorder in adults: management. *Nice Guideline, Last updated 15 June 2020*, 1–41. <https://www.nice.org.uk/guidance/cg113>
- Netshiheni, R. K., Omolola, A. O., Anyasi, T. A., & Jideani, A. I. O. (2020). Banana bioactives: absorption, utilisation and health benefits. In *Banana Nutrition - Function and Processing Kinetics*. <https://doi.org/10.5772/intechopen.83369>
- Nifinluri, C. M. B., Datu, O. S., Potalangi, N. O., & Pareta, D. N. (2019). Uji aktivitas anti-inflamasi ekstrak etanol kulit buah pisang kepok *Musa balbisiana* terhadap kaki tikus putih *Rattus novergicus*. [Anti-inflammatory activity test of ethanol extract of kepok banana peel *Musa balbisiana* on the feet of white rats *Rattus novergicus*.] *Jurnal Biofarmasetikal Tropis*, 2(2), 15–22.
- Pane, E. R. (2013). Uji aktivitas senyawa antioksidan dari ekstrak metanol kulit pisang raja (*Musa paradisiaca* Sapientum). [Testing the activity of antioxidant compounds from methanol extract of banana peel (*Musa paradisiaca* Sapientum)]. *Jurnal Kimia VALENSI*, 3(2), 75–80. <https://doi.org/10.15408/jkv.v3i2.502>
- Paredes, S. D., Rancan, L., García, C., Vara, E., & Tresguerres, J. A. F. (2016). Occurrence of serotonin, melatonin, and their derivatives in plants. In G. A. Ravishanker & A. Ramakrishna (Eds.), *Serotonin and Melatonin Their Functional Role in Plants, Food, Phytomedicine, and Human Health* (pp. 15–30). CRC Press.
- Parker, G., & Brotchie, H. (2011). Mood effects of the amino acids tryptophan and tyrosine: “food for thought” III. *Acta Psychiatrica Scandinavica*, 124(6), 417–426. <https://doi.org/10.1111/j.1600-0447.2011.01706.x>
- Parle, M., & Malik, S. (2010). Anti-depressant potential of banana fruit. *International Journal of Medical Sciences*, 2(2), 148–151.
- Paul, E. R., Schwieler, L., Erhardt, S., Boda, S., Trepci, A., Kämpe, R., Asratian, A., Holm, L., Yngve, A., Dantzer, R., Heilig, M., Hamilton, J. P., & Samuelsson, M. (2022). Peripheral and central kynurenine pathway abnormalities in major depression. *Brain, Behavior, and Immunity*, 101(December 2021), 136–145. <https://doi.org/10.1016/j.bbi.2022.01.002>
- Peng, S., Zhou, Y., Lu, M., & Wang, Q. (2022). Review of herbal medicines for the treatment of depression. *Natural Product Communications*, 17(11), 1–17. <https://doi.org/10.1177/1934578X221139082>
- Pereira, A., & Maraschin, M. (2015). Banana (*Musa spp*) from peel to pulp: ethnopharmacology, source of bioactive compounds and its relevance for human health. *Journal of Ethnopharmacology*, 160, 149–163. <https://doi.org/10.1016/j.jep.2014.11.008>
- Perrier, X., De Langhe, E., Donohue, M., Lentfer, C., Vrydaghs, L., Bakry, F., Carreel, F., Hippolyte,

- I., Horry, J. P., Jenny, C., Lebot, V., Risterucci, A. M., Tomekpe, K., Doutrelepont, H., Ball, T., Manwaring, J., De Maret, P., & Denham, T. (2011). Multidisciplinary perspectives on (*Musa spp.*) domestication. *Proceedings of the National Academy of Sciences of the United States of America*, 108(28), 11311–11318.
<https://doi.org/10.1073/PNAS.1102001110>
 8/-DCSUPPLEMENTAL/SFIG05.PDF
- Phuaklee, P., Ruangnoo, S., & Itharat, A. (2012). Anti-inflammatory and antioxidant activities of extracts from *Musa sapientum* peel. *Journal of the Medical Association of Thailand = Chotmaihet Thangphaet*, 95 Suppl 1.
- Prasad, S. S., & Das, U. (2024). Banana waste as a nutraceuticals product. In *Nutraceuticals from Fruit and Vegetable Waste* (pp. 175–194). Wiley.
<https://doi.org/10.1002/9781119803980.ch6>
- Pratiwi, I. Y., & Krisbianto, O. (2019). Kandungan gizi, beta karoten dan antioksidan pada tepung pisang tongka langit (*Musa troglodytarum* L.). [Nutritional content, beta carotene and antioxidants in tongka langit banana flour (*Musa troglodytarum* L.).] *Agritech*, 39(1), 48.
<https://doi.org/10.22146/agritech.32335>
- Qamar, S., & Shaikh, A. (2018). Therapeutic potentials and compositional changes of valuable compounds from banana-a review. *Trends in Food Science and Technology*, 79(june 2017), 1–9.
<https://doi.org/10.1016/j.tifs.2018.06.016>
- Rajesh, N. (2017). Medicinal benefits of *Musa paradisiaca* (Banana). *International Journal of Biology Research*, 2(2), 51–54.
- Ramakrishna, A., Giridhar, P., & Ravishankar, G. A. (2011). Phytoserotonin: a review. In *Plant Signaling and Behavior* (Vol. 6, Issue 6, pp. 800–809).
<https://doi.org/10.4161/psb.6.6.15242>
- Rebello, L. P. G., Ramos, A. M., Pertuzatti, P. B., Barcia, M. T., Castillo-Muñoz, N., & Hermosín-Gutiérrez, I. (2014). Flour of banana (*Musa AAA*) peel as a source of antioxidant phenolic compounds. *Food Research International*, 55, 397–403.
<https://doi.org/10.1016/j.foodres.2013.11.039>
- Reddy, A. J., Dubey, A. K., Handu, S., & Manocha, S. (2017). Effects of *Musa sapientum* stem extract on experimental models of anxiety. *Avicenna Journal of Phytomedicine*, 7(6), 495–501.
- Reuter, S., Gupta, S. C., Chaturvedi, M. M., & Aggarwal, B. B. (2010). Oxidative stress, inflammation, and cancer: how are they linked? *Free Radical Biology and Medicine*, 49(11), 1603–1616.
<https://doi.org/10.1016/j.freeradbiomed.2010.09.006>
- Richard, D. M., Dawes, M. A., Mathias, C. W., Acheson, A., Hill-kapczak, N., & Dougherty, D. M. (2009). L-Tryptophan : basic metabolic functions, behavioral research and therapeutic indications. *International Journal of Tryptophan Research*, 2, 45–60.
- Roth, W., Zadeh, K., Vekariya, R., Ge, Y., & Mohamadzadeh, M. (2021). Tryptophan metabolism and gut-brain homeostasis. *International Journal of Molecular Sciences*, 22(6), 2973.
<https://doi.org/10.3390/ijms22062973>
- Ruddick, J. P., Evans, A. K., Nutt, D. J., Lightman, S. L., Rook, G. A. W., & Lowry, C. A. (2006). Tryptophan metabolism in the central nervous system: medical implications. *Expert Reviews in Molecular Medicine*, 8(20), 1–27.
<https://doi.org/10.1017/S1462399406000068>
- Salako, O. A., Akindele, A. J., Balogun, A. O., & Adeyemi, O. O. (2019). Investigation of antidepressant, anxiolytic and sedative activities of the aqueous leaf extract of *Musa sapientum* Linn (Banana; Musaceae). *Drug Research*, 69(3), 136–143.
<https://doi.org/10.1055/a-0651-7978>
- Samad, N., Khaliq, S., Rehman, M. H. U., Ahmad, S., Alam, M., Mustafa, S., & Azizuddin. (2021). Exposure to noise augments behavioral deficits in mice: protective effect of banana peel extract. *Pakistan Journal of Pharmaceutical Sciences*, 34(6), 2333–2340.
<https://doi.org/10.36721/PJPS.2021.34.6.SUP.2333-2340.1>
- Samad, N., Khaliq, S., Rehman, M. H. U., Ahmad, S., Alam, M., Mustafa, S., Azizuddin, Sari, D. N. R., Rahmi, A., Hardi, N., Hevira, L., Peters, S. L., Biesiekierski, J. R., Yelland, G. W., Muir, J. G., Gibson, P. R., Bravo, R., Ugartemendia, L., Cubero, J., ... Bruins, M. E. (2018). Nutritional tables to improve mood disorders. *Nutrition Research*, 4(2), 1552–1558.
<https://doi.org/10.1016/j.nutres.2012.01.0>

- Samad, N., Muneer, A., Ullah, N., Zaman, A., Ayaz, M. M., & Ahmad, I. (2017). Banana fruit pulp and peel involved in antianxiety and antidepressant effects while invigorate memory performance in male mice: Possible role of potential antioxidants. *Pakistan Journal of Pharmaceutical Sciences*, 30(3), 989–995.
- Samad, N., Yasmin, F., Khaliq, S., Ahmad, S., Azizuddin, & Mustafa, S. (2020). Diet supplements of banana fruit pulp mitigates repeated noise stress induced behavioral deficits and oxidative stress. *Pakistan Journal of Pharmaceutical Sciences*, 33(5), 2293–2300.
<https://doi.org/10.36721/PJPS.2020.33.5.SUP.2293-2300.1>
- Schmidt, M. M., Prestes, R. C., Kubota, E. H., Scapin, G., & Mazutti, M. A. (2015). Evaluation of antioxidant activity of extracts of banana inflorescences (*Musa cavendishii*). *CYTA - Journal of Food*, 13(4), 498–505.
<https://doi.org/10.1080/19476337.2015.1007532>
- Setyarini, D., & Wasita, B. (2020). The effect of giving ambon banana (*Musa paradisiaca* Sp.) to decrease anxiety levels in adult schizophrenia. *Indian Journal of Public Health Research & Development*, 11(7), 1166–1171.
- Shaw, K. A., Turner, J., & Del Mar, C. (2002). Tryptophan and 5-hydroxytryptophan for depression. *Cochrane Database of Systematic Reviews*, 2010(1).
<https://doi.org/10.1002/14651858.CD003198>
- Sheng, Z., Dai, H., Pan, S., Ai, B., Zheng, L., Zheng, X., Prinyawiwatukul, W., & Xu, Z. (2017). Phytosterols in banana (*Musa spp.*) flower inhibit α -glucosidase and α -amylase hydrolyses and glycation reaction. *International Journal of Food Science and Technology*, 52(1), 171–179.
<https://doi.org/10.1111/ijfs.13263>
- Sidhu, J. S., & Zafar, T. A. (2018). Bioactive compounds in banana fruits and their health benefits. *Food Quality and Safety*, 2(4), 183–188.
<https://doi.org/10.1093/fqsafe/fyy019>
- Singh, B., Singh, J. P., Kaur, A., & Singh, N. (2016). Bioactive compounds in banana and their associated health benefits - a review. *Food Chemistry*, 206, 1–11.
<https://doi.org/10.1016/j.foodchem.2016.03.033>
- Smith, K. A., Fairburn, C. G., & Cowen, P. J. (1997). Relapse of depression after rapid depletion of tryptophan. *The Lancet*, 349(9056), 915–919.
[https://doi.org/10.1016/S0140-6736\(96\)07044-4](https://doi.org/10.1016/S0140-6736(96)07044-4)
- Smith, Q. R. (1988). Regulation of amino acid transport at the blood-brain barrier. In G. Huether (Ed.), *Amino Acid Availability and Brain Function in Health and Disease* (p. 58). Springer-Verlag.
- Soeliono, I., Puspitasari, I., Gani, A. P., & Hadisoewignyo, L. (2023). Antioxidant activity and tryptophan content of banana peel. *EnvironmentAsia*, 16(2), 99–108.
<https://doi.org/10.14456/ea.2023.24>
- Soltani, M., Alimardani, R., & Omid, M. (2011). Evaluating banana ripening status from measuring dielectric properties. *Journal of Food Engineering*, 105(4), 625–631.
<https://doi.org/10.1016/j.jfoodeng.2011.03.032>
- Strasser, B., Sperner-Unterwieser, B., Fuchs, D., & Gostner, J. M. (2016). Mechanisms of Inflammation-associated depression: immune influences on tryptophan and phenylalanine metabolisms. In *Brain Imaging in Behavioral Neuroscience* (Issue November 2011, pp. 95–115).
https://doi.org/10.1007/7854_2016_23
- Sukmawati, Yuliet, & Hardani, R. (2015). Uji Aktivitas antiinflamasi ekstrak etanol daun pisang ambon (*Musa paradisiaca* L.) terhadap tikus putih (*Rattus norvegicus* L.) yang diinduksi karagenan. [Anti-inflammatory activity of ethanol extract of Ambon banana leaves (*Musa paradisiaca* L.) on white rats (*Rattus norvegicus* L.) induced by carrageenan]. *Galenika Journal of Pharmacy*, 1(2), 126–132.
- Sundaram, S., Anjum, S., Dwivedi, P., & Rai, G. K. (2011). Antioxidant activity and protective effect of banana peel against oxidative hemolysis of human erythrocyte at different stages of ripening. *Applied Biochemistry and Biotechnology*, 164(7), 1192–1206.
<https://doi.org/10.1007/s12010-011-9205-3>
- Tayab, M. A., Islam, M. N., Ashfaq, K., & Chowdhury, A. (2022). Biomedicine & pharmacotherapy targeting neuroinflammation by polyphenols: a promising therapeutic approach against inflammation-associated

- depression. *Biomedicine & Pharmacotherapy*, 147, 112668. <https://doi.org/10.1016/j.biopha.2022.112668>
- Toh, P. Y., Leong, F. S., Chang, S. K., Khoo, H. E., & Yim, H. S. (2016). Optimization of extraction parameters on the antioxidant properties of banana waste. *Acta Scientiarum Polonorum, Technologia Alimentaria*, 15(1), 65–78. <https://doi.org/10.17306/JAFS.2016.1.7>
- Torrente, M. P., Gelenberg, A. J., & Vrana, K. E. (2012). Boosting serotonin in the brain: is it time to revamp the treatment of depression? *Journal of Psychopharmacology*, 26(5), 629–635. <https://doi.org/10.1177/02698811111430744>
- Vadodaria, K. C., Stern, S., Marchetto, M. C., & Gage, F. H. (2018). Serotonin in psychiatry: in vitro disease modeling using patient-derived neurons. *Cell and Tissue Research*, 371(1), 161–170. <https://doi.org/10.1007/s00441-017-2670-4>
- Vilela, C., Santos, S. A. O., Villaverde, J. J., Oliveira, L., Nunes, A., Cordeiro, N., Freire, C. S. R., & Silvestre, A. J. D. (2014). Lipophilic phytochemicals from banana fruits of several *Musa* species. *Food Chemistry*, 162, 247–252. <https://doi.org/10.1016/j.foodchem.2014.04.050>
- Vu, H. T., Scarlett, C. J., & Vuong, Q. V. (2018). Phenolic compounds within banana peel and their potential uses: A review. *Journal of Functional Foods*, 40(November 2017), 238–248. <https://doi.org/10.1016/j.jff.2017.11.006>
- Vu, H. T., Scarlett, C. J., & Vuong, Q. V. (2019). Changes of phytochemicals and antioxidant capacity of banana peel during the ripening process; with and without ethylene treatment. *Scientia Horticulturae*, 253(April), 255–262. <https://doi.org/10.1016/j.scienta.2019.04.043>
- Waalkes, T. P., Sjoerdsma, A., Creveling, C. R., Weissbach, H., & Udenfriend, S. (1958). Serotonin, norepinephrine, and related compounds in bananas. *Science*, 127(3299), 648–650. <https://doi.org/10.1126/science.127.3299.648>
- Waløen, K., Kleppe, R., Martinez, A., & Haavik, J. (2017). Tyrosine and tryptophan hydroxylases as therapeutic targets in human disease. *Expert Opinion on Therapeutic Targets*, 21(2), 167–180. <https://doi.org/10.1080/14728222.2017.1272581>
- Widoyanti, A. A. E., Chaikong, K., Rangsinth, P., Saengratwatchara, P., Leung, G. P. H., & Prasansuklab, A. (2023). Valorization of nam wah banana (*Musa paradisiaca* L.) byproducts as a source of bioactive compounds with antioxidant and anti-inflammatory properties: *in vitro* and *in silico* studies. *Foods*, 12(21). <https://doi.org/10.3390/foods12213955>
- Winiarska-mieczan, A., Kwiecie, M., & Donaldson, J. (2023). Anti-inflammatory, antioxidant, and neuroprotective effects of polyphenols — polyphenols as an element of diet therapy in depressive disorders. *International Journal of Molecular Sciences*, 24, 2258.
- Xiao, R., Beck, O., & Hjerdahl, P. (1998). On the accurate measurement of serotonin in whole blood. *Scandinavian Journal of Clinical and Laboratory Investigation*, 58(6), 505–510. <https://doi.org/10.1080/00365519850186319>
- Yakubu, M. T., Oyeyipo, T. O., Quadri, A. L., & Akanji, M. A. (2013). Effects of aqueous extract of *Musa paradisiaca* root on testicular function parameters of male rats. *Journal of Basic and Clinical Physiology and Pharmacology*, 24(2), 151–157. <https://doi.org/10.1515/jbcpp-2012-0059>
- Zou, F., Tan, C., Zhang, B., Wu, W., & Shang, N. (2022). The valorization of banana by-products: nutritional composition, bioactivities, applications, and future development. *Foods*, 11(20). <https://doi.org/10.3390/foods11203170>

Exploring the Mood-Boosting Potential of Bananas: A Comprehensive Review

ORIGINALITY REPORT

6%

SIMILARITY INDEX

3%

INTERNET SOURCES

7%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to Academic Library Consortium

Student Paper

1%

2

www.coursehero.com

Internet Source

1%

3

Gokare A. Ravishankar, Akula Ramakrishna. "Serotonin and Melatonin - Their Functional Role in Plants, Food, Phytomedicine, and Human Health", CRC Press, 2019

Publication

1%

4

Submitted to Konsorsium Perguruan Tinggi Swasta Indonesia

Student Paper

1%

5

"Handbook of Banana Production, Postharvest Science, Processing Technology, and Nutrition", Wiley, 2020

Publication

1%

6

T. K. Lim. "Edible Medicinal And Non Medicinal Plants", Springer Science and Business Media LLC, 2012

Publication

1%

7

"Harsh Environment and Plant Resilience", Springer Science and Business Media LLC, 2021

Publication

1%

8

novaprd-lb.newcastle.edu.au

Internet Source

1%

Exclude quotes	On	Exclude matches	< 1%
Exclude bibliography	On		