

In Vitro Antiviral Activity of Dried Red Jujube Fruit (*Ziziphus jujuba*) Ethanol Extract against DENV-2

by Silvia Sutandhio

Submission date: 02-Aug-2025 09:13PM (UTC+0700)

Submission ID: 2724119694

File name: Pharmacognj-16-5-1047.pdf (297.19K)

Word count: 3290

Character count: 16623

In Vitro Antiviral Activity of Dried Red Jujube Fruit (*Ziziphus jujuba*) Ethanol Extract against DENV-2

Niluh Suwasanti¹, Cecilia Putri Tedyanto², Silvia Sutandhio³, Teguh Hari Sucipto⁴

Niluh Suwasanti¹, Cecilia Putri Tedyanto², Silvia Sutandhio³, Teguh Hari Sucipto⁴

¹Department of Clinical Pathology, Faculty of Medicine, Universitas Katolik Widya Mandala Surabaya, Surabaya, INDONESIA
²Research Assistant, Faculty of Medicine, Universitas Katolik Widya Mandala Surabaya, Surabaya, INDONESIA

³Department of Microbiology and Parasitology, Faculty of Medicine, Universitas Katolik Widya Mandala Surabaya, Surabaya, INDONESIA

⁴Dengue Study Group, Institute of Tropical Disease, Universitas Airlangga, Surabaya, INDONESIA

Correspondence

Cecilia Putri Tedyanto

Faculty of Medicine, Universitas Katolik Widya Mandala Surabaya, Jalan Kalsaria Selatan 1, Surabaya, INDONESIA.
 E-mail: cecili@ukwms.ac.id

History

- Submission Date: 14-08-2024;
- Review completed: 22-09-2024;
- Accepted Date: 25-09-2024.

DOI : 10.5530/pj.2024.16.169

Article Available online

<http://www.phcogj.com/v16/i5>

Copyright

© 2024 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.



ABSTRACT

Dengue infection has become one of the most important mosquito-borne diseases worldwide and is caused by the dengue virus (DENV). Recently, neither patent drug, phytopharmaceutical medicine, nor standardized herbal medicine has been officially available against DENV. Dried red jujube fruit (*Ziziphus jujuba*) ethanol extract has been proven to have an antiviral effect, anti-inflammatory efficacy, and antioxidant properties, which have potential activity against DENV infection. This research was conducted to analyze the antiviral activity of dried red jujube fruit ethanol extract against DENV-2 in vitro. The half-maximal cytotoxic concentration (CC₅₀) and half-maximal inhibitory concentration (IC₅₀) were examined on Vero cells by a 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, whereas the half-maximal effective concentration (EC₅₀) was determined using luminescence assay. The selectivity index (SI) value was determined from the ratio of CC₅₀ and EC₅₀. Dried red jujube fruit ethanol extracts inhibited DENV-2 in 24.59%, 22.39%, 17.98%, 14.5%, 6.42%, and 1.28% at 80 µg/mL, 40 µg/mL, 20 µg/mL, 10 µg/mL, 5 µg/mL, and 2.5 µg/mL, respectively. The extract exhibited antiviral activity against DENV-2, showing a CC₅₀ of 67.73 µg/mL, an IC₅₀ value of 166.18 µg/mL, and an EC₅₀ of 64.87 µg/mL, with an SI of 1.04. The LD₅₀ value was 707.95 mg/kg. Dried red jujube fruit ethanol extract could be a potential candidate for developing an antiviral against DENV-2.

Keywords: *Ziziphus jujuba*, dried red jujube fruit, dengue virus, DENV-2, antiviral.

INTRODUCTION

Dengue is a primary arboviral global concern infection in tropical and subtropical regions, transmitted by *Aedes albopictus* and *Aedes aegypti* as their vectors. Humans can be infected through the bite of infected female mosquito. Dengue virus (DENV) consists of several serotypes, namely DENV-1, DENV-2, DENV-3, DENV-4, and more recently DENV-5.¹ Each DENV serotype can cause an acute febrile illness with several different clinical symptoms, ranging from dengue fever (DF) to dengue hemorrhagic fever (DHF) or dengue shock syndrome (DSS).²

Dengue infection has become one of the most important mosquito-borne diseases worldwide. The incidence of dengue infection, according to reported cases to the World Health Organization (WHO), increased from around 50,000 cases in 2000 to 5.2 million cases in 2019.³ The disease is endemic in more than 100 countries, with Asia presenting approximately 70% of the global disease burden. There were 390 million DENV infections per year, resulting in up to 36,000 deaths.⁴ There were at least 23 countries that reported dengue outbreaks in 2003, with approximately more than 5 million cases in 80 countries.⁵ This self-limiting disease usually only requires supportive therapy as primary therapy, but improper treatment due to inadequate sources of knowledge, the economy, or health access could worsen the disease.

Recently, neither patent drugs, phytopharmaceutical medicine, nor standardized herbal medicine have been officially available against DENV. Dried red jujube fruit (DRJF), or Chinese jujube (*Ziziphus jujuba*), has several

secondary active metabolites, such as flavonoids, phenolics, triterpenoids, and carotenoids.^{6,7} DRJF has proven to have an antiviral effect against the influenza virus, an anti-inflammatory effect, and a potential source of antioxidant properties.⁸⁻¹⁰ This research was conducted to establish the potential of DRJF ethanol extract as an antiviral against DENV-2 in vitro.

RESULTS AND DISCUSSION

Jujube, also known as Chinese jujube, is a fruit rich in nutrients and bioactive compounds. The fruit contains carbohydrates, proteins, fats, vitamins (A, B1, B2, B3, B6, and C), minerals (potassium, phosphorus, calcium, magnesium, sodium, iron, manganese, copper, and zinc), fiber, and bioactive compounds (phenolics, flavonoids, triterpenoids, and carotenoids).¹¹⁻¹³ Phenolics and flavonoids play the main role as antioxidants in jujube fruit.¹⁴⁻¹⁶

The maturity level of the fruit is generally divided into early, middle, and late stages. Its color can be evaluated from green at early maturity to red at late maturity. Research on fresh jujube fruits revealed that the total phenolic content (TPC) and total flavonoid content (TFC) were at their highest in early maturity and continued to decrease to the lowest in late maturity.¹⁴ The higher the level of maturity, the lower the levels of bioactive compounds.

Jujube is a fruit that is commonly found and consumed in red and dried form. The drying process affects the structure, texture, color, chemical components, and aroma of a product. Dried products also have lower nutritional value than fresh ones because several components are lost during drying. The processes that occur include degradation of proteins, Maillard

Cite this article: Suwasanti N, Tedyanto CP, Sutandhio S, Sucipto TH. In Vitro Antiviral Activity of Dried Red Jujube Fruit (*Ziziphus jujuba*) Ethanol Extract against DENV-2. *Pharmacogn J.* 2024;16(5): 1047-1050.

reactions, caramelization, lipid oxidation, and degradation of water-soluble and fat-soluble vitamins. Changes in these components are influenced by temperature, method, and drying duration.¹⁷ Previous studies revealed that red jujube fruit from fresh to dried experiences decreased levels of vitamins, TPC, TFC, and antioxidant activity.^{18,19} Despite the significant component changes, DRJF has been shown to have several health benefits.^{1,20}

The highest cell viability rate was found in DRJF ethanol extract at 2.5 µg/mL, with an amount of 126.16%. The cell viability rates of DRJF ethanol extract at 5 µg/mL, 10 µg/mL, 20 µg/mL, and 40 µg/mL were 104.3%, 78.81%, 72.52%, and 59.93%, respectively. The lowest cell viability rate was found in DRJF ethanol extract at 80 µg/mL, with an amount of 50.66%. Pearson correlation analysis showed a high negative correlation between the concentration of DRJF ethanol extract and the percentage of cell viability ($r = -0.809$, $p = 0.051$). The percentage of Vero cell viability decreased as the concentration of DRJF ethanol extract increased, forming a linear line, as seen in Figure 1.

The highest viral inhibition rate was found in DRJF ethanol extract at 80 µg/mL, with an amount of 24.59%. The viral inhibition rates of DRJF ethanol extract at 40 µg/mL, 20 µg/mL, 10 µg/mL, and 5 µg/mL were 22.39%, 17.98%, 14.5%, and 6.42%, respectively. The lowest viral inhibition rate was found in DRJF ethanol extract at 2.5 µg/mL, with an amount of 1.28%. Pearson correlation analysis showed a high positive correlation between the concentration of DRJF ethanol extract and the percentage of viral inhibition ($r = 0.824$, $p = 0.044$). The percentage of DENV-2 inhibition increased as the concentration of DRJF ethanol extract increased, forming a linear line, as seen in Figure 2.

The *in vitro* experimental results in Table 3 showed that DRJF ethanol extract had a CC_{50} value of 67.73 µg/mL, an IC_{50} value of 166.18 µg/mL, and an EC_{50} value of 64.87 µg/mL. The SI value from the ratio of CC_{50} and EC_{50} was 1.04. SI higher than 1 indicates that the extract efficacy against the virus is greater than the toxicity to normal cells, even though it was not a strong activity. The estimated LD_{50} value of DRJF ethanol extract was 707.95 mg/kg.

MATERIAL AND METHODS

Tools and Materials Preparation

Tools used in experimental study include biological safety cabinet (AIRTECH, Airtech Equipment Pte Ltd, Singapore), CO₂ incubator (Celculture, Esco Micro Pte. Ltd., Singapore), micropipette (NichipetEXII Multi, Nichiryo Co., Ltd., Japan), 96-well plate (NEST,




Figure 1. Percentage of Vero cell viability treated with various concentrations of dried red jujube fruit ethanol extract.




Figure 2. Percentage of DENV-2 inhibition by various concentrations of dried red jujube fruit ethanol extract.

Wuxi NEST Biotechnology Co., Ltd., China), T75 culture flask, refrigerated centrifuge, improved Neubauer counting chamber (Assistant, Hecht Glaswarenfabrik GmbH & Co. KG, Germany), light microscope (Nikon, Nikon Corp., Japan), hand tally counter, digital calculator, absorbance microplate reader (Bio-Rad, Bio-Rad Laboratories, Inc., California), luminescence microplate reader (Glomax, Promega Corp., USA), microcentrifuge tube, rotary evaporator, and oven.

Materials used in the experimental study include Vero cell, heat-inactivated DENV-3, active DENV-2, DRJF, ethanol 70%, 0.1% DMSO, and CMC-Na 1%.

Extract Preparation

DRJF were purchased from a local herbal shop and identified by the Technical Implementation Unit of the Herbal Laboratory, Materia Medica Batu, East Java, Indonesia. The DRJF ethanol extract was prepared at the Faculty of Pharmacy, Universitas Katolik Widya Mandala Surabaya, Surabaya, Indonesia, as described in the previous study.¹¹

Virus Preparation

DENV isolates were obtained from the Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia. The serotypes of DENV used in the experiment were active DENV-2 and heat-inactivated DENV-3. The virus titer was calculated using a hemocytometer under a light microscope and expressed in Foci-Forming-Unit (FFU). DENV-3 was inactivated using the heat treatment method at 56°C for 30 minutes.

Determination of Half-maximal Cytotoxic Concentration (CC_{50})

The half-maximal cytotoxic concentration (CC_{50}) study was performed to ensure that the DRJF ethanol extract was not toxic to the cell. The CC_{50} value was determined using a 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay based on cell viability after being treated with the extract. 2×10^4 cells per well of Vero cells were added to a 96-well plate and incubated at 37°C with 5% CO₂. After 24 hours, Vero cells were treated with 100 µL of diluted 0.1% DMSO with six various concentrations of DRJF ethanol extract (2.5 µg/mL, 5 µg/mL, 10 µg/mL, 20 µg/mL, 40 µg/mL, and 80 µg/mL), then incubated at 37°C with CO₂ 5%. After 48 hours, 20 µL of MTT solution was added to each well and incubated for four hours. The absorbances were measured using a microplate reader at 595 nm. The percentage of cell viability was calculated as $[(A / B) \times 100]$, where A and B are the absorbance of the extract test on the cell and the absorbance of the solvent test on the

Table 1. The CC_{50} , IC_{50} , EC_{50} , Selectivity Index, and LD_{50} values of dried red jujube fruit ethanol extract towards DENV-2 in Vero Cell.

| Concentration (μ g/mL) | Cell Viability (%) | CC_{50} (μ g/mL) | Viral Inhibition (%) | IC_{50} (μ g/mL) | EC_{50} (μ g/mL) | Selectivity Index ^a | LD_{50} ^b (mg/kg) |
|-----------------------------|--------------------|-------------------------|----------------------|-------------------------|-------------------------|--------------------------------|--------------------------------|
| 80 | 50.66 | 67.73 | 24.59 | 166.18 | 64.87 | 1.04 | 707.95 |
| 40 | 59.93 | | 22.39 | | | | |
| 20 | 72.52 | | 17.98 | | | | |
| 10 | 78.81 | | 14.5 | | | | |
| 5 | 104.3 | | 6.42 | | | | |
| 2.5 | 126.16 | | 1.28 | | | | |

DRJF Ethanol Extract: Dried Red Jujube Fruit Ethanol Extract; CC_{50} : Half-maximal Cytotoxic Concentration; IC_{50} : Half-maximal Inhibitory Concentration; EC_{50} : Half-maximal Effective Concentration; LD_{50} : Median Lethal Dose

^a Selectivity Index = CC_{50}/EC_{50}

^b $\log LD_{50} = 0.372 \log IC_{50} + 2.024$

cell, respectively. The CC_{50} value was estimated from a linear regression analysis between the percentage of cell viability and the concentration of DRJF ethanol extract.

Determination of Half-maximal Inhibitory Concentration (IC_{50})

The half-maximal inhibitory concentration (IC_{50}) study was performed as a screening to ensure that the DRJF ethanol extract has antiviral activity against DENV-2. The IC_{50} value was determined using an MTT assay based on viral titer after the cell was infected with DENV-2 and treated with the extract. All experiments were performed in triplicate. Vero cells were added at a density of 2×10^4 cells/well in a 96 well plate and incubated at 37°C with 5% CO_2 . After 24 hours, Vero cells were infected with DENV-2 at a multiplicity of infection (MOI) of 1 FFU/cell. After two hours, 100 μL of 0.1% DMSO with six various concentrations of DRJF ethanol extract were added. The plates were further incubated at 37°C with 5% CO_2 for two days, and 20 μL of MTT solution was added to each well and incubated for four hours. The absorbances were measured using a microplate reader at 595 nm. The percentage of viral inhibition was calculated as $[(A - B) / (C - B) \times 100\%]$, where A, B, and C are the absorbance of the extract on the virus-infected cell, the absorbance of the virus control, and the absorbance of the cell control, respectively. The IC_{50} value was estimated from a linear regression analysis between the percentage of viral inhibition and the concentration of DRJF ethanol extract.

Determination of Half-maximal Effective Concentration (EC_{50})

The half-maximal effective concentration (EC_{50}) study was performed to explore the effective concentration of DRJF ethanol extract against DENV-2. The EC_{50} value was determined by a luminescence assay using a microplate reader. The treatment was the same as previously described in the IC_{50} study.

Determination of Median Lethal Dose (LD_{50})

The median lethal dose (LD_{50}) value was obtained from the IC_{50} value ($\mu\text{g/mL}$) based on a regression formula approved by the Interagency Coordinating Committee on the Validation of Alternative Methods (ICCVAM).¹² The regression formula was calculated as $[\log LD_{50} = 0.372 \log IC_{50} + 2.024]$. The LD_{50} value was expressed in mg/kg.

Selectivity Index Analysis

The CC_{50} and EC_{50} values are the average of triplicate assays with six concentrations within the inhibitory range of the compounds. The selectivity index (SI) was determined from the ratio of CC_{50}/EC_{50} .

Statistical Analysis

A Pearson correlation analysis was conducted using IBM SPSS software version 25 (IBM Corp., New York).

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

REFERENCES

- Mustafa MS, Rasotgi V, Jain S, Gupta V. Discovery of fifth serotype of dengue virus (DENV-5): A new public health dilemma in dengue control. Med J Armed Forces India. 2015 Jan;71(1):67-70.
- Tayal A, Kabra SK, Lodha R. Management of dengue: An updated review. Indian J Pediatr. 2023 Feb;90(2):168-177.
- World Health Organization (WHO). Disease outbreak news: Dengue - Global situation. 2023.
- Yuan K, Chen Y, Zhong M, Lin Y, Liu L. Risk and predictive factors for severe dengue infection: A systematic review and meta-analysis. PLoS One. 2022 Apr 15;17(4):e0267186.
- Centers for Disease Control and Prevention (CDC). Dengue on the rise: Get the facts. 2024.
- Shi Q, Han G, Liu Y, Jiang J, Jia Y, Li X. Nutrient composition and quality traits of dried jujube fruits in seven producing areas based on metabolomics analysis. Food Chem. 2022 Aug 15;385:132627.
- Wu Z, Zhang S, Liu L, Wang L, Ban Z. The grade of dried jujube (Ziziphus jujuba Mill. cv. Junzao) affects its quality attributes, antioxidant activity, and volatile aroma components. Foods. 2023 Feb 26;12(5):989.
- Hong EH, Song JH, Kang KB, Sung SH, Ko HJ, Yang H. Anti-influenza activity of betulinic acid from Zizyphus jujuba on influenza A/PR/8 virus. Biomol Ther (Seoul). 2015 Jul;23(4):345-9.
- Mesaik AM, Poh HW, Bin OY, Elawad I, Alsayed B. In vivo anti-inflammatory, anti-bacterial and anti-diarrhoeal activity of Ziziphus jujuba fruit extract. Open Access Maced J Med Sci. 2018 May 15;6(5):757-766.
- Yang C, Zhu X, Liu W, Huang J, Xie Z, Yang F, Zhang L, Wei Y. Dietary dried jujube fruit powder (DJFP) supplementation improves growth performance, antioxidant stability, and meat composition in broilers. Foods. 2023 Mar 29;12(7):1463.
- Tedyanto CP, Wihanto L, Hendrata AP. Hepatoprotective effect of dried red jujube fruit extract against acetaminophen-induced acute hepatotoxicity. Cureus. 2023 Jan 2;15(1):e33272.
- ICCVAM. Peer review panel report: the use of in vitro basal cytotoxicity test methods for estimating starting doses for acute oral systemic toxicity testing. NIH publication no. 07-4519, Research Triangle Park, National Toxicology Program, 2006.
- Rashwan AK, Karim N, Shishir MRI, Bao T, Lu Y, Chen W. Jujube fruit: A potential nutritious fruit for the development of functional food products. J Funct Foods. 2020;75:104205.
- Wang BN, Huang Q, Venkatasamy C, Chai H, Gao H, Cheng N, et al. Changes in phenolic compounds and their antioxidant capacities in jujube (Ziziphus jujuba Miller) during three edible maturity stages. LWT Food Sci Technol. 2016;66:56-62.

15. Huang W, Wang Y, Jiang X, Sun Y, Zhao Z, Li S. Protective effect of flavonoids from *Ziziphus jujuba* cv. Jinsixiaozao against acetaminophen-induced liver injury by inhibiting oxidative stress and inflammation in mice. *Molecules*. 2017 Oct 20;22(10):1781.
16. Reche J, Almansa MS, Hernández F, Amorós A, Legua P. Physicochemical and antioxidant capacity of jujube (*Ziziphus jujuba* Mill.) at different maturation stages. *Agronomy*. 2021;11(1):132.
17. Guiné RPF. The drying of foods and its effect on the physical-chemical, sensorial, and nutritional properties. *Int J Food Eng*. 2018;4(2).
18. Tepe F, Ekinçi R. Drying characteristics and some quality parameters of whole jujube (*Zizyphus jujuba* Mill.) during hot air drying. *Ital J Food Sci*. 2021;33:1-15.
19. Wang C, Cao J, Jiang W. Effect of the drying method on browning of flesh, antioxidant compounds and antioxidant capacity of Chinese jujube (*Zizyphus jujuba* Mill.) fruit. *Curr Top Nutraceutical Res*. 2016;14:161-9.
20. Zhu D, Jiang N, Wang N, Zhao Y, Liu X. A literature review of the pharmacological effects of jujube. *Foods*. 2024 Jan 6;13(2):193.

Cite this article: Suwasanti N, Tedyanto CP, Sutandhio S, Sucipto TH. In Vitro Antiviral Activity of Dried Red Jujube Fruit (*Ziziphus jujuba*) Ethanol Extract against DENV-2. *Pharmacogn J*. 2024;16(5): 1047-1050.

In Vitro Antiviral Activity of Dried Red Jujube Fruit (Ziziphus jujuba) Ethanol Extract against DENV-2

ORIGINALITY REPORT

13%

SIMILARITY INDEX

3%

INTERNET SOURCES

11%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

- 1

Yun Jung Kim, Seong-Beom Lee, Hyang Burm Lee. "Oleic acid enhances keratinocytes differentiation via the upregulation of miR-203 in human epidermal keratinocytes", Journal of Cosmetic Dermatology, 2018
Publication

2%
- 2

Aidin Mohammadi Zonouz, Sahar Taghavi, Siroos Nekooei, Khalil Abnous et al. "Synthesis of targeted doxorubicin-loaded gold nanorod -mesoporous manganese dioxide core-shell nanostructure for ferroptosis, chemo-photothermal therapy in vitro and in vivo", International Journal of Pharmaceutics, 2024
Publication

1%
- 3

Submitted to Chonnam National University
Student Paper

1%
- 4

Ton Duc Thang University
Publication

1%
- 5

Knut Husmann, P.M Sexton, J.A Fischer, W Born. "Mouse receptor-activity-modifying proteins 1, -2 and -3: amino acid sequence, expression and function", Molecular and Cellular Endocrinology, 2000
Publication

1%
- 6

cuir.car.chula.ac.th
Internet Source

1%
- 7

Oliver, Jonathan, Paula Tennant, and Marc Fuchs. "Virus-Resistant Transgenic Horticultural Crops : Safety Issues and Lessons from Risk Assessment Studies",

1%

Transgenic Horticultural Crops Challenges and Opportunities, 2011.

Publication

| | | |
|---|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 8 | Yuxiang Ji, Gaozhe Cai, Cheng Liang, Zehang Gao, Weimin Lin, Zizhen Ming, Shilun Feng, Hongwei Zhao. "A microfluidic immunosensor based on magnetic separation for rapid detection of okadaic acid in marine shellfish", <i>Analytica Chimica Acta</i> , 2023 | 1 % |
|---|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|

Publication

| | | |
|---|--------------------------------------|-----|
| 9 | Submitted to universititeknologimara | 1 % |
|---|--------------------------------------|-----|

Student Paper

| | | |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 10 | Soha Gomaa, Mohamed Nassef, Randa El-Naggar, Ahmed Massoud, Mona El-Kholy. "Anti-tumoral Immunity and Chemo-preventive Effectiveness of Herbal Extracts of Curcumin, Ginger, Clove and Amygdaline in Ehrlich Ascites Carcinoma-Challenging Mice", <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2024 | 1 % |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|

Publication

| | | |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 11 | Bin Liu, Ting-Ting Gao, Xiao-Yu Fu, Zhen-Hao Xu, Hao Ren, Ping Zhao, Zhong-Tian Qi, Zhao-Ling Qin. "PTEN Lipid Phosphatase Activity Enhances Dengue Virus Production through Akt/FoxO1/Maf1 Signaling", <i>Virologica Sinica</i> , 2020 | 1 % |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|

Publication

| | | |
|----|-----------------------------------|-----|
| 12 | Submitted to Ngee Ann Polytechnic | 1 % |
|----|-----------------------------------|-----|

Student Paper

| | | |
|----|----------------------------------|-----|
| 13 | Submitted to University of Macau | 1 % |
|----|----------------------------------|-----|

Student Paper

| | | |
|----|------------------------------------|-----|
| 14 | Submitted to University of Reading | 1 % |
|----|------------------------------------|-----|

Student Paper

| | | |
|----|----------------------------------------------------------------------------------------------------------------------------|-----|
| 15 | Xiaoxue Xu, Sheng Guo, Jiangyan Chen, Yuhan Song, Xueli Wu, Feng Liu, Jin-Ao Duan. "Comparative pharmacokinetics of multi- | 1 % |
|----|----------------------------------------------------------------------------------------------------------------------------|-----|

components in normal and stomach cold syndrome rats after oral administration of Zingiberis Rhizoma – Jujubae Fructus herb pair and its single herb extracts by UHPLC-MS/MS", Journal of Chromatography B, 2024

Publication

16

Spyridon A Petropoulos, Ângela Fernandes, Lillian Barros, Isabel CFR Ferreira. " A comparison of the phenolic profile and antioxidant activity of different L. ecotypes ", Journal of the Science of Food and Agriculture, 2018

Publication

1%

17

Usman Sumo Friend Tambunan, Mochammad Arfin Fardiansyah Nasution, Fauziah Azhima, Arli Aditya Parikesit et al. "Modification of S-Adenosyl-L-Homocysteine as Inhibitor of Nonstructural Protein 5 Methyltransferase Dengue Virus Through Molecular Docking and Molecular Dynamics Simulation", Drug Target Insights, 2017

Publication

1%

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On