

Paulus Supit

1. CLINICAL PROFILE OF STROKE PATIENTS IN GOTONG ROYONG HOSPITAL

 Paulus Supit

Document Details

Submission ID

trn:oid:::3618:114882749

Submission Date

Sep 29, 2025, 5:07 PM GMT+7

Download Date

Sep 29, 2025, 5:10 PM GMT+7

File Name

1-Clinical_profile_of_stroke_patients_-1.pdf

File Size

206.6 KB

13 Pages

4,749 Words

24,293 Characters

1% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- Bibliography
- Small Matches (less than 11 words)

Exclusions

- 5 Excluded Sources
- 8 Excluded Matches

Match Groups

-  **2 Not Cited or Quoted 1%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 1%  Internet sources
- 0%  Publications
- 0%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 2** Not Cited or Quoted 1%
 Matches with neither in-text citation nor quotation marks
- 0** Missing Quotations 0%
 Matches that are still very similar to source material
- 0** Missing Citation 0%
 Matches that have quotation marks, but no in-text citation
- 0** Cited and Quoted 0%
 Matches with in-text citation present, but no quotation marks

Top Sources

- 1% Internet sources
- 0% Publications
- 0% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

- 1 Internet

mmrjournal.biomedcentral.com <1%
- 2 Internet

tsitn.org <1%

CLINICAL PROFILE OF STROKE PATIENTS IN GOTONG ROYONG HOSPITAL

Paulus Supit^{1,2*}, Nita Kurniawati^{1,2}, Desy Kartikasari^{1,2}, Evander Aloysius Raymond Desun²,
Ferdinand Erwin^{2,3}

*Correspondent author email: paulus.supit@ukwms.ac.id

DOI: <https://doi.org/10.33508/jwmj.v6i3.5781>

ABSTRACT

Introduction: Stroke is a global health issue due to its high prevalence, mortality, and associated disabilities. The Global Burden of Disease Study identifies ischemic and hemorrhagic strokes as significant contributors to global health burdens. Understanding these factors in different populations is crucial for developing targeted interventions and improving patient outcomes. **Method:** This retrospective descriptive study analyzed medical records of stroke patients at Gotong Royong Hospital from January to December 2023. Patients aged 18 years or older with acute stroke confirmed by CT or MRI were included. **Result:** Out of 224 stroke patients, 58% were male and 42% female. Most patients were over 70 years old (31.7%), with 29% aged 51-60 and 26.3% aged 61-70. Ischemic stroke was predominant (91.5%), while 8.5% had hemorrhagic stroke. Many patients (40.2%) arrived at the emergency department more than 24 hours after symptom onset. Hypertension was present in 68.8% of patients, diabetes in 36.6%, and dyslipidemia in 24.1%. The average length of stay was 4.13 days. Blood pressure measurements showed high prevalence of Grade 2 hypertension in systolic (37.05%) and prehypertension in diastolic (30.80%). Laboratory results revealed mean random blood glucose of 174.3 mg/dL, fasting blood glucose of 110.42 mg/dL, HbA1C of 8.06%, LDL cholesterol of 120.31 mg/dL, and triglycerides of 151.26 mg/dL. **Conclusion:** The majority of acute stroke patients at Gotong Royong Hospital are male, over 60, with ischemic stroke, and present more than 24 hours post-onset. These patients typically have a history of hypertension but no previous strokes, with an average hospital stay of 4 days. Many exhibit Grade 2 hypertension in systolic blood pressure and prehypertension in diastolic pressure. Understanding these profiles can guide better management and preventive strategies for stroke patients.

Keywords: profile, risk factor, elderly, hemorrhagic stroke, ischemic stroke

¹ Department of Neurology, Faculty of Medicine, Widya Mandala Catholic University Surabaya

² Gotong Royong Hospital Surabaya

³ Faculty of Medicine, Widya Mandala Catholic University Surabaya

INTRODUCTION

Stroke is a global health problem due to its high prevalence, mortality, and the disabilities it causes. According to the Global Burden of Disease Study in 2013, ischemic and hemorrhagic strokes collectively contribute significantly to the global health burden.¹ The prevalence of stroke varies depending on factors such as stroke type, study design, and geographic location. Several studies have shown that stroke prevalence ranges from 7% to 67.3%, with hypertension, dyslipidemia, lack of physical activity, and coronary artery disease identified as significant risk factors.²⁻⁴ Furthermore, the risk of recurrent stroke is approximately 4% within the first month and 12% within the first year after the initial stroke, often associated with atherosclerotic and cardiovascular risk factors.⁵

Various studies have highlighted the importance of understanding stroke prevalence in different populations. For instance, research in China has demonstrated geographic variations in stroke prevalence, influenced by urban-rural differences over time.⁶ Some studies have shown that stroke prevalence is affected by sociodemographic factors, lifestyle, and comorbid conditions such as hypertension, diabetes mellitus, and cardiovascular diseases.⁷⁻⁹ The impact of stroke after an acute attack leaves

individuals with a history of stroke at an increased risk of developing dementia and cognitive impairment.¹⁰

In addition, stroke significantly contributes to mortality rates. In Indonesia, stroke is the leading cause of death, with a mortality rate of 131.8 per 100,000 population. Globally, stroke is the second leading cause of death after coronary heart disease.¹¹ Understanding the risk factors associated with stroke, such as hypertension, diabetes, and dyslipidemia, is crucial for preventive measures and improving health outcomes for at-risk individuals.^{7,9,12} This comprehensive approach can inform tailored prevention strategies, enhance early detection, and optimize treatment protocols, ultimately contributing to better management of stroke patients and reducing the overall burden of this debilitating condition. Thus, the present study aims to provide a detailed analysis of these factors among stroke patients at Gotong Royong Hospital, offering valuable data to guide clinical practices.

METHOD

This study is a retrospective descriptive study based on the medical records of stroke patients at Gotong Royong Hospital from January to December 2023. The study included patients aged 18 years or older with acute

stroke, verified by head computerized tomography (CT) or magnetic resonance imaging (MRI). Patients who did not meet the inclusion criteria were excluded from the study. Total sampling was used in this research by including data from all patients who met the inclusion criteria.

1 The study aimed to observe patients' characteristics, unmodifiable risk factors, and modifiable risk factors. Unmodifiable risk factors included age and sex, while modifiable risk factors encompassed a history of hypertension, diabetes, previous stroke, and dyslipidemia. Investigations performed included initial blood pressure at the emergency room, fasting and random blood glucose levels, HbA1C, and a fasting lipid profile (triglycerides and LDL). Other data collected included length of stay, onset-to-presentation time, and initial blood pressure at the emergency room. The data collected provided a comprehensive profile of stroke patients treated at Gotong Royong Hospital during the specified period.

RESULT

Gender

Out of 224 stroke patients, the majority are male, with 130 patients (58%), while 94 patients (42%) are female. This indicates that stroke incidence is higher in males than in females at Gotong Royong Hospital.

Age

The age distribution of stroke patients shows that the age group above 70 years is the most prevalent, with 71 patients (31.7%). The age group 51-60 years is also significant with 65 patients (29%), followed by the age group 61-70 years with 59 patients (26.3%). The age group 41-50 years has 23 patients (10.3%), and the age group under 40 years is the least, with only 6 patients (2.7%).

Type of Stroke

The majority of patients suffer from ischemic stroke, with 205 patients (91.5%), while 19 patients (8.5%) suffer from hemorrhagic stroke. This shows that ischemic stroke is more dominant among stroke patients at this hospital.

Stroke Onset

Based on stroke onset, most patients arrived at the hospital more than 24 hours after the onset of symptoms, with 89 patients (39.7%). Patients who arrived within less than 4.5 hours numbered 49 (21.9%), followed by those who arrived between 4.5 to 12 hours with 48 patients (21.4%), and patients who arrived within 12 to 24 hours numbered 38 (17%).

Medical History

Common medical histories among stroke patients include hypertension, found

in 154 patients (68.8%). Diabetes mellitus is also quite common with 82 patients (36.6%), followed by dyslipidemia in 54 patients (24.1%). Most patients do not have a history of previous strokes, with 136

patients (60.7%). A total of 71 patients (31.7%) have had one previous stroke, and 17 patients (7.6%) have a history of more than one stroke.

Tabel 1. Clinical Profile of Stroke Patients in Gotong Royong Hospital

Variable	N (%)	Mean ± SD	Median (Min-Max)		
Gender					
Male	130 (58%)	63.98 ± 12.54	64 (29-91)		
Female	94 (42%)				
Age					
< 40 years	6 (2.7%)				
41-50 years	23 (10.3%)				
51-60 years	65 (29%)				
61-70 years	59 (26.3%)				
> 70 years	71 (31.7%)				
Type of Stroke					
Ischemic	205 (91.5%)	-			
Hemorrhagic	19 (8.5%)				
Stroke Onset					
< 4.5 hours	49 (21.8%)	-			
4.5 – 12 hours	47 (21%)				
12 – 24 hours	38 (17%)				
> 24 hours	90 (40.2%)				
Medical History					
Hypertension	154 (68.8%)	-			
Diabetes Mellitus	82 (36.6%)				
Dyslipidemia	54 (24.1%)				
Previous Stroke History					
No history of stroke	136 (60.7%)	-			
1 time	71 (31.7%)				
>1 time	17 (7.6%)				
Length of treatment (days)	-	4.13 ± 2.64	4 (1-17)		

Length of Stay

The average length of stay for stroke patients at Gotong Royong Hospital is 4.13 days with a standard deviation of $\pm$ 2.64 days. The median length of stay is 4 days, with a range of stay between 1 to 17 days. This indicates that most patients are treated for about 4 days, although there is

considerable variability in the duration of stay depending on each patient's condition and needs.

Blood Pressure

The initial blood pressure profile of stroke patients at Gotong Royong Hospital

reveals significant findings (Table 2). The mean systolic blood pressure was 150.64 ± 30.22 mmHg, with a median of 150 mmHg (range 90-260 mmHg). Notably, 8.48% of patients had normal systolic blood pressure, while 22.32% were classified as prehypertensive. The majority of patients were hypertensive, with 32.14% falling into Hypertension Grade 1 and 37.05% into

Hypertension Grade 2. Similarly, the mean diastolic blood pressure was 88.46 ± 14.31 mmHg, with a median of 90 mmHg (range 60-120 mmHg). Among the patients, 14.29% had normal diastolic blood pressure, and 30.80% were prehypertensive. Hypertension Grade 1 and Grade 2 were prevalent, observed in 26.79% and 28.13% of patients, respectively.

Table 2. Initial Blood Pressure Profile of Stroke Patients in Gotong Royong Hospital

Initial Blood Pressure	N (%)	Mean ± SD	Median
Systolic Blood Pressure			
Normal (<120 mmHg)	19 (8.48%)	150.64 ± 30.22	150 (90-260)
Prehypertension (120-139 mmHg)	50 (22.32%)		
Hypertension Grade 1 (140-159 mmHg)	72 (32.14%)		
Hypertension Grade 2 (≥ 160 mmHg)	83 (37.05%)		
Diastolic Blood Pressure			
Normal (<80 mmHg)	32 (14.29%)	88.46 ± 14.31	90 (60-120)
Prehypertension (80-89 mmHg)	69 (30.80%)		
Hypertension Grade 1 (90-99 mmHg)	60 (26.79%)		
Hypertension Grade 2 (≥ 100 mmHg)	63 (28.13%)		

Laboratory Examination

The laboratory examination results in this study included blood glucose profiles (random blood glucose, fasting blood glucose, and HbA1C) and lipid profiles (LDL and triglycerides), as shown in Table 3. Random blood glucose levels were measured in 198 out of 224 patients, with a mean value of 174.3 mg/dL and a standard deviation of 102.96 mg/dL. Fasting blood glucose and HbA1C levels were assessed in

83 and 85 patients, respectively. The mean fasting blood glucose level was 110.42 mg/dL with a standard deviation of 35.46 mg/dL, and the mean HbA1C level was 8.06% with a standard deviation of 2.59%. Lipid profile examinations were conducted on 170 patients, revealing a mean LDL level of 120.31 ± 43.85 mg/dL, and a mean triglyceride level of 151.26 ± 96.10 mg/dL.

Tabel 3. Laboratory Profile of Stroke Patients in Gotong Royong Hospital

Laboratory Examination	Mean $\pm$ SD	Median (Min-Max)
Random Blood Sugar (n = 198)	174.3 $\pm$ 102.96	137 (32-704)
Fasting Blood Sugar (n = 83)	110.42 $\pm$ 35.46	101 (75-297)
HbA1C (n = 85)	8.06 $\pm$ 2.59	7.2 (3.9-15.1)
LDL Cholesterol (n = 170)	120.31 $\pm$ 43.85	120.5 (17-239)
Triglycerides (n = 170)	151.26 $\pm$ 96.10	125 (35-668)

DISCUSSION

This study reveals that the majority of stroke patients at Gotong Royong Hospital are male (58%), consistent with previous research indicating that men are more frequently affected by stroke.¹³⁻¹⁵ While men have a higher incidence of stroke, women tend to experience worse outcomes and higher mortality rates post-stroke. Women show higher 1 and 10-year mortality and recurrence rates and tend to have less favorable outcomes within the first year after a stroke.¹⁶ Additionally, women often present with greater disability and worse long-term outcomes, partly due to more severe initial neurological symptoms and higher susceptibility to neurological deterioration.¹⁷ Risk factors also differ by sex, with men are more likely to have modifiable risk factors such as smoking, alcohol consumption, and

dyslipidemia, which contribute significantly to stroke recurrence in males.^{18,19} Conversely, women are more likely to have hypertension, diabetes, and atrial fibrillation, which are associated with higher stroke severity and worse outcomes.^{19,20}

Our study also shows that most stroke patients are over 70 years old (31.7%), followed by those aged 51-60 years (29%) and 61-70 years (26.3%). This finding differs from previous studies, which indicated that the 61-70 age group had the highest incidence of stroke.^{14,21,22} Research indicates that most strokes occur in older adults, with 75% affecting individuals aged 65 and above, and those over 75 experiencing higher hospitalization and mortality rates.²³ Arterial stiffness and carotid intima-media thickness, indicators of vascular health, are more pronounced in middle-aged stroke patients compared to younger ones, further complicating the clinical picture as age advances.²⁴

The data indicates that ischemic stroke is overwhelmingly the most common type of stroke, affecting 205 patients (91.5%), while hemorrhagic stroke affects only 19 patients (8.5%). This finding is consistent with previous research conducted by Stevano et al. in Jakarta, which showed that ischemic stroke comprised the majority (89.4%) of stroke cases treated in hospitals.²⁵ The Global

Burden of Disease (GBD) 2017 study further highlighted that in 2017, there were 11.9 million incident strokes, with ischemic stroke constituting 65%, primary intracerebral hemorrhage 26%, and subarachnoid hemorrhage 9%.²⁶

In this study, most stroke patients (40.2%) presented to the emergency department more than 24 hours after stroke onset, followed by 21.8% who arrived within 4.5 hours of onset. Among ischemic stroke patients, 42.93% presented more than 24 hours after onset, with only 19.02% arriving within the golden period. Various studies have highlighted the proportion of patients arriving within this timeframe. For instance, a study at Yale-New Haven Hospital from 2010-2020 found that 37.5% of ischemic stroke patients presented early, within 4.5 hours.²⁷ The Get With The Guidelines–Stroke database analysis indicated that 28.3% of ischemic stroke patients arrived within 60 minutes, and 31.7% within 61 to 180 minutes, highlighting the importance of early arrival for better outcomes.²⁸ Additionally, a study on the impact of family knowledge on stroke management found that 58.2% of respondents with high knowledge levels brought patients to the hospital within the golden period.²⁹

The study also shows that more than half of the stroke patients had a history of hypertension (68.8%). Diabetes mellitus

was present in 36.6% of patients, and dyslipidemia in 24.1%. A study by Thrinetrapriya in 2021 showed that 70% of stroke patients had comorbid hypertension, consistent with our findings. That study also reported that 48% of patients had diabetes mellitus and 38% had dyslipidemia, while our study shows a higher percentage of patients with comorbid dyslipidemia compared to diabetes mellitus.¹⁴

The initial blood pressure examination of stroke patients in the emergency department revealed that the majority exhibited Grade 2 hypertension, followed by Grade 1 hypertension in systolic blood pressure. In terms of diastolic blood pressure, prehypertension was most common, followed by Grade 2 hypertension. Only 8.48% and 14.29% of patients had normal systolic and diastolic blood pressures, respectively. The mean systolic and diastolic blood pressures in this study were similar to those reported by Retnaningsih, with mean values of 150.97 ± 22.97 mmHg for systolic and 86.86 ± 13.30 mmHg for diastolic blood pressure.³⁰ Previous study by Dinata also found that only 8.33% of stroke patients presenting to the emergency department had normal blood pressure.³¹ Elevated blood pressure is common immediately following acute ischemic stroke (AIS), likely due to a physiological response to neurological insult and cerebral autoregulation failure.³²

The mean random blood glucose (174.3 ± 102.96 mg/dL) and fasting blood glucose (110.42 ± 35.46 mg/dL) levels did not show significant elevations among the patients. However, the mean HbA1C level was elevated ($8.06 \pm 2.59\%$). Research indicates that diabetes significantly increases the risk of stroke, with higher fasting glucose levels correlating with a greater incidence of stroke events.³³ HbA1C, reflecting long-term glycemic control, has been associated with stroke severity and infarct volume, with higher levels correlating with more severe strokes and larger infarct sizes.³⁴ Additionally, HbA1C levels are crucial in identifying undiagnosed diabetes or prediabetes in stroke patients.³⁵ Both fasting glucose and HbA1C levels are associated with intracranial atherosclerotic stenosis (ICAS), a common cause of stroke, indicating that elevated levels of these glycemic measures increase the risk of ICAS.³⁶

The mean LDL cholesterol level in this study was 120.31 ± 43.85 mg/dL, while triglycerides had a mean of 151.26 ± 96.10 mg/dL. LDL cholesterol, which transports cholesterol to cells, is known as a primary atherogenic lipoprotein when oxidized, significantly increasing the risk of atherosclerosis. Previous studies have highlighted the significance of serum LDL cholesterol levels in the context of ischemic stroke, with elevated LDL cholesterol being

associated with an increased risk of ischemic stroke³⁷⁻³⁹. Additionally, Kanagarajan et al. (2020) reported a direct association between increased triglyceride levels and ischemic stroke risk, emphasizing the role of triglycerides as a potential risk factor for stroke.³⁹ A report by Yaghi et al. found that total cholesterol and LDL levels have the strongest association with ischemic stroke occurrence.⁴⁰

CONCLUSION

This study shows that the majority of acute stroke patients at Gotong Royong Hospital are male, over the age of 60, present with ischemic stroke, and arrive at the emergency department more than 24 hours after the onset of symptoms. These patients commonly have a history of hypertension, but no previous history of stroke, with an average length of stay of 4 days. Many patients exhibit Grade 2 hypertension in systolic blood pressure and prehypertension in diastolic blood pressure.

REFERENCES

1. Feigin VL, Krishnamurthi R V., Parmar P, Norrving B, Mensah GA, Bennett DA, et al. Update on the Global Burden of Ischemic and Hemorrhagic Stroke in 1990-2013: The GBD 2013 Study. *Neuroepidemiology*. 2015;45(3):161-76. doi:

- 10.1159/000441085
2. Barbay M, Diouf M, Roussel M, Godefroy O. Systematic Review and Meta-Analysis of Prevalence in Post-Stroke Neurocognitive Disorders in Hospital-Based Studies. *Dement Geriatr Cogn Disord*. 2018;46(5–6):322–34. doi: 10.1159/000492920
3. Yi X, Luo H, Zhou J, Yu M, Chen X, Tan L, et al. Prevalence of stroke and stroke related risk factors: a population based cross sectional survey in southwestern China. *BMC Neurol*. 2020; doi: 10.21203/rs.2.17357/v3
4. Wu L, Wang A, Wang X, Zhao X, Wang C, Liu L, et al. Factors for short-term outcomes in patients with a minor stroke: results from China National Stroke Registry. *BMC Neurol*. 2015 Dec 9;15(1):253. doi: 10.1186/s12883-015-0505-z
5. Putri K, Tri Antika Rizki, Rahayu LP, Agustina EN. Stroke Recurrence Based on Stroke Prognosis Instrument II (SPI-II) and The Attack Number of Stroke. *KnE Life Sci*. 2019 Oct 9; doi: 10.18502/cls.v4i13.5352
6. Ru X, Wang W, Sun H, Sun D, Fu J, Ge S, et al. Geographical Difference, Rural-urban Transition and Trend in Stroke Prevalence in China: Findings from a National Epidemiological Survey of Stroke in China. *Sci Rep*. 2019 Nov 22;9(1):17330. doi: 10.1038/s41598-019-53848-1
7. Ahangar AA, Saadat P, Heidari B, Taheri ST, Alijanpour S. Sex difference in types and distribution of risk factors in ischemic and hemorrhagic stroke. *Int J Stroke*. 2018 Jan 31;13(1):83–6. doi: 10.1177/1747493017724626
8. Thammasarn K, Loahasiriwong W, Mahato RK, Sornlom K. Spatial Association Between Sociodemographic, Environmental Factors and Prevalence of Stroke Among Diabetes and Hypertension Patients in Thailand. *Open Public Health J*. 2022 Dec 2;15(1). doi: 10.2174/18749445-v15-e2210191
9. Li Q, Wu H, Yue W, Dai Q, Liang H, Bian H, et al. Prevalence of Stroke and Vascular Risk Factors in China: a Nationwide Community-based Study. *Sci Rep*. 2017 Jul 25;7(1):6402. doi: 10.1038/s41598-017-06691-1
10. De Ronchi D, Palmer K, Pioggiosi P, Atti AR, Berardi D, Ferrari B, et al. The Combined Effect of Age, Education, and Stroke on Dementia and Cognitive Impairment No Dementia in the Elderly. *Dement Geriatr Cogn Disord*. 2007;24(4):266–73. doi: 10.1159/000107102

11. World Health Organization. WHO methods and data sources for country-level causes of death 2000-2019. Geneva; 2020.
12. Hu L, Huang X, Zhou W, You C, Liang Q, Zhou D, et al. Associations between resting heart rate, hypertension, and stroke: A population-based cross-sectional study. *J Clin Hypertens*. 2019 May 5;21(5):589–97. doi: 10.1111/jch.13529
13. Balagopal K, Mathew SR, Koshy RA. Clinical profile of patients with acute ischaemic stroke undergoing thrombolysis in a tertiary hospital- A retrospective observational study. *IP Indian J Neurosci*. 2024 Apr 28;10(1):14–21. doi: 10.18231/j.ijn.2024.004
14. Thrinetrapriya NA, Jagadeesan M. Prevalence, Clinical Profile and Outcome of Patients Presenting with Stroke. *J Pharm Res Int*. 2021 Nov 9;146–57. doi: 10.9734/jpri/2021/v33i48B33273
15. Bhutta ZA, Akhtar N, Pathan SA, Castren M, Harris T, Ganesan GS, et al. Epidemiological profile of stroke in Qatar: Insights from a seven-year observational study. *J Clin Neurosci*. 2024 May;123:30–5. doi: 10.1016/j.jocn.2024.03.014
16. Guo X, Xiong Y, Huang X, Pan Z, Kang X, Chen C, et al. Sex-based differences in long-term outcomes after stroke: A meta-analysis. *Nasreldein A, editor. PLoS One*. 2023 Apr 27;18(4):e0283204. doi: 10.1371/journal.pone.0283204
17. Ryu WS, Chung J, Schellingerhout D, Jeong SW, Kim HR, Park JE, et al. Biological Mechanism of Sex Difference in Stroke Manifestation and Outcomes. *Neurology*. 2023 Jun 13;100(24). doi: 10.1212/WNL.0000000000207346
18. Dimitrov GT, Danovska MP, Marinova DL, Stoev PG, Simeonova YI. GENDER-RELATED DIFFERENCES IN MODIFIABLE RISK FACTORS AND ISCHEMIC STROKE SUBTYPE IN YOUNG AND MIDDLE-AGED PATIENTS – A PROSPECTIVE STUDY. *J IMAB - Annu Proceeding (Scientific Pap*. 2023 Jan 3;29(1):4793–9. doi: 10.5272/jimab.2023291.4793
19. Chung JY, Lee BN, Kim YS, Shin BS, Kang HG. Sex differences and risk factors in recurrent ischemic stroke. *Front Neurol*. 2023 Jan 26;14. doi: 10.3389/fneur.2023.1028431
20. Lappe C, Reinecke H, Feld J, Köppe J. Sex Differences in Ischemic Cerebral Infarction: A Nationwide German Real-Life Analysis from 2014 to 2019. *Clin Transl Neurosci*.

- 2022 Sep 12;6(3):23. doi: 10.3390/ctn6030023
21. Sreevani S, Paramjyothi P. INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH CLINICAL PROFILE OF STROKE PATIENTS IN TERTIARY CARE CENTRE General Medicine Dr . Sapuri Sreevani * Dr . Pitta Paramjyothi ABSTRACT Diabetismellitus Dyslipidemia Hdl (high Density Lipoproteins) Hypertension S. Int J Sci Res. 2018;7(7):30–2. doi: <https://www.doi.org/10.36106/ijrs>
22. Itagi A, G. B, Biradarpatil P, Pattanashetti L. A study on clinical profile of acute stroke. Int J Adv Med. 2020 Nov 23;7(12):1809. doi: 10.18203/2349-3933.ijam20204966
23. Simmons CA, Poupore N, Nathaniel TI. Age Stratification and Stroke Severity in the Telestroke Network. J Clin Med. 2023 Feb 14;12(4):1519. doi: 10.3390/jcm12041519
24. Saeed S, Waje-Andreassen U, Naess H, Fromm A, Nilsson PM. The impact of age and 24-h blood pressure on arterial health in acute ischemic stroke patients: The Norwegian stroke in the young study. J Clin Hypertens. 2021 Oct 7;23(10):1922–9. doi: 10.1111/jch.14361
25. Stevano R, Margono JT, Sutanto A. Clinical Profile And Risk Factors Of Stroke: A Comparative Analytical Study Between Young And Old Onset. Magna Neurol. 2023 Jan 1;1(1):13–6. doi: 10.20961/magnaneurologica.v1i1.470
26. Krishnamurthi R V., Ikeda T, Feigin VL. Global, Regional and Country-Specific Burden of Ischaemic Stroke, Intracerebral Haemorrhage and Subarachnoid Haemorrhage: A Systematic Analysis of the Global Burden of Disease Study 2017. Neuroepidemiology. 2020;54(2):171–9. doi: 10.1159/000506396
27. Mac Grory B, Xian Y, Solomon N, Matsouaka R, Decker-palmer M, Fonarow GC, et al. Abstract 35: Unmet Need In Acute Ischemic Stroke Patients Arriving After 4.5 Hours. Stroke. 2022 Feb;53(Suppl_1). doi: 10.1161/str.53.suppl_1.35
28. Saver JL, Smith EE, Fonarow GC, Reeves MJ, Zhao X, Olson DM, et al. The “Golden Hour” and Acute Brain Ischemia. Stroke. 2010 Jul;41(7):1431–9. doi: 10.1161/STROKEAHA.110.583815
29. Ambarika R, Anggraini NA. Golden Period in Terms of Knowledge Family of Early Detection of Stroke for Stroke Patients in Installation of

- Emergency. Malaysian J Med Res. 2022;06(03):30–4. doi: 10.31674/mjmr.2022.v6i03.004
30. Retnaningsih R, Hendartono TK. PROFIL PASIEN STROKE ISKEMIK AKUT DENGAN TERAPI RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR DI RSUP DR. KARIADI SEMARANG. Maj Kedokt Neurosains Perhimpun Dr Spes Saraf Indones. 2020 Sep 29;36(4). doi: 10.52386/neurona.v36i4.86
 31. Dinata CA, Safrita YS, Sastri S. Gambaran Faktor Risiko dan Tipe Stroke pada Pasien Rawat Inap di Bagian Penyakit Dalam RSUD Kabupaten Solok Selatan Periode 1 Januari 2010 - 31 Juni 2012. J Kesehat Andalas. 2013 May 1;2(2):57. doi: 10.25077/jka.v2i2.119
 32. Gordon W, Ilorah C. Abstract 90: Meta-analysis Of Early Treatment Of Hypertension In Acute Ischemic Stroke And Functional Outcome. Stroke. 2022 Feb;53(Suppl_1). doi: 10.1161/str.53.suppl_1.90
 33. Mahapatra U, Ganai S, Pulai D, Saha D, Swaika BC. Distribution of aetiological types of stroke among the diabetic adults admitted with stroke in a referral hospital: an observational study. Int J Basic Clin Pharmacol. 2021 Jan 25;10(2):182. doi: 10.18203/2319-2003.ijbcp20210188
 34. Ambresh A, K R M. Association between HBA1C and NIHSS score in stroke. IP Indian J Neurosci. 2021 Sep 28;7(3):192–7. doi: 10.18231/j.ijn.2021.034
 35. Matz K, Tuomilehto J, Teuschl Y, Dachenhausen A, Brainin M. Comparison of oral glucose tolerance test and HbA1c in detection of disorders of glucose metabolism in patients with acute stroke. Cardiovasc Diabetol. 2020 Dec 5;19(1):204. doi: 10.1186/s12933-020-01182-6
 36. Wang YL, Leng XY, Dong Y, Hou XH, Tong L, Ma YH, et al. Fasting glucose and HbA1c levels as risk factors for the presence of intracranial atherosclerotic stenosis. Ann Transl Med. 2019 Dec;7(24):804–804. doi: 10.21037/atm.2019.12.56
 37. Rahman A, Dasgupta R, Quraishi F, Saha U, Ali Z, Hossain S, et al. Association between Plasma Homocysteine Level and Ischemic Stroke. J Bangladesh Coll Physicians Surg. 2014 Nov 27;31(3):128–33. doi: 10.3329/jbcps.v31i3.20978
 38. Kar A, Dash C, Murmu M, Mahesh K. P. K. Study of lipid profile in cases of non-diabetic stroke. Int J Adv Med. 2018 Jan 18;5(1):82. doi: 10.18203/2349-3933.ijam20180063
 39. Kanagarajan NG, Ningthoukhongjam

- R, Ningshen R, Singh TG, Khruomo V. A cross-sectional study of impact of dyslipidemia in stroke. *Int J Adv Med*. 2020 Oct 21;7(11):1670. doi: 10.18203/2349-3933.ijam20204518
40. Yaghi S, Elkind MSV. Lipids and Cerebrovascular Disease. *Stroke*. 2015 Nov;46(11):3322–8. doi: 10.1161/STROKEAHA.115.011164