


Dessy Dessy

1-The_effect_of_submaximal_exercise_

 The effect of submaximal

Document Details

Submission ID

trn:oid:::3618:116765324

Submission Date

Oct 15, 2025, 1:17 PM GMT+7

Download Date

Oct 15, 2025, 1:28 PM GMT+7

File Name

1-The_effect_of_submaximal_exercise_.pdf

File Size

299.1 KB

10 Pages

2,945 Words

15,170 Characters

2% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.





Filtered from the Report

- Bibliography
- Quoted Text




Exclusions

- 13 Excluded Matches

Match Groups

-  **4 Not Cited or Quoted 2%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 2%  Internet sources
- 0%  Publications
- 2%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 4 Not Cited or Quoted 2%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 2% Internet sources
- 0% Publications
- 2% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

- 1** **Student papers**
Universitas Katolik Widya Mandala Surabaya on 2025-09-29 **1%**
- 2** **Student papers**
University of Auckland on 2005-01-31 **<1%**
- 3** **Internet**
academic.oup.com **<1%**

THE EFFECT OF SUBMAXIMAL EXERCISE ON THE ATTENTION OF HEALTHY ADULTS

Kartikasari Desy^{1*}

*Corresponding author email:

DOI: <https://doi.org/10.33508/jwmj.v6i3.5645>

ABSTRACT

Introduction: Submaximal exercise uses intensity below 85% of the predicted maximum pulse for age. One example of a submaximal exercise is the Three-minutes step test (TMST) which is practical, easy to implement in the subject's routine, and has clear measurement standards. After TMST is done, VO_2 max can be calculated which is the main indicator used to determine cardiorespiratory capacity. There have been no studies evaluating the effect of submaximal exercise and VO_2 max on domain attention in healthy adults. **Aim:** This study aims to evaluate the effect of submaximal exercise on the attention of healthy adults. **Method:** Using a controlled before-and-after study, sixty healthy adults, aged 25-40 years, were divided into control and intervention groups (TMST). Attention function was measured with SDMT (Symbol Digit Modalities Test) before and after TMST or 3 minutes rest. **Result:** There was a significant difference in the intervention and control groups (p-values <0.001 and 0.0029). The values of SDMT in the intervention group with the control group were significant with the p-value <0.0001. The effect of the TMST intervention on SDMT was significant, with p-value <0.0001. **Conclusion:** Submaximal exercise can improve attention function in healthy adults.

Keywords: Submaximal Exercise, TMST, Attention, VO_2 max, Healthy Adults

¹ Department of Neurology, Faculty of Medicine, Widya Mandala Catholic University

INTRODUCTION

Cognitive function is the main human capital in activities of daily life. This function is divided into five major domains, namely attention, memory, visuospatial, language, and executive functions, which are constantly interconnected. Good cognitive function is vital for maintaining a satisfactory quality of life and also work performance.¹

Attention is an increase in brain activity in the form of sorting and categorizing received stimuli¹. Posner et al divide attention into three distinct pathways, namely alerting, orientation, and executive control. The latter pathway assists in the resolution of conflicts and decisions involving attention and mainly involves the anterior cingulate and frontal areas of the brain.²

Disorders of attention often occur among many professionals, especially in productive age groups such as early (26-35 years) and late adulthood (36-45 years). Several strategies have been developed to improve the function of attention, one of which is physical exercise, which can improve cognitive function through indirect and direct pathways. The indirect pathway affects systemic and central nervous system inflammation and increases cerebral blood flow. The direct pathway exerts its effect through 2 mechanisms, namely increasing neurotrophic development factors

(including BDNF/brain-derived neurotrophic factor) and increasing neurotransmitters (especially norepinephrine and dopamine). Increased BDNF will help neuroplasticity, neurogenesis, and neuroprotection.³

Submaximal exercise is a potential intervention that can be implemented for the improvement of cognitive function among professional young adults, as it is performed below 85% of the predicted maximum pulse according to age, and does not cause fatigue⁴. After submaximal exercises, the VO₂max value of individuals can be measured through the use of a simple formula. To date, VO₂ max is the main indicator used to determine cardiorespiratory capacity and function. Traditionally, this is measured by indirect calorimetry, subjects are required to perform a maximal test for causing fatigue and oxygen consumption. This test can be quite expensive and time-consuming. So, submaximal exercise can be an option that is more practical.⁴

Various studies have been developed showing a positive effect of submaximal exercise on improving cognitive function.^{3,5} A potential test is the three-minute step test, as it is practical, requires minimal time and equipment, has clear measurement standards, and is hence easily implemented in routine activities⁵. To date,

no studies have looked at the effects of TMST on cognitive function despite said qualities. In this study, we aimed to assess the effects of the three-minute step test (TMST) exercise on cognitive domains, is attention domains, of healthy young adult professionals.

METHOD

Study Design and Data Collection

This was a *controlled before-and-after study* conducted between February to April 2022 on healthy adults. The study has been approved by the Human Research Ethics Committee of the Medical Faculty of Hasanuddin University, South Sulawesi, Indonesia (Ethics number 125/UN4.6.4.5.31/PP36/2022, Protocol number UH22020051).

The inclusion criteria in this study were: (1) healthy adult working as professional medical staff, (2) age between 25-40 years, and (3) willing to participate in the study with signed informed consent. Criteria for exclusion include: (1) visual and hearing impairment, (2) diagnosed with any type of dementia, (3) had a comorbid disease, such as hypertension, diabetes mellitus, dyslipidemia, heart disease, (3) leg pain, (4) brain disorder, such as stroke, tumor, trauma, (5) pregnancy, (6) consumed sedative drugs and alcohol, (7) had consumed caffeine in <6 hours before the test, (8) after TMST, maximal pulse

>85%. Subjects who withdrew their willingness to participate in this study or failed to complete the entire series of the study procedure were considered dropout subjects.

Subjects were divided into 2 groups, namely the intervention group (TMST) and the control group (3 minutes rest). TMST was performed by asking the subject to move up and down the bench four steps per cycle (right leg up-left foot up-right foot down-left foot down). Within 3 minutes, male subjects had to complete 24 cycles regularly with the metronome set to 96 strokes per minute. Meanwhile, female subjects had to complete 22 cycles regularly with the metronome set to 88 strokes per minute for 3 minutes. Before and after TMST or resting for 3 minutes, physical and neurological examinations were carried out as well as measurements of attention using the Symbol Digit Modalities Test (SDMT).

Smartphone-Based Symbol Digit Span Modality Test (SDMT)

The Symbol Digit Modalities Test (SDMT) is a substitution test that is used specifically to evaluate attention⁶. Within 90 seconds, the subject must be able to match a specific number to the given geometric image. This test was invented by Smith in 1984⁷. The sensitivity and specificity of computerized SDMT were

71% and 84%, respectively, while the sensitivity and specificity of paper SDMT were 67% and 95%, respectively⁸.

In this mobile smartphone application, the initial display (Figure 1A) is the homepage with 'Start' to begin the test, 'Instruction' contains an explanation of the test steps, and selecting 'Language: English' will change the language to Indonesian. Figure 1B shows a test display that contains the keys, questions, and the numbers keypad after pushing the 'Start' button. The key contains numbers 1 until 9 with their pair of geometric symbols (randomly assigned to prevent memorization) above each number. Subjects were asked to fill in the blank box in the 'test' with numbers 1 to 9 according to the key. Subjects can empty the boxes that are considered difficult and jump to the next box by pressing the target box. Press 'next' to go to the next test page with the same key. The duration of this test is 90 seconds, questions automatically cannot be refilled when time runs out (Figure 1C) and by pressing 'Show Score', the number of correct answers will be seen (Figure 1D). When 'Back to Home' is selected, the test begins again with randomized symbols to avoid bias (immediate memory).

The assessment of the test is made by the number of correct answers. A decrease of 4 points or 10% or 0.5 standard deviations is considered significant

cognitive deterioration. On the other hand, an increase of 4 points or 10% or 0.5 standard deviation is considered an improvement in cognitive function³.




Figure 1. The SDMT mobile smartphone application: (A) Home screen, (B) The test with key above, (C) Time out, (D) Total score.

Measurement of VO₂max

In the intervention group, VO₂max was calculated using the formula for men = $70,597 - 0.246 \times (\text{age}) + 0.077 \times (\text{height}) - 0.222 \times (\text{weight}) - 0.147 \times (\text{pulse})$, or women = $70,597 - 0.185 \times (\text{age}) + 0.097 \times (\text{height}) - 0.246 \times (\text{weight}) - 0.122 \times (\text{pulse})$. VO₂max is then compared with the difference between SDMT and BDS.

Data Analysis

All data was analyzed using GraphPad Prism v9. Changes in scores

before and after treatment (intervention group) and after rest (control group) on each test were assessed using the Wilcoxon test. Comparison of the Δ SDMT between the intervention and control group was assessed using the Mann-Whitney test. The chi-square analysis was used to assess associations between SDMT with submaximal exercise in intervention group, compared with the control group. In the intervention group, see relationship between the VO₂max value with the SDMT was assessed using the Spearman collaboration. A p-value <0.05 was considered to indicate statistical significance for all tests.

RESULT

Table 1 shows the distribution of subjects based on gender, age, BMI, time, and type of exercise performed in a week. In the intervention group, the number of male subjects was 16 people (53.3%) while

the female subjects were 14 people (46.7%). Meanwhile, in the control group, the number of male and female subjects was the same, each 15 people (50%). The mean age in the intervention group was 32.23 years, while in the control group was 32.23 years.

Most of the subjects (70% in the intervention group and 66.7% in the control group) exercised 1 to 150 minutes in 1 week. There were 16.7% in the intervention group and 23.3% in the control group who never exercised at all. There were 13.3% in the intervention group and 10% in the control group who exercised >150 minutes in 1 week. The main mode of exercise in was jogging, performed by 43.4% of subjects in the intervention group and 40% in the control group. In addition, other common exercises include football, badminton, cycling, HIIT, swimming, fitness, table tennis, and yoga.

Table 1. Baseline characteristics of study subjects.

Variable	Intervention Group		Control Group		p value
	n	%	n	%	
Gender					
Male	16	53,3%	15	50%	>0,9999
Female	14	46,7%	15	50%	
Age					
25-29 years	4	13,3%	8	26,6%	0,2185
30-34 years	19	63,3%	13	43,3%	
35-39 years	6	20%	9	30%	
Mean	32,23 ± 2,64		32,23 ± 3,18		
One week exercise					0,9069
0 minutes	5	16,7%	7	23,3%	

Variable	Intervention Group		Control Group		p value
	n	%	n	%	
1-150 minutes	21	70%	20	66,7%	
>150 minutes	4	13,3%	3	10%	
Type of exercise					
Cycling	1	3,3%	1	3,3%	
HIIT	1	3,3%	1	3,3%	
Badminton	2	6,7%	5	16,7%	
Football	8	26,7%	6	20%	
Jogging	13	43,4%	12	40%	
Swimming	1	3,3%	0	0%	
Fitness	1	3,3%	0	0%	
Table tennis	1	3,3%	0	0%	
Yoga	0	0%	1	3,3%	
Heart rate					
• Before (mean)	85,43		77,97		
• After (mean)	115,37		77,60		

A significant change in the SDMT score of subjects was observed in both the intervention (p-value < 0.001) and control (p-value < 0.0029) groups can be seen in **Figure 2**. It is apparent that the intervention results in a larger improvement of SDMT scores relative to the control.




Figure 2. SDMT Score on Intervention and Control Group

The Mann-Whitney test was then used to compare the values of Δ SDMT in the intervention group with the control group, with a result of p < 0.0001 for Δ SDMT (**Figure 5**). Chi-square analysis was also performed to see an effect between attention function with the TMST intervention. The p-value < 0.0001 on the SDMT score was obtained. This indicates that there is an influence between attentional function with the TMST intervention.




Figure 3. Comparison of Δ SDMT scores between interventional and control groups.

DISCUSSION

Regular exercise, especially aerobic exercise, can increase the efficiency of the aerobic energy production system by increasing maximal oxygen uptake and cardiorespiratory endurance. Physical activity can increase the activity of the prefrontal cortex, namely attention, short-term memory, problem-solving, cognitive flexibility, decision-making, and inhibition control. Acute exercise or just one exercise can increase the activity of the prefrontal cortex (related to cognitive function), improve mood, and reduce stress levels (Basso, J. & Suzuki, W., 2017).

WHO in 2020 recommended doing at least 150-300 minutes of moderate aerobic physical activity or 75-150 minutes of vigorous aerobic activity in 1 week. The subjects of this study were only 7 people (11.7%) who exercised regularly ≥ 150 minutes in 1 week. This shows that the research subjects are still exercising less than what the body needs.

The data above shows that there are only 10 subjects (16.7%) who have a normal backward digit span at the beginning of the examination. The rest have short-term memory disorders. Therefore, it is necessary to do physical exercise to improve cognitive function. The physical exercise carried out in this study was TMST submaximal exercise.

In this study, there were significant differences in SDMT scores before and after the intervention, with p-values < 0.0001 . In the control group, there was a significant difference in SDMT scores before and after a 3-minute rest with a p-value of 0.029. However, the p-value in the control group was higher than that in the intervention group.

The significant difference in the SDMT test in the control group was probably caused by the time gap between the pretest and post-test examinations which was too close (3-5 minutes) so that the subject still had an immediate memory of the questions being worked on. In addition, it may also be because the subject has become accustomed to working on the problem so that he is more capable. In the SDMT test application, the final score is immediately displayed. This allows the subject to be motivated to work on the SDMT questions better on the post-test.

Chang, Y, et al. 2014 investigated the effect of acute exercise on cognitive function and found a positive effect, especially on the domains of attention and executive function, on all fitness levels of young adults⁹. While Manadhar et al. in his 2020 study in Nepal on the Effect of the Three Minutes Step Test on Cognitive Function of Medical Students concluded that there was an increase in cognitive

function in the domains of attention and executive function as assessed by the Stroop test⁵. This is consistent with this study, which also found an increase in cognitive function, but in a different domain (attention). The increase in cognitive function in this study may be associated with an increase in blood flow which will improve brain health, one of which is cognitive function³.

Research conducted by Flodin P, et al. in 2017 regarding the effect of VO₂max on cognitive function showed that VO₂max is not related to cognitive function, possibly due to differences in imaging tools with previous studies, differences in study design and analysis with previous studies, and different types of exercise¹⁰. In our study, the type of exercise that the subjects did in the control and intervention groups also varied.

CONCLUSION

From this study, it can be concluded that there are differences in changes in SDMT scores between the intervention and control groups, and there is an effect of submaximal exercise on increasing SDMT scores in healthy adults. Therefore, TMST (Three Minutes Step Test) can be performed in daily life to improve short-term memory function and attention.

REFERENCES

1. Anindhita T, Wiratma W. *Buku Neurologi Ajar*. 1st ed. (Jakarta, ed.). Departemen Neurologi FKUI; 2017.
2. Hussain F, Wood S. Modelling the Efficiencies and Interactions of Attentional Networks. *Lect Notes Comput Sci*. 2009;5395.
3. Canepa P, Pedullà L, Bisio A, Ruggeri P, Bove M. Is the 12 minute-walk/run test a predictive index of cognitive fitness in young healthy individuals? A pilot study on aerobic capacity and working memory in a real-life scenario. *Neurosci Lett*. Published online 2020:728.
4. Kieu NT Van, Jung SJ, Shin SW, et al. The Validity of the YMCA 3-Minute Step Test for Estimating Maximal Oxygen Uptake in Healthy Korean and Vietnamese Adults. *J Lifestyle Med*. 2020;10(1):21-29.
5. Manandhar S, Chettri S, Pandey KR, Limbu N, Baral D, Pandey DR. EFFECT OF THREE MINUTE STEP TEST ON COGNITION AMONG MEDICAL STUDENTS. *MNJ (Malang Neurol Journal)*. 2021;7(2):120-124.
6. Garcia S, Alosco ML, Spitznagel MB, et al. Cardiovascular fitness associated with cognitive performance in heart failure patients

- enrolled in cardiac rehabilitation.
BMC Cardiovasc Disord. Published online 2013.
7. Kreutzer JS, DeLuca J, Caplan B. *Encyclopedia of Clinical Neuropsychology*. 2nd ed. (Kreutzer JS, DeLuca J, Caplan B, eds.). Springer Cham; 2018.
 8. Akbar N, Honarmand K, Kou N, Feinstein A. Validity of a computerized version of the symbol digit modalities test in multiple sclerosis. *J Neurol.* Published online 2011:258.
 9. BA W, A C, C A, et al. *Current Diagnosis & Treatment: Geriatrics, Second Edition*. McGraw-Hill Education; 2014.
<https://accessmedicine.mhmedical.com/content.aspx?bookid=953§ionid=53374629>
 10. Basso JC, Suzuki WA. The Effects of Acute Exercise on Mood, Cognition, Neurophysiology, and Neurochemical Pathways: A Review. *Brain Plast.* 2017;2(2):152.