

# Yudhiakuari Sincihu

## 3-The Method\_occurrence\_health\_

 Yudhiakuari Sincihu3

---

### Document Details

Submission ID

trn:oid::3618:129709528

Submission Date

Mar 3, 2026, 8:47 AM GMT+7

Download Date

Mar 3, 2026, 2:07 PM GMT+7

File Name

3-The Method\_occurrence\_health\_.pdf

File Size

955.1 KB

16 Pages

10,691 Words

59,503 Characters

# 8% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

## Exclusions

- ▶ 35 Excluded Sources
- ▶ 205 Excluded Matches

## Match Groups

-  **46 Not Cited or Quoted 4%**  
Matches with neither in-text citation nor quotation marks
-  **41 Missing Quotations 4%**  
Matches that are still very similar to source material
-  **0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

## Top Sources

- 3%  Internet sources
- 6%  Publications
- 4%  Submitted works (Student Papers)

## Integrity Flags

0 Integrity Flags for Review

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

### Match Groups

- 46 Not Cited or Quoted 4%**  
Matches with neither in-text citation nor quotation marks
- 41 Missing Quotations 4%**  
Matches that are still very similar to source material
- 0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

### Top Sources

- 3% Internet sources
- 6% Publications
- 4% Submitted works (Student Papers)

### Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

<b>1</b>	Publication	Anna Ida Sunaryo Purwiyanto, Tri Prartono, Ety Riani, Yuli Naulita, Muhammad R...	<1%
<b>2</b>	Publication	Fatemeh Sadat Kashfi, Bahman Ramavandi, Hossein Arfaeinia, Azam Mohammadi...	<1%
<b>3</b>	Publication	Longyi Shao, Yaowei Li, Tim Jones, M. Santosh et al. "Airborne microplastics: A rev...	<1%
<b>4</b>	Publication	Nanthi S. Bolan, M.B. Kirkham, Claudia Halsband, Dayanthi Nugegoda, Yong Sik O...	<1%
<b>5</b>	Publication	Alvise Vianello, Rasmus Lund Jensen, Li Liu, Jes Vollertsen. "Simulating human ex...	<1%
<b>6</b>	Student papers	Higher Education Commission Pakistan on 2025-05-03	<1%
<b>7</b>	Publication	Paramastri Rahmi Syafina, Adyati Pradini Yudison, Emenda Sembiring, Mohamma...	<1%
<b>8</b>	Student papers	Indonesia International Institute for Life Sciences on 2025-12-18	<1%
<b>9</b>	Publication	Xinran Zhao, Yupeng Zhou, Chenzhe Liang, Jianchen Song, Siyun Yu, Gengxuan Li...	<1%
<b>10</b>	Internet	sajs.co.za	<1%

11	Publication	"Micro-Nano Plastics Exposure, Environmental Degradation and Public Health Cri...	<1%
12	Student papers	Wayne State University on 2025-10-06	<1%
13	Internet	www.cwejournal.org	<1%
14	Publication	"Microplastics: Ecological and Food Security Implications", Springer Science and B...	<1%
15	Student papers	Asian Institute of Technology on 2024-05-05	<1%
16	Student papers	Universiti Malaysia Terengganu UMT on 2025-06-16	<1%
17	Publication	"Microplastics in African and Asian Environments", Springer Science and Business...	<1%
18	Student papers	Asian Institute of Technology on 2021-07-23	<1%
19	Internet	www.dovepress.com	<1%
20	Student papers	Finger Lakes Community College on 2024-10-20	<1%
21	Publication	Lei Su, Bingxu Nan, Nicholas J. Craig, Vincent Pettigrove. "Temporal and spatial v...	<1%
22	Publication	Udomratana Vattanasit, Jira Kongpran, Atsuko Ikeda. "Airborne microplastics: A ...	<1%
23	Publication	Horgnies, M., and J.J. Chen. "Superhydrophobic concrete surfaces with integrated ...	<1%
24	Student papers	Rutgers University, New Brunswick on 2023-02-13	<1%

25	Publication	Tiffany Eberhard, Gaston Casillas, Gregory M. Zarus, Dana Boyd Barr. "Systematic ...	<1%
26	Student papers	University of Arizona on 2025-06-04	<1%
27	Student papers	Cork Institute of Technology on 2024-12-11	<1%
28	Publication	Rahmi Amtha, Ferry Sandra, Rosalina Tjandrawinata, Indrayadi Gunardi, Anggrae...	<1%
29	Publication	Tonni Agustiono Kurniawan, Abdelkader Anouzla. "Handbook of Microplastic Poll...	<1%
30	Internet	ndl.ethernet.edu.et	<1%
31	Publication	Nasim Zolfaghari Firouzsalari, Farhad Ghayurdoost, Akbar Gholampour. "A narrat...	<1%
32	Internet	www.canada.ca	<1%
33	Internet	www.researchgate.net	<1%
34	Internet	www.springerpflege.de	<1%
35	Student papers	Coventry University on 2020-08-21	<1%
36	Publication	Guanglong Chen, Qingyuan Feng, Jun Wang. "Mini-review of microplastics in the ...	<1%
37	Publication	Huirong Yang, Yinglin He, Yumeng Yan, Muhammad Junaid, Jun Wang. "Character...	<1%
38	Publication	Steigvilė Byčėnkienė, Ieva Uogintė, Lina Davulienė, Sergej Šemčuk et al. "Compar...	<1%

39	Student papers	University College London on 2025-01-16	<1%
40	Student papers	University of Bristol on 2023-05-17	<1%
41	Student papers	University of Gloucestershire on 2021-05-16	<1%
42	Student papers	Unversity of Christchurch International College on 2020-07-30	<1%
43	Publication	Xuan Zhu, Wei Huang, Mingzhu Fang, Zhonglu Liao et al. "Airborne Microplastic C...	<1%
44	Internet	enveurope.springeropen.com	<1%
45	Publication	"Microplastic Pollution", Springer Science and Business Media LLC, 2021	<1%
46	Publication	A. Torres-Agullo, A. Karanasiou, T. Moreno, S. Lacorte. "Overview on the occurren...	<1%
47	Student papers	Heriot-Watt University on 2024-05-28	<1%
48	Student papers	Kennesaw State University on 2020-07-17	<1%
49	Publication	Kexin Zheng, Peng Wang, Xiaoming Lou, Zhen Zhou et al. "A review of airborne m...	<1%
50	Student papers	Mapua Institute of Technology on 2024-08-21	<1%
51	Publication	MohammadAmin Azizi, Zahra Khashi, Razieh Ashoori, Farzaneh Rostami, Aboalf...	<1%
52	Publication	Pei-Wei Yeh, Chien-Cheng Jung. "Concentrations, characteristics, influencing fact...	<1%

53	Student papers	Southern Illinois University Edwardsville on 2025-04-16	<1%
54	Student papers	University of Hertfordshire on 2024-11-11	<1%
55	Student papers	University of Technology, Sydney on 2022-11-11	<1%
56	Internet	e-journal.lp2m.uinjambi.ac.id	<1%
57	Internet	eaht.org	<1%
58	Internet	ebin.pub	<1%
59	Internet	ebuah.uah.es	<1%
60	Internet	www.cetjournal.it	<1%
61	Publication	Eunsu Lim, Hirofumi Tanaka, Yuan Ni, Yifan Bai, Kazuhide Ito. "Microplastics/micr...	<1%
62	Publication	Yaowei Li, Longyi Shao, Wenhua Wang, Mengyuan Zhang, Xiaolei Feng, Wenjun Li...	<1%
63	Publication	Feride Öykü Sefiloglu, Marthinus Brits, Martin J. M. van Velzen, Eelco N. Pieke, On...	<1%

## SYSTEMATIC REVIEW

# The Method, Occurrence, Health Risk, and Prevention of Airborne Microplastics in Indonesia: A Systematic Review

Kusuma S. Lestari<sup>1</sup>, M. Zainul Asrori<sup>2</sup>, Yudhiakuari Sincihu<sup>3</sup>, Gerry Andhikaputra<sup>4</sup>, Gondo Mastutik<sup>5</sup>

<sup>1</sup> Department of Environmental Health, Faculty of Public Health, Universitas Airlangga, 60115 Mulyorejo, Surabaya Indonesia

<sup>2</sup> Department of Physics, Faculty of Science and Data Analytics, Institut Teknologi Sepuluh Nopember, 60111 Sukolilo, Surabaya Indonesia

<sup>3</sup> Faculty of Medicine, Widya Mandala Surabaya Catholic University, 60112 Mulyorejo, Surabaya Indonesia

<sup>4</sup> Department of Environmental Engineering, College of Engineering, Chung Yuan Christian University, 320314 Taoyuan, Zhongli, Taiwan

<sup>5</sup> Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga, 60132 Tambaksari, Surabaya, Indonesia

## ABSTRACT

**Introduction:** Microplastics originate from production in small forms (micro) and degraded plastic waste. Initially, most research on microplastics focused on their presence in oceans and lakes. Recently, researchers have begun investigating airborne microplastics, as their size, ranging from 500 to 2000  $\mu\text{m}$ , allows them to be carried by the wind. There is a paucity of research on airborne microplastics conducted in Indonesia. This study aims to evaluate the current level of airborne microplastic research in Indonesia. **Materials and methods:** The method involved searching the literature that included the terms "airborne," "atmospheric," or "ambient microplastics" in Indonesia, following PRISMA guidelines. Data sources used the Google Scholar database from 2019 to 2024 and search papers from May 15, 2024, until August 12, 2024. A total of 15 studies were reviewed. **Results:** All studies reviewed confirmed the presence of airborne microplastics, with fibers and fragments being the most common forms, varying in size, and polyethylene terephthalate (PET), polyethylene (PE), and polyester being the most frequently identified polymers. **Conclusion:** The sampling methods used were active and passive; variations in location, time, duration, and equipment can influence the outcomes. Additionally, factors such as weather conditions, research site characteristics, and human activities also have an impact. The health risks associated with the tiny size of microplastics are similar to those of particulate matter, which can easily enter the respiratory system and cause non-communicable diseases. There is a need to establish threshold levels for microplastic concentrations in the air, both indoors and outdoors. *Malaysian Journal of Medicine and Health Sciences* (2025) 21(SUPP7): 220-235. doi:10.47836/mjmhs.21.s7.26

**Keywords:** Airborne microplastic, Method, Occurrence, Health risk, Prevention

## Corresponding Author:

Kusuma S. Lestari, M.KM.

Email: kusumalestari@fkm.unair.ac.id

Tel : +62 821-1551-1145

## INTRODUCTION

The issue of plastic pollution was first raised many years ago, (1) with the initial discovery of plastic fragments of plastic in the stomachs of *albatrosses*, an arge seabird (2). Ingestion of plastics occurs in fish, seabirds, marine animals, and various food chain members (3). These instances involve not only the toxicity of plastics but also physical interference, such as entanglement in nets and lines, choking after consuming plastic debris, or ingesting plastic, which can lead to a false sense of fullness and ultimately starvation (1). The term "microplastics" (4) was

first introduced in 2004, referring to granular and fibrous plastic particles with a diameter of 20  $\mu\text{m}$  collected from Currently, microplastics are prevalent in various environments, particularly beaches and the ocean. Microplastics have been recognised as a significant contributor to biodiversity loss and were included in the UNEP Year Book 2014 as one of ten emerging environmental issues (5). Currently, microplastics are prevalent in various environments, particularly in the oceans (4), the deep ocean (6), freshwater lakes (7), rivers (8–10), and terrestrial ecosystems (11,12).

The global expansion in plastic manufacturing over the past few decades has led to a significant increase in the quantity of microplastic pollution in the environment, a trend that is expected to persist (13). Improperly managed plastic waste can ultimately find

45  
58  
15  
53  
its way into water bodies. Each year, millions of tons of plastic waste from 192 coastal nations enter the ocean, and this figure is anticipated to rise in tandem with increasing plastic production. (14,15). According to a study conducted across 192 countries, Indonesia ranks second in contributing plastic waste to marine environments (11). Plastic particles can become trapped in or on bubbles formed at the surface due to wave action, and when these bubbles burst upon contact with the air, they release contaminants, including plastic particles, into the atmosphere (16). Once in the atmosphere, plastic particles can be transported by the wind over considerable distances, similar to dust, sand, or sea salt (17). Depending on their density and weight, microplastics can be carried by air or water currents throughout the environment (18).

25  
25  
19  
16  
Airborne microplastics were first identified by Dris in Paris, France, in 2015, confirming the presence of microplastics in freshwater, sewage, and atmospheric fallout in urban environments (19). Subsequent research has been conducted across Europe, including studies in France (Paris (20,21), Pyrenees Mountains (22); UK in London (23), Nottingham (24), Yorkshire (25); Germany in Hamburg (26); Denmark in Aarhus (27); Spain in Barcelona (28); Portugal in Aveiro (29); US in California (30); Australia in Sydney (31), Victoria (32); Asia: China in Dongguan (33), Yantai (34), Shanghai (35–37), Beijing (38), Wenzhou (39); Iran in Tehran (40), Asaluyeh (41), Bushehr port and Shiraz (42); Turkey in Sakarya (43); India in Chennai (44), Nagpur (45); Nepal in Kathmandu (46); Japan in Kusatsu (46); and the Arctic regions of the Swiss Alps and Bremen, Bavaria (47). Research on airborne microplastics in Southeast Asia has also been conducted in several countries, including Vietnam, in Da Nang (46) Ho Chi Minh (48); Malaysia in Klang, Petaling and Gombak (49), Kuala Lumpur (50), Kuala Lumpur (51), Trengganu (52); and Thailand in Bangkok, Pathum Thani, and Samut Prakan (53), Pathum Thani (54), Nakhon Si Thammarat (55), Pathum Thani (56), Bangkok (57).

61  
In addition, several studies have investigated the differences in airborne microplastics between indoor and outdoor environments (21,29,30,39,57), as well as between urban and suburban areas (20,26,32,39,45). Research has also focused on the atmospheric deposition rates of microplastics (19,20,23,31,33–35,52,56), their abundance in the air (19,20,36–42,44–46,21,47,48,51,55,57,22,25,26,28–30,32), and their various shapes (19,20,30–36,38–40,22,41–47,49–51,23,52–55,57,24–29). Additionally, studies have examined the sizes (19,20,39,40,45–47,50,53–56,22,57,23,26,27,30,31,34,38), and colors (34,35,52,55,40,41,43,45,46,49–51) of these polymer particles (20,21,33–36,40,42–46,23,47,53,57,25–28,30–32).

Airborne microplastics can easily disperse and

contribute to air pollution, as well as exacerbate global warming and climate change. It is noted that many airborne microplastics may increase the risk of global warming and climate change (58). These microplastics are so minuscule that they can be readily ingested and inhaled by living organisms (33,39). Numerous studies have examined microplastics in water, food, the sea, and lakes. Unfortunately, research on airborne microplastic in Indonesia remains limited, particularly concerning the quantification of airborne microplastics, their sources, and their impact on human health. Furthermore, policies related to microplastics, waste management, and both indoor and outdoor pollution are not sufficiently regulated. As the fourth most populous country, Indonesia ranks as the seventh-largest producer of waste globally. This study aims to evaluate the existing research on airborne microplastics in Indonesia, focusing on methods, prevalence, health risks, and prevention strategies.

## MATERIALS AND METHODS

### Source

All published, peer-reviewed journal articles focusing on microplastics were systematically searched using Boolean operators following PRISMA guidelines.

### Formulation of the Research Question

The research question for this systematic review was formulated using the PICO framework. A modified version of PICO, referred to as Problems, Interest, and Context, was employed. The review of studies must include three primary components: microplastics in the air (problems), their shapes, sizes, colors, and polymers (interest), and the geographical context of Indonesia.

### Eligibility Criteria

Papers written in both English and Indonesian were included in the search; however, mini-theses and theses were excluded.

### Information Sources

To locate published papers on airborne microplastics in Indonesia, a web search was conducted using Google Scholar, focusing on publications from the past five years (2019–2024). Google Scholar was chosen to broaden the search for relevant papers that align with the research criteria, given the study's exclusive focus on Indonesia.

### Search Strategy

The database search was conducted using the following search phrases with Boolean operators: "Airborne AND Microplastic AND in Indonesia," "Atmospheric AND Microplastic AND in Indonesia," "Ambient AND Microplastic AND in Indonesia," and "Mikroplastik di Udara AND Indonesia," yielding approximately 1,940 results, 6,250 results, 2,120 results, and 857 results, respectively.

1

**Selection Process**

A preliminary screening based on titles was conducted on all retrieved publications. Duplicate papers were removed, and those that were not freely accessible were excluded from this analysis (N = 27). A total of 22 free full-text papers were downloaded. Further screening led to the exclusion of additional papers, resulting in 15 articles available for review. The data presented are categorized by location of study, sampling methods, identification and quantification of microplastics, occurrence of airborne microplastics—including shapes, sizes, colors, and polymers—atmospheric microplastic deposition rates, total suspended particles (TSP), PM10, PM2.5, and microplastic concentrations; as well as health risks and prevention strategies. Figure 1. illustrates a diagram of the literature selection process.

59

7

28




Figure 1: PRISMA Flow Chart

**RESULTS**

**Location Site**

Research on airborne microplastics has been conducted in six major cities across Indonesia: 13 sites on Java Island and two on Sumatra Island. The study on Java Island included Jakarta, Bandung, Surabaya, Yogyakarta, Sidoarjo, Mojokerto, and Gresik, while the Sumatra Island locations were Medan and Bandar Lampung. This research took place from 2019 to 2023. The geographical distribution of the study locations across Indonesia is depicted in Figure 2.

16

55

27




Figure 2: Location Site (Pink province level; Orange city level)

In Jakarta, microplastic sampling was conducted in Ancol, North Jakarta, adjacent to Jakarta Bay. The sampling occurred at a high elevation to capture atmospheric microplastics without interference from trees or buildings (59). In Bandung, in 2019 (60), sampling was first conducted in Sarana Olahraga Ganesha (SARAGA), a green belt area designed to mitigate debris from passing traffic. The monitoring area was chosen due to its bowl-like topography and central location, which restricts horizontal air circulation, leaving vertical convection as the primary mechanism for air movement. That same year (61), a comparison of airborne microplastics was undertaken in two distinct spatial typologies within the Bandung Metropolitan Area: the Lebak Siliwangi area, representing an urban area, and the Cibabat area, representing a suburban area. In the following year (2022-2023), sampling occurred on the roof of an 8-story building at the Institute of Technology Bandung (ITB) and was compared to Osaka in 2021 on the roof of a 3-story building at Osaka Metropolitan University (62). This study was conducted across two cities in two nations: Bandung, Indonesia, and Osaka (Sakai City), Japan. Airborne microplastic research in Yogyakarta (63) and Surabaya (64, 64,65) was carried out along roads with high vehicle congestion due to microplastics produced from rubber tire friction (66). In Yogyakarta, sampling took place at four locations along the intersections of Yogyakarta's arterial road, known as the ring road (north, south, east, and west). In Surabaya (2019), sampling was performed on three streets: Urip Sumoharjo Street, representing a road with high vehicle congestion; Mayjend Sungkono Street, representing a road with medium congestion; and Embong Malang, representing a road with low congestion. Other studies (67, 67,68) were conducted indoors in three different settings (office, school, and apartment) to analyze the quantity of microplastics and the number of inhabitants in various indoor spaces. In Medan (69), sampling was conducted in six industrial areas to analyze outdoor airborne microplastics. Meanwhile, in Bandar Lampung (70), sampling was carried out near three roads representing different environments: industrial areas, residential zones, busy roads such as provincial roads, and the city center. In Sidoarjo, Gresik, and Mojokerto (71–74), sampling took place at various sites, including a tofu industry, a plastic waste recycling plant, a waste treatment industry incinerator, a temporary landfill, and a Waste to Energy Power Plant.

**Sampling**

There are two types of sampling methods: active and passive (75). Active sampling methods were employed in Bandung, Yogyakarta, Surabaya, Medan, and Bandar Lampung. All devices used were High Volume Air Samplers (HVAS) with different sampling durations: 24 hours (HVAS SIBATA HV 1000F) (61,63), 8 hours (70), and 3 hours (60). The flow rates varied among studies, including 68.82 m<sup>3</sup>/h (60), 20 L/min (62), 1.2

13 m<sup>3</sup>/second (63), 207.38 mL/min, and 446.93 mL/min (64,65). Vacuum pumps (Vacuubrand pumps) and a vacuum cleaner (Krisbow Turbo Tiger) were used in Surabaya (64,65,67,68). In Bandung, the sampling aimed to measure both microplastics and Total Suspended Particulate (TSP) (60,61). In both Bandung and Osaka, a multi-nozzle cascade impactor sampler (Tokyo Dylec) was used to collect three size fractions of ambient aerosols (PM<sub>2.5</sub>, PM<sub>10</sub>, and TSP) on Teflon-coated glass fiber filters (TX40HI20-WW, Pallflex) (62). In Yogyakarta, TSP, PM<sub>10</sub>, and PM<sub>2.5</sub> were measured (63), while in Medan, only PM<sub>2.5</sub> was assessed (69).

1 In Jakarta, for instance, 10 mL of distilled water is added to the rain gauge to collect atmospheric microplastics. This water is tested using the blank method to ensure it is free of microplastics. The rain gauge, with its mouth opening angled at 15° against the wall, operates for ninety-six hours (59). Sampling involves cutting

Whatman filter paper to fit petri dishes with diameters of 12 or 15 cm. The coated petri dishes are then placed in various locations according to the cardinal directions. In Sidoarjo, Gresik, and Mojokerto, the passive sampling methods are employed not only to measure microplastics but also PM<sub>2.5</sub> levels (71–74).

### Identification and Quantification of Microplastics

The identification of microplastics was conducted using various microscopes, including a binocular microscope (69,70), a Leica M205C (59), an Olympus UTV0.5XC-3 (63,64), a DinoLite AM2111 digital microscope (60,61,67,68), a stereo microscope (71,72,74) and a trinocular digital microscope Dw-tc-y Black edition (73). The occurrence of airborne microplastics, including shapes, sizes, colours, polymers, as well as the atmospheric microplastic deposition rate, Total Suspended Particles (TSP), PM<sub>10</sub>, PM<sub>2.5</sub> and microplastic concentration is presented in Table I.

**Table I: Occurrence of Airborne Microplastic in Indonesia 2019 – 2024**

Study Location	Shape	Size	Colour	Polymer	TSP Concentration	Deposition Rate	Concentration	Year	Reference
Jakarta	foam<fragment<fibres	500–1000 µm < 300–500 µm	N/A	<ul style="list-style-type: none"> <li>•Polyester,</li> <li>•Polystyrene (PS),</li> <li>•Polybutadiene, and</li> <li>•Polyethylene (PE).</li> </ul>	N/A	<ul style="list-style-type: none"> <li>•3-40 particles/m<sup>2</sup>/d, average of 15 ± 13 particles/m<sup>2</sup>/d</li> <li>•The highest mean in the rainy season (23.422 ± 9.065 particles/m<sup>2</sup>/d), and the lowest in the dry season (5.745 ± 1.984 particles/m<sup>2</sup>/d).</li> </ul>	N/A	2022	(52)
Bandung	Fibres in an urban area is greater than in the suburban area	<ul style="list-style-type: none"> <li>•Fibres size in urban samples is 1000–1400 µm</li> <li>•Fibres size in suburban area, the range is 600–1000 µm.</li> </ul>	<ul style="list-style-type: none"> <li>•Black,</li> <li>•brown,</li> <li>•red,</li> <li>•green, and</li> <li>•transparent</li> </ul>	N/A	<ul style="list-style-type: none"> <li>•Urban area TSP average: 186.52 ± 45.76 µg/m<sup>3</sup></li> <li>•Suburban area TSP average: 154.30 ± 74.02 µg/m<sup>3</sup></li> </ul>	N/A	<ul style="list-style-type: none"> <li>•Urban: 0.3–0.6 particles/m<sup>3</sup></li> <li>•Suburban: 0.1–0.3 particles/m<sup>3</sup></li> </ul>	2022	(53)
Bandung	Fibres and fragment	N/A	N/A	N/A	TSP: 78.28 + 2.50 µg/Nm <sup>3</sup>	N/A	N/A	2019	(54)

CONTINUE

**Table 1: Occurrence of Airborne Microplastic in Indonesia 2019 – 2024 (CONT.)**

Study Location	Shape	Size	Colour	Polymer	TSP Concentration	Deposition Rate	Concentration	Year	Reference
Bandung and Osaka	Bandung and Osaka: Fragment, granule, fibres (<fragment)	•Bandung: 3.14 - 512 µm •Osaka: 3.50 - 728.15 µm	N/A	Bandung: •Ethylene Vinyl Acetate (EVA), •Polybutadiene Rubber (PBR), •Polycaprolactone (PCL), •Polydiallyl Phthalate (PDAP), •Polyethylene (PE), •Polyethylene Terephthalate (PET), and •Polyvinyl Acetate (PVA).  Osaka: •Ethylene Vinyl Acetate (EVA), •Polyethylene (PE), •PE/Polypropylene (PP) Copolymer, •Polyethyl Methacrylate (PEMA), •Polyethylene Terephthalate (PET), •Polymethyl Methacrylate (PMMA), •Polypropylene (PP), •Polystyrene (PS), and •Polyvinyl Chloride (PVC).	N/A	N/A	•Bandung:1.03 to 14.27 particles/m <sup>3</sup> •Osaka:0.63 to 3.29 particles/m <sup>3</sup> Average: •Bandung: 6.64 particles/m <sup>3</sup> •Osaka:1.65 particles/m <sup>3</sup>	2023	(55)
Yogyakarta	Fibres< film<fragment	N/A	•Black, •Blue, •Brown, •Clear/ Transparent, •Green, •Orange, •Purple, •Red, And •Yellow	N/A	N/A	N/A	Four locations (road) are •0.86 particles/ Nm <sup>3</sup> •0.82 particles/ Nm <sup>3</sup> •0.59 particles/ Nm <sup>3</sup> •0.42 particles/ Nm <sup>3</sup>	2024	(56)
Surabaya	Fibres, fragments and film	N/A	N/A	•Polyethylene Terephthalate (PET), •Polyester, and •Cellophane	N/A	N/A	•First location: 174.97 and 130.50 particles/ m <sup>3</sup> •Second location: 131.75 and 68.36 particles/m <sup>3</sup> •Third location: 94.69 and 55.93 particles/m <sup>3</sup>	2019	(57,58)

CONTINUE

63

18

**Table 1: Occurrence of Airborne Microplastic in Indonesia 2019 – 2024 (CONT.)**

Study Location	Shape	Size	Colour	Polymer	TSP Concentration	Deposition Rate	Concentration	Year	Reference
35 Surabaya	Film<frag- ment<fi- bres	3000–3500 µm	N/A	N/A	N/A	N/A	Amount of micro- plastics Office: 1.334 particles on a weekday 242 particles on a weekend 2.351 particles on a weekday 252 particles on a weekend School 1.290 particles on a weekday and 239 particles on a weekend 2.321 particles on a weekday and 257 particles on a weekend Apartment 1.133 particles on a weekday and 127 particles on a weekend 2.108 particles on a weekday and 95 particles on a weekend	2020	(60)
31  39 Surabaya	Office: fibres and fragments  School and apartment: fibres	N/A	N/A	•Cellophane, •Clipboard, •Alkyd Resin, •Polyester, •Polyethylene Terephthalate (PET)	N/A	N/A	Office: •709.09 particles/ m <sup>3</sup> on Monday and Tuesday, •1186.36 parti- cles/m <sup>3</sup> on Thurs- day and Friday, •509.09 particles/ m <sup>3</sup> on weekend. School: •1004.55 parti- cles/m <sup>3</sup> on Mon- day and Tuesday, •590.09 particles/ m <sup>3</sup> on Thursday and Friday •227.27 particles/ m <sup>3</sup> on weekend. Apartment: •595.45 particles/ m <sup>3</sup> on Monday and Tuesday, •231.82 particles/ m <sup>3</sup> on Thursday and Friday, •236.36 particles/ m <sup>3</sup> on weekend	2020	(61)

CONTINUE

**Table 1: Occurrence of Airborne Microplastic in Indonesia 2019 – 2024 (CONT.)**

Study Location	Shape	Size	Colour	Polymer	TSP Concentration	Deposition Rate	Concentration	Year	Reference
Medan	N/A	N/A	N/A	N/A	PM <sub>2.5</sub> concentrations in Medan range from 25.78 – 60.36 µg/m <sup>3</sup> .	N/A	The amount of microplastics at 6 locations: 1.78 2.41 3.41 4.39 5.27 6.97 particles Abundance of microplastics at six locations: 1.1.18 2.0.62 3.0.6 4.0.57 5.0.41 6.1.4 particles/m <sup>3</sup>	2024	(62)
Bandar Lampung	Fragment<fibres	•500-1000 µm (42%), •100-500 µm (29%), •1000-1500 µm (15%), •1500-2000 µm (6%), •2000-2500 µm (4%), and •>2500 µm (4%)	•Blue, •Grey, •Black, •Brown, •Red, •Green, •Purple, •Pink, •Light Blue, •Yellow, •Orange, and •Transparent	Polyethylene Terephthalate (PET)	PM <sub>2.5</sub> : 25.78 – 60.36 µg/m <sup>3</sup>	N/A	Abundance of microplastics 0.0021-0.0199 particles/m <sup>3</sup> Average 0.008 ± 0.005 particles/m <sup>3</sup> of air or 49.812 ± 31.435 particles/day.	2024	(63)
Sidoarjo	Fragment<film<fibres	N/A	N/A	N/A	PM <sub>2.5</sub> : 1. 29 µg/m <sup>3</sup> 2.1876 µg/m <sup>3</sup>	N/A	Amount of microplastics 1.1243 particles 2. 15 particles	2021	(64)
Sidoarjo, Gresik, Mojokerto	Fragment, fibres, film	N/A	N/A	N/A	PM <sub>2.5</sub> : 1. 44 µg/m <sup>3</sup> 2. 821 µg/m <sup>3</sup> 3. 36 µg/m <sup>3</sup> 4. 50 µg/m <sup>3</sup> 5. 58 µg/m <sup>3</sup> 6. 40 µg/m <sup>3</sup>	N/A	Amount of microplastics at 6 locations: 1. 10 particles 2. 4 particles 3. 3 particles 4. 3 particles 5. 9 particles 6. 4 particles	2021	(65)
Sidoarjo	Fibres<fragment<film	N/A	N/A	N/A	PM <sub>2.5</sub> : 1. 96 µg/m <sup>3</sup> 2.224 µg/m <sup>3</sup> 3.153 4.302	N/A	Amount of microplastics at 4 locations: 911 particles	2023	(66)
Sidoarjo	Fragment<film<fibres	N/A	N/A	N/A	N/A	N/A	Location 1.3.7 particle/gr 2.7.8 particle/gr 3.8.9 particle/gr 4.17.2 particle/gr	2023	(67)




Figure 3: Illustration Passive (A) and Active (B) Sampling Method

using various sampling techniques. Furthermore, the concentration levels of microplastics and other empirical variables may be influenced by these different methods. Active sampling techniques necessitate the use of pumping sampler devices, which draw a fixed volume of air over a designated area. As air passes through the sampler, particulate matter is collected on a substrate or filter. Passive sampling methods are considered to be the most effective for atmospheric microplastic deposition due to their simplicity, affordability, ease of use, and reliance on common laboratory equipment. Additionally, since passive sampling techniques do not require electricity or other power sources, they can be employed for extended outdoor sampling that lasts for weeks or even months (75).

### Identification and Quantification of Microplastics

Microscopes enabled for the identification of microplastic shapes (fiber, foam, granule, and fragment), sizes of microplastics (<300 µm, 300–500 µm, 500–1000 µm, and >1000 µm), and colors (transparent, crystalline, white, clear-white-cream, red, orange, blue, opaque, black, gray, brown, green, pink, tan, yellow, and pigmented) (33,78).

Microplastics primarily exist in three forms: pellets, fragments, and other generic types. Pellets typically exhibit shapes such as cylinders, disks, flats, ovoids, and spheres. In contrast, fragments are most commonly found in angular, subrounded, rounded, and subangular shapes. Generally, microplastics are characterized by their elongated, uneven, and rough surfaces, often featuring broken edges (33). The diameter of microplastics could be accurately measured using Image-J software (60–62,70,79). One effective method for assessing minute variations in microplastic size while minimizing background noise is through image analysis (80,81).




Figure 4: Schematic Representation of Occurrence Airborne Microplastic, Prevention, and Health Risk

## DISCUSSION

### Location Site

All cities mentioned are provincial capitals in Indonesia, except for Sidoarjo, Gresik, and Mojokerto. The largest population in Indonesia is located in West Java, particularly in the cities of Depok, Bogor, and Bandung. Surabaya, in East Java, has the second largest population. Meanwhile, Jakarta, as the country's capital, experiences congestion that is twice as high during the day due to its status as an economic and government center. Medan ranks fourth in terms population size in Indonesia. Additionally, microplastic levels in urban areas are significantly higher than in suburban areas (19,23), attributed to the greater density of buildings and a higher volume of activities involving plastics and textiles (20,76,77).

### Sampling

Due to variations in the form and density of particles, different types of microplastics in the air can be identified

For Fourier Transform Infrared (FTIR) analysis, particles that were extracted from the filter were selected at random (82,83). FTIR analysis with the Nicolet iS10 from Thermo Fisher Scientifics. FTIR analysis was conducted using the Nicolet iS10 from Thermo Fisher Scientific. By analyzing the spectrum of the functional groups generated, this process determined the composition of each sample. The generated spectra were compared with those found in the Thermo Fisher Scientifics Nicolet databases (64,68). The concentration and properties of microplastics, including their shape, physical diameter, and polymer type, were examined using Micro-Fourier Transform Infrared (µ-FTIR) spectroscopy (Spotlight 400/Spectrum 3 PerkinElmer). The detector operated within a wave number range of 4,000–695 cm<sup>-1</sup>. Initially, the infrared peak bands of C-H stretching (3,000–2,700 cm<sup>-1</sup>) and C=O stretching (1,740–1,710 cm<sup>-1</sup>) were identified to screen for microplastic candidates (84). Raman Spectroscopy was employed to characterize the polymer (70).

## Occurrence of Airborne Microplastics

### Shapes

The findings indicate that the most common shapes of microplastics are fibers and fragments. Fibers are particularly prevalent due to their high production and waste rates, with an annual production rate of 6.6% (85). Furthermore, they are typically generated from activities such as the garment industry, washing, laundry, and clothing (20,76,77,86), and are often associated with dust from PET, a primary polymer used in textile production (19,87,88). Fibrous microplastics are classified as secondary microplastics; they rarely disintegrate in the environment. The primary concern regarding fibrous microplastics is their potential to disperse over vast distances (41).

Both indoor and outdoor environments predominantly contain fibers, which are released from textiles, carpets, laundry, synthetic clothing during washing, and automobile abrasion (primarily from tire treads) (87,88). Sixty-seven percent of indoor fibers are composed of natural materials (21). The concentration of fibers indoors is higher than that outdoors, as people spend over 70% of their time inside, where conditions are influenced by airflows, ventilation, and room layout. Consequently, elevated concentrations of indoor microplastics are often a result of inadequate air renewal rates (89). Furthermore, outdoor fiber concentrations are significantly lower than indoor levels, likely due to the migration of outdoor microplastics into the buildings caused by wind and air movement (29). During the rainy season, fiber concentrations are higher than in the dry season, probably because rainfall enhances the deposition of atmospheric microplastics (20,21). Various factors, including the materials used in clothing, bedding, carpets, and furniture, the methods and frequency of cleaning, and the turbulence and filtration of airflow caused by human activity and ventilation systems, are proposed account for the geographic variance of indoor airborne microplastics (21,37,90). Although fiber particles found in surface dust and airborne particles belong to the same category, fibers in surface dust are generally larger than those in the air. This observation suggests that fibers larger than 20  $\mu\text{m}$  are less likely to remain airborne for extended periods, indicating that this size range is particularly significant (38).

Fibers are predominantly found in various locations around the world, including in Europe: France, the UK (London, Yorkshire, Nottingham), Portugal (Aveiro), Spain (Barcelona), Turkey (Sakarya); the US (California – indoor building); Australia: Sydney, Victoria; and Asia: China (Dongguan, Yantai, Shanghai, Beijing), Japan (Kusatsu), Vietnam (Ho Chi Minh, Da Nang), Nepal (Kathmandu), India (Nagpur), Iran (Bushehr and Shiraz);

as well as the Arctic regions, including the Swiss Alps and Bremen, Bavaria (19,20,37,38,42,43,45–48,21,23–25,32–34,36). Fragments are primarily found in Europe: France, Germany (Hamburg), the US: California – outdoor building), and Asia: Iran (Tehran), China (Shanghai, Wenzhou), and India (Chennai) (22,26,30,35,39,40,44). The most prevalent forms of non-fibrous microplastics are fragments and films. Potential sources of fragments include thicker plastic products that can be recycled and disposable plastic bags (33). These items can develop from the physical abrasion and mechanical crushing of larger polymers over time, particularly those commonly found in construction materials and billboards (42). Since plastics constitute the majority of the materials discussed above, frequent exposure to heat and sunlight causes them to deteriorate (91). The breakdown of polyethylene and polypropylene bags and wrappings is typically the primary source of microplastic films (92). Expanded polystyrene (PS) products have the potential to emit foam microplastics (33).

### Sizes

In most studies, the size of microplastics in Indonesia ranges from 300 – 1000  $\mu\text{m}$ . Fiber sizes reported in other studies range from 200 – 400  $\mu\text{m}$ , 400 – 600  $\mu\text{m}$ , and up to 750  $\mu\text{m}$ , with these sizes being predominant; fibers in larger size ranges are considered rare. Some microplastics are smaller than 20  $\mu\text{m}$ , with additional size categories including 20 to 25  $\mu\text{m}$  and 50 to 200  $\mu\text{m}$  (20,22,23,28,31,38,47). Fragments sizes are reported as less than 63  $\mu\text{m}$ , 63 to 300  $\mu\text{m}$ , and more than 300  $\mu\text{m}$ , variation variations 10 – to 1,000  $\mu\text{m}$  (22, 28). smaller than 100  $\mu\text{m}$  dominated airborne microplastics (39). The prevalence of smaller airborne microplastics suggests that size plays a crucial role in controlling the mobilization and transportation of these particles in the atmosphere. Generally, particles remain suspended in the air longer if it is smaller (93). Smaller size fractions of microplastics dominate the airborne microplastics samples; larger microplastics tend to settle rapidly due to gravity, whereas smaller microplastics remain in the atmosphere for longer. On air sample filters, larger plastic fibers are more noticeable than tiny pieces, fragments, which might be why fibers have frequently drawn greater attention (39).

As size increases, the relative quantity of fibers decreases. The majority of research on indoor airborne microplastics supports this trend (21,27,39). Hartmann et al. (94) reported that the dominant size range varies across studies, indicating that multiple factors impact the size distribution. Among these variables are the sampling location, sampling technique, identification method, and visual detection threshold. The primary determinant of plastic particles' destiny in the environment and their ability to penetrate biological organisms is their size (93).

## Colors

15 It has been reported that airborne microplastics are  
2 found in a wide range of colors. Transparent, white, and  
black are the most common colours among them (95).  
The varying sizes and colors of microplastics originate  
from a diverse array of industrial, automotive, and  
household sources (41). One possible explanation for  
2 the prevalence of lighter-colored microplastics is the  
widespread use of various disposable plastics, such as  
plastic bags, in both residential and commercial settings  
(42).

## Polymer

14 The most commonly found polymers were PET, PE, PS,  
and polyester. PET was the predominant polymer in  
most of the fibers, polyethylene was primarily found in  
fragments, and polystyrene was mainly used in foams  
(34). PET is a plastic polymer extensively used to  
produce textiles, fabrics, and synthetic fibers (22,34). PE,  
PS, polyester fibers were more prevalent in residences  
where carpet served as the primary flooring material (31).  
Polystyrene is frequently used in the fast-food industry  
as a packaging material and for thermal insulation (96).  
11 Polyethylene is the most widely produced type of plastic  
and is also extensively used as a packaging material (97).

## The Atmospheric Microplastic Deposition Rate and Microplastic Concentration

41 These results showed that the atmospheric microplastic  
deposition rate and microplastic concentration differed  
across several locations. Variations in the quantity  
of airborne microplastics between urban and rural  
areas were also linked to changes in population and  
land-use variables (21). The transport and dispersion  
20 of airborne microplastics can also be influenced by  
30 various meteorological factors, such as temperature,  
precipitation, wind speed, wind direction, and the  
vertical gradient in pollutant concentration, which can  
affect how microplastics behave and move through the  
atmosphere and surroundings (98).

The "urban heat island effect" significantly influences  
the distribution of air pollutants in urban areas. (93),  
which may also affect the dispersion of airborne  
microplastics. Compared to rural residents, urban  
inhabitants face a higher risk of inhaling microplastics  
due to the elevated concentration of these particles in  
metropolitan environments. Additionally, the amount  
21 of accumulated rainfall may be a crucial climatic factor  
to consider for future monitoring, as indicated by the  
temporal dynamics of microplastic abundances (32).

17 In indoor areas, microplastics can be released by  
various sources, including carpets, toys, clothes,  
cleaning methods, and furniture made of synthetic  
materials (39,90). Indoor microplastics may accumulate  
as a result of air conditioning systems (39). The  
placement of buildings, human activity, and population  
density similarly affect the concentration and size of  
36

microplastics. Conversely, outdoors, the concentration  
of microplastics is impacted by heavy traffic volume  
and the presence of industrial and workshop units (99).  
Because arid-windy conditions favor the atmospheric  
transport of particles, climate and wind patterns are  
expected to influence the concentrations of microplastics  
in the air. Greater air microplastic deposition would  
occur in regions downwind of a recognized source area  
(such as big cities) than in areas upwind (39).

Consequently, microplastics are likely being released  
into the environment via several sources, such as  
exposed plastic that degrades in open landfills, or  
plastic debris that wears down during garbage transfer  
and processing processes (23). These studies carried  
out at waste recycling plants, waste treatment industry  
incinerators, temporary landfills, and Waste to Energy  
Power Plants had caused air pollution, which is called a  
false solution to plastic management (71–74).

This finding is similar across various countries, as all  
citizens use plastic intentionally in a variety of activities.  
Moreover, microplastics likely travel through the air  
due to wind deflation. Compared to a remote, pristine  
environment, the predicted mobility in the prior research  
was lower (22), indicating that microplastics may have  
traveled up to 95 km. However, the differing settings  
probably experience distinct climatic conditions due  
to varying assumptions (size/aerodynamic equivalent  
diameter and density) made about the particles.  
Additionally, considering the small sample size, the  
results of the current and earlier studies should be  
evaluated cautiously, and further research is suggested  
to strengthen this body of data. Airborne microplastics  
should be considered alongside particulate matter (as  
coarse, fine, or ultra-fine particles), which are regarded  
as significant air pollutants, in addition to SO<sub>x</sub>, NO<sub>x</sub>, and  
CO<sub>x</sub> emissions. These have been closely monitored for  
some time and should therefore be carefully examined  
for their effects on human health. As a result, although it  
is difficult to pinpoint exactly how much microplastics a  
location should have, it is conceivable to state that the  
atmosphere does contain some of them, albeit at varying  
levels. (43).

## Health Risk

Microplastic exposure can increase the risk of non-  
communicable diseases such as asthma and cancer.  
In Indonesia, the prevalence of asthma and cancer in  
2023 was 1.8% and 1.2%, respectively (100). In recent  
years, human exposure to microplastics has grown  
into a serious problem. Chronic exposure to airborne  
particulates through skin contact, inhalation, and  
ingestion negatively affects one's health (89). The most  
significant ways that microplastics are encountered are  
through ingestion and inhalation (31,101). Fibres are  
more prevalent (90%) and smaller (5–250 μm) (21,25)  
and are found in a wide variety of manufactured and  
natural materials used in daily living, such as furniture

9 24 and textiles. The sorts of fibres that are most likely to infiltrate the human body can disrupt its physiological processes (21,25). They can enter the human lungs by inhalation (102), are durable (103), and 20 µm in length cannot be cleansed by the human lungs (104). Although the discovered fibres are allegedly too large to be breathed, exposure can nonetheless occur through ingestion, especially for young infants. When inhaled, microplastics build up in the human lungs and directly contact respiratory organs, potentially causing long-term discomfort.

5 5 17 A recent study has identified disorders associated with plastics and their derivatives (86). Particles lodged in the nasal and pulmonary mucus can be expelled through sneezing, coughing, blowing the nose, and spitting out or swallowing the mucus. One of the most significant aspects of atmospheric microplastics is their potential for inhalation and subsequent passage through the alveoli of the lungs (breathable particles). The inhalability of particles is influenced by their size and shape; it appears that only fibrous particles and those smaller than 5 µm can become lodged in the deep lung (85). While mucociliary clearance in the upper airways effectively removes larger inhalable particles, some particles manage to evade this process and reach the deep lung. Particularly with longer fibers, these particles tend to evade clearance (105) and demonstrate remarkable resilience in physiological fluids, likely continuing to accumulate with ongoing inhalation (103).

47 62 6 22 6 There is a significant chance that occupational workers who handle para-aromatic amides, polyesters, or nylon fibers may experience coughing, dyspnea, and a reduction in lung capacity. (86) Several employees at plastic processing plants have suffered from respiratory and health issues (such as wheezing, coughing, dyspnea, and occupational asthma), potentially caused by long-term exposure to microplastics. (105) These health considerations suggest that regularly consuming organic fibers can lead to inflammation by triggering localized biological responses. (85,106) Greim et al. (107) proposed that the interaction of cells with vitreous fibers or particles could serve as an approximate measure of toxicity. They discovered that this interaction can release cytotoxic substances and intracellular messengers, which may promote lung inflammation. The ongoing production of reactive oxygen species can subsequently result in secondary genotoxicity (85,107), which can cause oxidative DNA damage (102,108) and is also a factor inducing cancer (105). Previous studies identified the presence of synthetic fibers, including cellulosic, cotton, nylon, polyamide, and plastic fibers (like polyester) in human lungs (102,103). Notably, 97% of malignant lung specimens contained fibers (21), all polymeric particles in the microplastics found in human lung tissues taken during autopsies were less than 5.5 µm, and the fibers varied in size from 8.12 to 16.8 µm. Polyethylene and polypropylene were the most

frequently identified polymers. (109)

### Prevention

Indirect and direct methods are employed to reduce the presence of microplastics in the environment. Examples of indirect solutions include installing appropriate filters in both new and existing ventilation or air conditioning systems, as well as utilizing air purifiers in private homes that lack air conditioning or are used infrequently. (110). The direct method involves reducing the sources of microplastics. (111). Public education and awareness regarding microplastics are essential, as is collaboration in minimizing or eliminating plastic usage.

Moreover, to strengthen efforts in reducing the sources of microplastics, the government needs to establish policies on airborne microplastics. The framework involves implementing proper waste management, the 3R principles (reuse, reduce, and recycle), regulating quality standards both indoors and outdoors, and monitoring airborne microplastics alongside particulate matter from indoor and outdoor areas. Indoor area monitoring is more focused on high-activity times and occupants on weekdays than on weekends, while outdoor area monitoring primarily targets the streets. Industries can recycle plastics to minimize the generation of existing plastic waste. These strategies will succeed through collaboration between the government, industry players, and the community.

### CONCLUSION

It described the presence of airborne microplastics in our environment and daily lives, with common shapes being fibers and fragments of varying sizes. The predominant polymers identified were PET, PE, and polyester. The duration of data collection (measurement), measurement device, meteorological conditions, location (indoor or outdoor, suburban or urban), human activity, and surrounding goods impacted these research findings. Most research was conducted on Java Island. Several cities and districts, such as Surabaya, Bali, and Banjarmasin, have implemented bans on plastic bags in groceries. This policy could minimize plastic use. Moreover, it is important to formulate these policies nationally to address airborne microplastics, to set thresholds for the concentrations of microplastics both indoors and outdoors, and to monitor their presence. It requires commitment from the government, industry, and the community. The health effects of non-communicable disease relate to the size, shape, and polymer of microplastics, but exact exposures could not be determined. Further research on the link between airborne microplastics and health effects on at-risk populations should be conducted.

### ACKNOWLEDGEMENT

The authors are sincerely grateful to the Department

of Environmental Health, Faculty of Public Health, Universitas Airlangga for their assistance in writing this review.

## REFERENCES

1. Schmid C, Cozzarini L, Zambello E. Microplastic's story. *Mar Pollut Bull* [Internet]. 2021;162:111820. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0025326X20309383?via%3Dihub>
2. Karl W. Kenyon EK. Laysan Albatrosses swallow indigestible matter. *Ornithology*. 1969;339–43.
3. Ryan PG. Ingestion of Plastics by Marine Organisms. In: *Hazardous Chemicals Associated with Plastics in the Marine Environment* [Internet]. SpringerLink; 2016. p. 235–266. Available from: [https://link.springer.com/chapter/10.1007/698\\_2016\\_21](https://link.springer.com/chapter/10.1007/698_2016_21)
4. Thompson RC, Olson Y, Mitchell RP, Davis A, Rowland SJ, John AWG, et al. Lost at Sea: Where Is All the Plastic? *Science* (80-). 2004;304(5672):838.
5. Gall SC, Thompson RC. The impact of debris on marine life. *Mar Pollut Bull* [Internet]. 2015;92(1–2):170–9. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0025326X14008571>
6. Cauwenberghe L Van, Vanreusel A, Mees J, Janssen CR. Microplastic pollution in deep-sea sediments. *Environ Pollut* [Internet]. 2013;182:495–9. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0269749113004387>
7. Faure F, Corbaz M, Baecher H, De Alencastro LF. Pollution due to plastics and microplastics in lake Geneva and in the Mediterranean sea. *Arch des Sci*. 2012;65(1–2):157–64.
8. Amanda McCormick, Hoellein TJ, Mason SA, Schlupe J, Kelly JJ. Microplastic is an Abundant and Distinct Microbial Habitat in an Urban River. *Environ Sci Technol* [Internet]. 2014;48(20):11863–71. Available from: <https://pubs.acs.org/doi/10.1021/es503610r>
9. Moore CJ, Lattin GL, Zellers AF. Quantity and type of plastic debris flowing from two urban rivers to coastal waters and beaches of Southern California. *Rev Gestro Costeira Integr*. 2011;11(1):65–73.
10. Morrith D, Stefanoudis P V., Pearce D, Crimmen OA, Clark PF. Plastic in the Thames: A river runs through it. *Mar Pollut Bull* [Internet]. 2014;78(1–2):196–200. Available from: <http://dx.doi.org/10.1016/j.marpolbul.2013.10.035>
11. Jambeck Jenna R, Geyer Roland, Wilcox Chris, Siegler Theodore R, Perryman Miriam, Andrady Anthony, et al. Plastic waste inputs from land into the ocean. *Science* (80- ) [Internet]. 2015;347(6223):768–70. Available from: <http://www.scopus.com/inward/record.url?eid=2-s2.0-84954204572&partnerID=40&md5=28a97ef4a4fdee6db9ef2fe507a1a02a>
12. Rillig MC. Microplastic in Terrestrial Ecosystems and the Soil? *Environ Sci Technol*. 2012;46(12):6453–4.
13. Brandon JA, Jones W, Ohman MD. Multidecadal increase in plastic particles in coastal ocean sediments. *Sci Adv*. 2019;5(9):1–6.
14. Geyer R, Jambeck JR, Law KL. Production, use, and fate of all plastics ever made. *Sci Adv*. 2017;3(7):25–9.
15. Meijer LJJ, van Emmerik T, van der Ent R, Schmidt C, Lebreton L. More than 1000 rivers account for 80% of global riverine plastic emissions into the ocean. *Sci Adv*. 2021;7(18):1–13.
16. Allen S, Allen D, Moss K, Le Roux G, Phoenix VR, Sonke JE. Examination of the ocean as a source for atmospheric microplastics. *PLoS One*. 2020;15(5):1–14.
17. Allen S, Allen D, Baladima F, Phoenix VR, Thomas JL, Le Roux G, et al. Evidence of free tropospheric and long-range transport of microplastic at Pic du Midi Observatory. *Nat Commun*. 2021;12(1).
18. Haque F, Fan C. Fate and Impacts of Microplastics in the Environment: Hydrosphere, Pedosphere, and Atmosphere. *Environ - MDPI*. 2023;10(5).
19. Dris R, Gasperi J, Rocher V, Saad M, Renault N, Tassin B. Microplastic contamination in an urban area: A case study in Greater Paris. *Environ Chem*. 2015;12(5):592–9.
20. Dris R, Gasperi J, Saad M, Mirande C, Tassin B. Synthetic fibers in atmospheric fallout: A source of microplastics in the environment? *Mar Pollut Bull* [Internet]. 2016;104(1–2):290–3. Available from: <http://dx.doi.org/10.1016/j.marpolbul.2016.01.006>
21. Dris R, Gasperi J, Mirande C, Mandin C, Guerrouache M, Langlois V, et al. A first overview of textile fibers, including microplastics, in indoor and outdoor environments. *Environ Pollut*. 2017;221:453–8.
22. Allen S, Allen D, Phoenix VR, Le Roux G, Pilar Durántez Jiménez, Anaëlle Simonneau SB& DG. Atmospheric transport and deposition of microplastics in a remote mountain catchment. *Nat Geosci* [Internet]. 2019;12:339–344. Available from: <https://www.nature.com/articles/s41561-019-0335-5>
23. Wright SL, Ulke J, Font A, Chan KLA, Kelly FJ. Atmospheric microplastic deposition in an urban environment and an evaluation of transport. *Environ Int* [Internet]. 2020;136(December 2019):105411. Available from: <https://doi.org/10.1016/j.envint.2019.105411>
24. Stanton T, Johnson M, Nathanail P, MacNaughtan W, Gomes RL. Freshwater and airborne textile fibre populations are dominated by 'natural', not microplastic, fibres. *Sci Total Environ* [Internet]. 2019;666:377–89. Available from: <https://doi.org/10.1016/j.scitotenv.2019.02.278>
25. Jenner LC, Sadofsky LR, Danopoulos E, Rotchell JM. Household indoor microplastics within the

- Humber region (United Kingdom): Quantification and chemical characterisation of particles present. *Atmos Environ* [Internet]. 2021;259(June):118512. Available from: <https://doi.org/10.1016/j.atmosenv.2021.118512>
26. Klein M, Fischer EK. Microplastic abundance in atmospheric deposition within the Metropolitan area of Hamburg, Germany. *Sci Total Environ* [Internet]. 2019;685:96–103. Available from: <https://doi.org/10.1016/j.scitotenv.2019.05.405>
27. Vianello A, Jensen RL, Liu L, Vollertsen J. Simulating human exposure to indoor airborne microplastics using a Breathing Thermal Manikin. *Sci Rep* [Internet]. 2019;9(1):1–11. Available from: <http://dx.doi.org/10.1038/s41598-019-45054-w>
28. Torres-Agullo A, Karanasiou A, Moreno T, Lacorte S. Airborne microplastic particle concentrations and characterization in indoor urban microenvironments. *Environ Pollut* [Internet]. 2022;308(March):119707. Available from: <https://doi.org/10.1016/j.envpol.2022.119707>
29. Prata JC, Castro JL, da Costa JP, Duarte AC, Rocha-Santos T, Cerqueira M. The importance of contamination control in airborne fibers and microplastic sampling: Experiences from indoor and outdoor air sampling in Aveiro, Portugal. *Mar Pollut Bull* [Internet]. 2020;159(July):111522. Available from: <https://doi.org/10.1016/j.marpolbul.2020.111522>
30. Gaston E, Woo M, Steele C, Sukumaran S, Anderson S. Microplastics Differ Between Indoor and Outdoor Air Masses: Insights from Multiple Microscopy Methodologies. *Appl Spectrosc*. 2020;74(9):1079–98.
31. Soltani NS, Taylor MP, Wilson SP. Quantification and exposure assessment of microplastics in Australian indoor house dust. *Environ Pollut*. 2021;283.
32. Su L, Nan B, Craig NJ, Pettigrove V. Temporal and spatial variations of microplastics in roadside dust from rural and urban Victoria, Australia: Implications for diffuse pollution. *Chemosphere* [Internet]. 2020;252:126567. Available from: <https://doi.org/10.1016/j.chemosphere.2020.126567>
33. Cai L, Wang J, Peng J, Tan Z, Zhan Z, Tan X, et al. Characteristic of microplastics in the atmospheric fallout from Dongguan city, China: preliminary research and first evidence. *Environ Sci Pollut Res*. 2017;24(32):24928–35.
34. Zhou Q, Tian C, Luo Y. Various forms and deposition fluxes of microplastics identified in the coastal urban atmosphere. *Chinese Sci Bull* [Internet]. 2017;62(33):3902–9. Available from: <https://www.sciengine.com/CSB/doi/10.1360/N972017-00956;JSESSIONID=51c89a88-4a09-4a97-bab4-f47e39ac6e7c>
35. Liu K, Wang X, Wei N, Song Z, Li D. Accurate quantification and transport estimation of suspended atmospheric microplastics in megacities: Implications for human health. *Environ Int* [Internet]. 2019;132(July):105127. Available from: <https://doi.org/10.1016/j.envint.2019.105127>
36. Liu K, Wang X, Fang T, Xu P, Zhu L, Li D. Source and potential risk assessment of suspended atmospheric microplastics in Shanghai. *Sci Total Environ* [Internet]. 2019;675:462–71. Available from: <https://doi.org/10.1016/j.scitotenv.2019.04.110>
37. Zhang Q, Zhao Y, Du F, Cai H, Wang G, Shi H. Microplastic Fallout in Different Indoor Environments. *Environ Sci Technol*. 2020;54(11):6530–9.
38. Li Y, Shao L, Wang W, Zhang M, Feng X, Li W, et al. Airborne fiber particles: Types, size and concentration observed in Beijing. *Sci Total Environ* [Internet]. 2020;705:135967. Available from: <https://doi.org/10.1016/j.scitotenv.2019.135967>
39. Liao Z, Ji X, Ma Y, Lv B, Huang W, Zhu X, et al. Airborne microplastics in indoor and outdoor environments of a coastal city in Eastern China. *J Hazard Mater* [Internet]. 2021;417(May):126007. Available from: <https://doi.org/10.1016/j.jhazmat.2021.126007>
40. Dehghani S, Moore F, Akhbarizadeh R. Microplastic pollution in deposited urban dust, Tehran metropolis, Iran. *Environ Sci Pollut Res*. 2017;24(25):20360–71.
41. Abbasi S, Keshavarzi B, Moore F, Turner A, Kelly FJ, Dominguez AO, et al. Distribution and potential health impacts of microplastics and microrubbers in air and street dusts from Asaluyeh County, Iran. *Environ Pollut* [Internet]. 2019;244:153–64. Available from: <https://doi.org/10.1016/j.envpol.2018.10.039>
42. Kashfi FS, Ramavandi B, Arfaeinia H, Mohammadi A, Saeedi R, De-la-Torre GE, et al. Occurrence and exposure assessment of microplastics in indoor dusts of buildings with different applications in Bushehr and Shiraz cities, Iran. *Sci Total Environ*. 2022;829.
43. Tunahan Kaya A, Yurtsever M, Ziftzi Bayraktar S. Ubiquitous exposure to microfiber pollution in the air. *Eur Phys J Plus*. 2018;133(11).
44. Patchaiyappan A, Dowarah K, Zaki Ahmed S, Prabakaran M, Jayakumar S, Thirunavukkarasu C, et al. Prevalence and characteristics of microplastics present in the street dust collected from Chennai metropolitan city, India. *Chemosphere* [Internet]. 2021;269:128757. Available from: <https://doi.org/10.1016/j.chemosphere.2020.128757>
45. Narmadha VV, Jose J, Patil S, Farooqui MO, Srimuruganandam B, Saravanadevi S, et al. Assessment of Microplastics in Roadside Suspended Dust from Urban and Rural Environment of Nagpur, India. *Int J Environ Res* [Internet]. 2020;14(6):629–40. Available from: <https://doi.org/10.1007/s41742-020-00283-0>
46. Yukioka S, Tanaka S, Nabetani Y, Suzuki Y, Ushijima T, Fujii S, et al. Occurrence and

- characteristics of microplastics in surface road dust in Kusatsu (Japan), Da Nang (Vietnam), and Kathmandu (Nepal). *Environ Pollut* [Internet]. 2020;256:113447. Available from: <https://doi.org/10.1016/j.envpol.2019.113447>
47. Bergmann M, Mützel S, Primpke S, Tekman MB, Trachsel J, Gerdts G. White and wonderful? Microplastics prevail in snow from the Alps to the Arctic. *Sci Adv*. 2019;5(8):1–10.
48. Thinh TQ, Sang TTN, Viet TQ, Tam LTM, Dan NP, Strady E, et al. Preliminary assessment on the microplastic contamination in the atmospheric fallout in the Phuoc Hiep landfill, Cu Chi, Ho Chi Minh city. *Vietnam J Sci Technol Eng*. 2020;62(3):83–9.
49. Dewika M, Markandan K, Nagaratnam S, Irfan NA, Abdah MAAM, Ruwaida JN, et al. Assessing the concentration, distribution and characteristics of suspended microplastics in the Malaysian indoor environment. *Sci Total Environ* [Internet]. 2025;959(September 2024):178049. Available from: <https://doi.org/10.1016/j.scitotenv.2024.178049>
50. Haida N, Mohmad A, Norruwaida J, Dewika M, Sara YY, Norfatih I, et al. Environmental Influences on Microplastics and Particulate Matter Resuspension in Indoor Air : A Study of Office Settings with Air Conditioning. 2024;8(March):44–9.
51. Hashim HA, Jamian NR, Yusuf SY, Naidu D, Paad KM, Ishak NF. Evaluation of Deposited Atmospheric Microplastic Characteristic within Malaysia Cities Airshed. *Chem Eng Trans*. 2023;106(July):1105–10.
52. Chenappan NK, Ibrahim YS, Anuar ST, Yusof KMKK, Jaafar M, Ahamad F, et al. Quantification and characterization of airborne microplastics in the coastal area of Terengganu, Malaysia. *Environ Monit Assess* [Internet]. 2024;196(3). Available from: <https://doi.org/10.1007/s10661-024-12381-z>
53. Sarathana D, Winijkul E. Concentrations of Airborne Microplastics during the Dry Season at Five Locations in Bangkok Metropolitan Region, Thailand. *Atmosphere (Basel)*. 2023;14(1).
54. Limsiriwong K, Winijkul E. Exploring Personal Exposure to Airborne Microplastics across Various Work Environments in Pathum Thani Province, Thailand. *Int J Environ Res Public Health*. 2023;20(24).
55. Myat YN, Kongpran J, Vattanasit U, Tanaka S. Airborne microplastics in the roadside and residential areas of Southern Thailand. *Case Stud Chem Environ Eng* [Internet]. 2024;9(February):100682. Available from: <https://doi.org/10.1016/j.csee.2024.100682>
56. Winijkul E, Latt KZ, Limsiriwong K, Pussayanavin T, Prapasongsa T. Depositions of airborne microplastics during the wet and dry seasons in Pathum Thani, Thailand. *Atmos Pollut Res* [Internet]. 2024;15(10):102242. Available from: <https://doi.org/10.1016/j.apr.2024.102242>
57. Dahal Y, Babel S. Abundance and characteristics of atmospheric microplastics deposition in indoor and outdoor environments in Bangkok, Thailand. *Air Qual Atmos Heal*. 2024;
58. Parvez MS, Ullah H, Faruk O, Simon E, Czjdl H. Role of Microplastics in Global Warming and Climate Change: A Review [Internet]. Vol. 235, Water, Air, and Soil Pollution. Springer International Publishing; 2024. 1–31 p. Available from: <https://doi.org/10.1007/s11270-024-07003-w>
59. Purwiyanto AIS, Prartono T, Riani E, Naulita Y, Cordova MR, Koropitan AF. The deposition of atmospheric microplastics in Jakarta-Indonesia: The coastal urban area. *Mar Pollut Bull* [Internet]. 2022;174(November 2021):113195. Available from: <https://doi.org/10.1016/j.marpolbul.2021.113195>
60. Syafina PR, Yudison AP, Sembiring E, Irsyad M, Tomo HS. Identification of fibrous suspended atmospheric microplastics in Bandung Metropolitan Area, Indonesia. *Chemosphere* [Internet]. 2022;308(P2):136194. Available from: <https://doi.org/10.1016/j.chemosphere.2022.136194>
61. Yudison A. Investigating Atmospheric Microplastic in Total Suspended Particulate ( TSP ): Preliminary Study. 2020;(June).
62. Hidayat NAA, Kitano K, Tani Y, Lestari P, Iriana W, Fujii Y, et al. Airborne microplastics in Bandung and Osaka: concentration and characteristics. *E3S Web Conf*. 2024;485:1–12.
63. Jannah BR, Maharani HA, Rahmawati S, Nugroho AR, Abdull NB. Occurrence and characteristic of microplastics in suspended particulate, a case study in street of Yogyakarta. *E3S Web Conf*. 2024;485.
64. Syafei AD, Nurasrin NR, Assomadi AF, Boedisantoso R. Microplastic Pollution in the Ambient Air of Surabaya, Indonesia. *Curr World Environ*. 2019;14(2):290–8.
65. Asrin N, Dipareza A. Microplastics in Ambient Air (Case Study : Urip Sumoharjo Street and Mayjend Sungkono Street of Surabaya City , Indonesia). *IAETSD J Adv Res Appl Sci*. 2019;6(54):54–7.
66. Luo Z, Zhou X, Su Y, Wang H, Yu R, Zhou S, et al. Environmental occurrence, fate, impact, and potential solution of tire microplastics: Similarities and differences with tire wear particles. *Sci Total Environ*. 2021;795.
67. Bahrina I, Syafei AD, Satoto R, Jiang JJ, Nurasrin NR, Assomadi AF, et al. An Occupant-Based Overview of Microplastics in Indoor Environments in the City of Surabaya, Indonesia. *J Ecol Eng*. 2020;21(8):236–42.
68. Pratiwi A, Syafei AD, Assomadi AF, Boedisantoso R, Hermana J. Microplastic characterization based on the number of occupants. *AIP Conf Proc*. 2020;2296(November).
69. Suryati I, Hasibuan NH, Silalahi RA, Malau R, Tambun Y. Microplastic Distribution Model in

- Ambient Air PM<sub>2.5</sub> Around the Medan Industrial Area, North Sumatra. *E3S Web Conf.* 2024;519:1–6.
70. Kartika Sari N, Daffa Shiddiq M. Atmospheric Microplastic Particulate in Urban Roadside: Case of Bandar Lampung City, Indonesia. 2024;1–16.
71. Fathulloh MZ, Minanurrohman MR, Mahmudah R. False Solution Plastic Managemen. *Environ Pollut J.* 2021;1(3):208–16.
72. Syarifah N, Sari BSE, Hanapi A. Identifikasi Mikroplastik Di Udara Pada False Solution Technology. *Environ Pollut J.* 2022;1(3):199–207.
73. Titan Memory Yuhana. Identifikasi Mikroplastik pada Tahu di Sentra Industri Tahu Desa Tropodo Kecamatan Krian Kabupaten Sidoarjo. *Environ Pollut J.* 2023;3(3):770–85.
74. Yuhana TM. Identifikasi Mikroplastik pada Tahu di Sentra Industri Tahu Desa Tropodo Kecamatan Krian Kabupaten Sidoarjo. *Environ Pollut J.* 2023;3(3).
75. Shao L, Li Y, Jones T, Santosh M, Liu P, Zhang M, et al. Airborne microplastics: A review of current perspectives and environmental implications. *J Clean Prod.* 2022;347(March).
76. Atis S, Tutluoglu B, Levent E, Ozturk C, Tunaci A, Sahin K, et al. The respiratory effects of occupational polypropylene flock exposure. *Eur Respir J.* 2005;25(1):110–7.
77. Liebezeit G, Liebezeit E. Origin of synthetic particles in honeys. *Polish J Food Nutr Sci.* 2015;65(2):143–7.
78. Akhbarizadeh R, Dobaradaran S, Amouei Torkmahalleh M, Saeedi R, Aibaghi R, Faraji Ghasemi F. Suspended fine particulate matter (PM<sub>2.5</sub>), microplastics (MPs), and polycyclic aromatic hydrocarbons (PAHs) in air: Their possible relationships and health implications. *Environ Res [Internet].* 2021;192(September 2020):110339. Available from: <https://doi.org/10.1016/j.envres.2020.110339>
79. Tokunaga Y, Okochi H, Tani Y, Niida Y, Tachibana T, Saigawa K, et al. Airborne microplastics detected in the lungs of wild birds in Japan. *Chemosphere [Internet].* 2023;321(November 2022):138032. Available from: <https://doi.org/10.1016/j.chemosphere.2023.138032>
80. Cowger W, Gray A, Christiansen SH, DeFrono H, Deshpande AD, Hemabessiere L, et al. Critical Review of Processing and Classification Techniques for Images and Spectra in Microplastic Research. *Appl Spectrosc.* 2020;74(9):989–1010.
81. Valente T, Ventura D, Matiddi M, Sbrana A, Silvestri C, Piermarini R, et al. Image processing tools in the study of environmental contamination by microplastics: reliability and perspectives. *Environ Sci Pollut Res [Internet].* 2023;30(1):298–309. Available from: <https://doi.org/10.1007/s11356-022-22128-3>
82. Qiu Q, Peng J, Yu X, Chen F, Wang J, Dong F. Occurrence of microplastics in the coastal marine environment: First observation on sediment of China. *Mar Pollut Bull [Internet].* 2015;98(1–2):274–80. Available from: <http://dx.doi.org/10.1016/j.marpolbul.2015.07.028>
83. Vianello A, Boldrin A, Guerriero P, Moschino V, Rella R, Sturaro A, et al. Microplastic particles in sediments of Lagoon of Venice, Italy: First observations on occurrence, spatial patterns and identification. *Estuar Coast Shelf Sci [Internet].* 2013;130:54–61. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0272771413001480>
84. Wang Y, Okochi H, Tani Y, Hayami H, Minami Y, Katsumi N, et al. Airborne hydrophilic microplastics in cloud water at high altitudes and their role in cloud formation. *Environ Chem Lett [Internet].* 2023;21(6):3055–62. Available from: <https://doi.org/10.1007/s10311-023-01626-x>
85. Gasperi J, Wright SL, Dris R, Collard F, Mandin C, Guerrouache M, et al. Microplastics in air: Are we breathing it in? *Curr Opin Environ Sci Heal [Internet].* 2018;1:1–5. Available from: <https://doi.org/10.1016/j.coesh.2017.10.002>
86. Prata JC. Airborne microplastics: Consequences to human health? *Environ Pollut [Internet].* 2018;234:115–26. Available from: <https://doi.org/10.1016/j.envpol.2017.11.043>
87. Alasdair Carmichael. Man-Made Fibers Continue To Grow [Internet]. 2015. Available from: <https://www.textileworld.com/textile-world/fiber-world/2015/02/man-made-fibers-continue-to-grow/>
88. Liu C, Li J, Zhang Y, Wang L, Deng J, Gao Y, et al. Widespread distribution of PET and PC microplastics in dust in urban China and their estimated human exposure. *Environ Int.* 2019;128(April):116–24.
89. Wright SL, Kelly FJ. Plastic and Human Health: A Micro Issue? *Environ Sci Technol.* 2017;51(12):6634–47.
90. Zhang Y, Kang S, Allen S, Allen D, Gao T, Sillanpää M. Atmospheric microplastics: A review on current status and perspectives. *Earth-Science Rev [Internet].* 2020;203(September 2019):103118. Available from: <https://doi.org/10.1016/j.earscirev.2020.103118>
91. Cai L, Wang J, Peng J, Wu Z, Tan X. Observation of the degradation of three types of plastic pellets exposed to UV irradiation in three different environments. *Sci Total Environ [Internet].* 2018;628–629:740–7. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0048969718304698?via%3Dihub>
92. Nor NHM, Jeffrey Philip Obbard. Microplastics in Singapore's coastal mangrove ecosystems. *Mar Pollut Bull [Internet].* 2014;79(1–2):278–83. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0025326X13007261>
93. Veron F. Ocean spray. *Annu Rev Fluid Mech.*

- 2015;47:507–38.
94. Hartmann NB, Hüffer T, Thompson RC, Hasselluv M, Verschoor A, Daugaard AE, et al. Are We Speaking the Same Language? Recommendations for a Definition and Categorization Framework for Plastic Debris. *Environ Sci Technol*. 2019;53(3):1039–47.
  95. Wang Y, Huang J, Zhu F, Zhou S. Airborne Microplastics: A Review on the Occurrence, Migration and Risks to Humans. *Bull Environ Contam Toxicol* [Internet]. 2021;107(4):657–64. Available from: <https://doi.org/10.1007/s00128-021-03180-0>
  96. Song XC, Lin QB, Zhang YC, Li Z, Zeng Y, Chen ZF. Rapid classification of virgin and recycled EPS containers by Fourier transform infrared spectroscopy and chemometrics. *Altmetric Artic Rapid Classif virgin Recycl EPS Contain by Fourier Transform infrared Spectrosc Chemom* [Internet]. 2018;35(11):2220–9. Available from: <https://www.tandfonline.com/doi/full/10.1080/19440049.2018.1515502>
  97. Andrady AL. The plastic in microplastics: A review. *Mar Pollut Bull* [Internet]. 2017;119(1):12–22. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0025326X1730111X?via%3Dihub>
  98. Kaur S, Nieuwenhuijsen MJ, Colvile RN. Fine particulate matter and carbon monoxide exposure concentrations in urban street transport microenvironments. *Atmos Environ* [Internet]. 2007;41(23):4781–810. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S1352231007001343?via%3Dihub>
  99. Nematollahi MJ, Zarei F, Keshavarzi B, Zarei M, Moore F, Busquets R, et al. Microplastic occurrence in settled indoor dust in schools. *Sci Total Environ* [Internet]. 2022;807(3):150984. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0048969721060629>
  100. BPS. Dalam Angka Dalam Angka. Kota Kediri Dalam Angka. 2018;1–68.
  101. Zhang J, Wang L, Kannan K. Microplastics in house dust from 12 countries and associated human exposure. *Environ Int* [Internet]. 2020;134(June 2019):105314. Available from: <https://doi.org/10.1016/j.envint.2019.105314>
  102. Lippmann M. Effects of fiber characteristics on lung deposition, retention, and disease. *Environ Health Perspect*. 1990;88:311–7.
  103. Law BD, Bunn W. B, Hesterberg TW. Solubility of Polymeric Organic Fibers and Manmade Vitreous Fibers in Gambles Solution. *Inhal Toxicol* [Internet]. 1990;2(4). Available from: <https://www.tandfonline.com/doi/epdf/10.3109/08958379009145261?needAccess=true>
  104. Swaen GGMH, Teggeler O, Van Amelsvoort LGPM. False positive outcomes and design characteristics in occupational cancer epidemiology studies. *Int J Epidemiol*. 2001;30(5):948–54.
  105. Warheit DB, Hart GA, Hesterberg TW, Collins JJ, Dyer WM, Swaen GM, et al. Potential Pulmonary Effects of Man-Made Organic Fiber (MMOF) Dusts. *Crit Rev Toxicol* [Internet]. 2001;31(6). Available from: <https://www.tandfonline.com/doi/abs/10.1080/20014091111965>
  106. Pauly JL, Stegmeier SJ, Allaart HA, Cheney RT, Zhang PJ, Mayer AG, et al. Inhaled cellulosic and plastic fibers found in human lung tissue. *Cancer Epidemiol Biomarkers Prev*. 1998;7(5):419–28.
  107. Greim H, Borm P, Schins R, Donaldson K, Driscoll K, Hartwig A, et al. Toxicity of Fibers and Particles? Report of The Workshop Held In Munich, Germany, 26-27 October 2000. *Inhal Toxicol* [Internet]. 2008;13(9). Available from: <https://www.tandfonline.com/doi/epdf/10.1080/08958370118273?needAccess=true>
  108. WHO. Air quality guidelines for Europe, 2nd edition [Internet]. 2000. Available from: <https://www.who.int/publications/i/item/9789289013581>
  109. Amato-Lourenço LF, Carvalho-Oliveira R, Júnior GR, dos Santos Galvao L, Ando RA, Mauad T. Presence of airborne microplastics in human lung tissue. *J Hazard Mater*. 2021;416(April).
  110. Kacprzak S, Tijning LD. Microplastics in indoor environment: Sources, mitigation and fate. *J Environ Chem Eng* [Internet]. 2022;10(2):107359. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S2213343722002329>
  111. Patil S, Bafana A, Naoghare PK, Krishnamurthi K, Sivanesan S. Environmental prevalence, fate, impacts, and mitigation of microplastics—a critical review on present understanding and future research scope. *Environ Sci Pollut Res*. 2021;28(5):4951–74.