

BAB IV

KESIMPULAN DAN SARAN

4.1. Kesimpulan

Jenis produk teh hitam umumnya ada 2 yaitu *tea bag* dan *loose tea*. Kedua jenis produk tersebut memiliki aktivitas antioksidan dan kandungan polifenol yang berbeda. Aktivitas antioksidan dan kandungan polifenol juga dipengaruhi oleh suhu dan waktu penyeduhan. Berdasarkan studi literatur yang dilakukan, diketahui bahwa jenis produk teh hitam *tea bag* dan *loose tea* dari berbagai merk memiliki aktivitas antioksidan dan kandungan polifenol yang berbeda – beda. Pada kondisi penyeduhan yang sama, jenis *tea bag*, memiliki aktivitas antioksidan dan kandungan polifenol yang lebih tinggi dibandingkan dengan *loose tea*. Suhu dan waktu penyeduhan pada kedua jenis produk teh hitam tersebut berpengaruh terhadap aktivitas antioksidan dan kandungan polifenolnya. Semakin tinggi suhu (30°C – 90°C) dan semakin lama waktu penyeduhan (3 – 10 menit), menghasilkan aktivitas antioksidan dan kandungan polifenol yang semakin tinggi. Suhu yang tinggi, menyebabkan proses ekstraksi teh berlangsung lebih cepat.

4.2. Saran

Suhu dan waktu penyeduhan teh hitam yang digunakan sabaiknya berkisar antara 90 – 100°C dengan waktu 3 menit, dan pada suhu 80°C dengan waktu 5 menit untuk *tea bag* dan 60 menit dengan suhu 80°C untuk *loose tea*, agar aktivitas antioksidan dan kandungan polifenol dalam teh hitam dapat memberikan manfaat kesehatan yang lebih maksimal bagi tubuh.

DAFTAR PUSTAKA

- Anonim. 2018. Black Tea (Loose). <https://www.philzcoffee.com/teas-black> (Diakses tanggal 27 Mei 2018)
- Anonim. 2007. *Superoxide Dismutase Kit*. Reagent kit for the analysis of Superoxide Dismutase incell extracts.
- Apak, R., S. Gorinstein, V. Bohm, K. M. Schaich, M. Ozyurec, and K. Guclu. 2013. Methods of Measurement and Evaluation of Natural Antioxidant Capacity / Activity (IUPAC Technical Report). Istanbul: *Departement of Chemistry, Istanbul University*
- Asosiasi Industri Minuman Ringan. 2014. <https://asrim-indonesia.org/> (Diakses tanggal 29 Juli 2018)
- Astil, C., M. R. Birch, C. Dacombe, P. G. Humprey, and P.T. Martin. 2001. Factors Affecting The Caffeine and Polyphenol Content of Black and Green Tea Infusion. *Journal of Agricultural Food Chemistry* 49: 5340 - 5347.
- Balentine, D.A., S.A. Wiseman, and L.C.M. Bouwens. 1997. The Chemistry of Tea Flavonoids. *Critical Review Food Science Nutrition.*, 37, 693–704
- Blocknina, O., E. Virolainen, and K.V. Fagerstedt. 2003. Antioxidants Oxidative Damage and Oxygen Deprivation Stress. *Review* 91 : 179-194.
- Celestino, S.B., and S. Augustin. 2000. Proanthocyanidins and Tannin Like Compounds Nature Occurrence, Dietary Intake and Effects on Nutrition and Health. *Journal of Food Agriculture* 80:1094-1117.
- Chu, D.C., and L. R. Juneja. 1997. General Chemical Composition of Green Tea and its Infusion. in: Chemistry and Applications of Green Tea (eds. T. Yamamoto, L.R. Juneja, D.C. Chu, M. Kim). CRC, New York, pp. 13–22.
- Cook, N. C. and S. Samman, 1996, Flavonoids : Chemistry, Metabolism, Carsioprotective Effects, and Dietary Sources, *Nutrition Biochemistry* 7: 66-67.

- Di Castelnuovo, A., R. Digiuseppe, L. Iacoviello, and G. Degaetano. 2011. Consumption of Cocoa, Tea, and Coffe and Risk of Cardiovascular Disease. *Internal Medicine*: 1-11.
- Fernandez P.L., F. Pablos, M. J. Martin, and A. G. Gonzales. 2002. Study of Catechin and Xantine Tea Profiles as Geographical Tracers. *Journal of Agricultural and Food Chemistry* 50: 1833-1839.
- Folin, Octo, Ciocalteu, and Vintila, 1944. On Tyrosine and Tryptophane Determinations in Proteins, *Journal Biochemistry.*, 73 : 627-650, 1927
- Gupta, S., B. Saha, and A.K. Giri. 2002. Comparative Antimutagenic and Anticlastogenic Effects of Green Tea and Black Tea: A Review. *Mutation Research* 512:37-65.
- Hahn, D.H., L. W. Rooney, and C.F. Earp. 1984. Tannins and Phenols of Sorghum. *Cereal Foods World* 29: 776-779.
- Han, X., T. Shen, and L. Hongxiang. 2007. Dietary Polyphenols and Their Biological Significance. *International Journal of Molecular Science* 8: 950-988.
- Khokhar, S., And S.G.M. Magnusdottir. 2002. Total Phenol, Catechin, and Caffeine Contents of Teas Commonly Consumed in The United Kingdom. *Journal of Agricultural and Food Chemistry* 50: 565-570.
- Kunarto, B. 2005. *Teknologi Pengolahan Teh Hitam (Camellia sinensis L. Kuntze) Sistem Orthodox*. Semarang: University Press Semarang
- Lagiou P, J. Wuu, and A. Trichopoulou. 2004. Diet and Benign Prostatic Hyperplasia: a Study in Greece. *Urology* 54: 424-428.
- Litescu, S. C., S. A.V. Eremia, A. Tache, I. Vasilescu, dan G. L. Radu. 2014. *The Use of Oxygen Radical Absorbance Capacity (ORAC) and Trolox Equivalent Antioxidant Capacity (TEAC) Assays in the Assessment of Beverages' Antioxidant Properties*. Bucharest: National Institute for Biological Sciences.
- Madhavi, D. L., S. S. Deshpande, and D. K. Salunkhe. 1996. *Technological Aspects of Food Antioxidants, Toxicological and Health Perspectives*. Inc, New York. pp 159-265.

- Mermelstein, N.H. 2007. Determining Antioxidant Activity. *Food Technology*. p. 63-66
- Miyachi, Y. 1995. Photoaging from an oxidative standpoint. *Journal of Dermatology Science* 9:79-86.
- Nikniaz, Z., R. Mahdavi, S. J. Ghaemmaghami, N. L. Yagin, and L. Nikniaz. 2015. Effect of Different Brewing Times on Antioxidant Activity and Polyphenol Content of Loosely Packed and Bagged Black Teas (*Camellia Sinensis L.*). *Avicenna Journal of Phytomedicine (AJP)* 6(3): 313 – 321.
- Nazaruddin dan Paimin. 1993. *Pembudidayaan dan Pengolahan Teh*. Jakarta: Penerbit Penebar Swadaya.
- Polydera, A. C., N. G. Stoforos, and P. S. Taoukis. 2005. Effect of High Hydrostatic Pressure Treatment on Post Processing Antioxidant Activity of Fresh Navel Orange Juice. *Journal of Agricultural and Food Chemistry* 91:495-503.
- Pratt, D. E. 1992. Natural Antioxidants from Plant material. In Phenolic Compounds in Food and their Effects on Health (Vol. II). Antioxidants and Cancer Prevention; Huang, M-T., Ho, C-T., Lee, C., Eds.: ACS Symposium Series 507; American Chemical Society: Washington, DC.
- Rosyadi, A. I. 2001. *Efisiensi Penggunaan Sumber Daya Untuk Memproduksi Teh Hitam Berkelanjutan*. Bandung: Universitas Padjajaran
- Roginsky, V., Tatyana B., Chyong F. H., and Paul A. K. 2003. Chain-Breaking Antioxidant Activity and Cyclic Voltammetry Characterization of Polyphenols in a Range of Green, Oolong, and Black Teas. *Journal of Agricultural and Food Chemistry* 51(19) : 1864 – 1873
- Shahidi, F., and P. D. Wanasundara. 1992. Phenolic antioxidants. *Critical Review in Food Science and Nutrition* 1: 67-103.
- Shahidi, F. and M. Naczk. 1995. *Food Phenolics: Sources, Chemistry, Effects and Applications*. Technomic Publication Company, Inc., Lancaster.
- Shalaby and Sanaa. 2013. Antioxidant Compounds, Assays of Determination and Mode of Action. Africa: *Academic Journals*

- Shonisani, N. 2010. *Effects of Brewing Temperature and Duration on Quality of Black Tea (*Camellia Sinensis*) and Equal (50:50) Combination of Bush Tea (*Athrixia Phylicoides DC.*) and Black Tea*. University of Limpopo.
- Singleton, V.L. and J. A. Rossi, 1965. Colorimetry of Total Phenolic with Phosphomolybdic-Phosphotungstic Acid Reagent. *American Journal of Enology and Viticulture* 16: 147.
- Su, X.G., J. Duan, Y. M. Jiang, J. Shi, and Y. Kakuda. 2006. Effects of Soaking Conditions on the Antioxidant Potentials of Oolong Tea. *Journal of Food Composition* 19:348-353.
- Su, X., J. Duan, J. Y. Jiang, D. Xuewa, and F. Chen. 2007. Polyphenolic Profile and Antioxidant Activities of Oolong Tea Infusion Under Various Steeping Conditions. *International Journal of Molecular Science* 8:1196 – 1205.
- Susanti, R. M. 2016. *Analisis Aktivitas Antioksidan Teh Hitam Celup Menggunakan Metode Superoksida Dismutase (SOD)*. Bandung: Universitas Pasundan.
- Swapna, L. A., K. Pradeep, P. Reddy, K. Deepak, and S. Goyal. 2014. Antioxidants and Their Implication in Oral Health and General Health. *International Journal of Case Reports* 5(4): 258 – 263.
- Total Merchandise. 2018. *Label Tea bags*.
<https://www.totalmerchandise.co.uk/products/details/Label-Tea-Bags>
 (Diakses tanggal 27 Mei 2018)
- Turkmen, N., Y. S. Velioglu, F. Sari, dan G. Pola. 2007. Effect of Extraction Conditions on Measured Polyphenol Contents, Antioxidant and Antibacterial Activities of Black Tea. *Molecules* 12: 484-496.
- Venditti, E., T. Bacchetti, L. Tiano, P. Carloni, L. Greci, E. Damiani. 2010. Hot vs. Cold Water Steeping of Different Teas: Do They Affect Antioksidant Activity?. *Food Chemistry* 119: 1597–1604
- Winarsi, H. 2007. *Antioksidan Alami dan Radikal Bebas*. Yogyakarta: Kanisius

- Zuo, Y., H. Chen, and Y. Deng. 2002. Simultaneous Determination of Catechins Caffeine and Gallic Acid in Green, Oolong, Black and Puerh Teas Using Hplc With Aphotodiode Array Detector. *Talanta* 57:3017-4012.