

# The Lipid Lowering Effect of Water Extract of *Andrographis paniculata*, Water Extract of *Syzygium polyanthum* and its Combination in Alloxan-Induced Diabetic Rats

*by* Sumi Wijaya

---

FILE	22PI-THE_LIPID_LOWERING.PDF (6.29M)	WORD COUNT	2645
TIME SUBMITTED	05-DEC-2019 02:34PM (UTC+0700)	CHARACTER COUNT	14499
SUBMISSION ID	1227573080		

The Lipid Lowering Effect of Water Extract of *Andrographis paniculata*, Water Extract of *Syzygium polyanthum* and its Combination in Alloxan-Induced Diabetic Rats

Sumi Wijaya\* and Farida L.Darsono\*

\*Faculty of Pharmacy, Widya Mandala Catholic University, Surabaya Indonesia

Correspondence should be addressed to Sumi Wijaya, sumiwijaya@yahoo.com

Background: Combination of water extract of *Andrographis paniculata* and water extract of *Syzygium polyanthum* (6:1) have been reported to reduce blood glucose in normal and alloxan rats. This combination also gave protection effects on kidney, liver and pancreas animals tested from alloxan toxicity

Methods : Eighty male white albino rats were divided into eight different groups including normal control, diabetic control, 4 treated groups with the water extract of *Andrographis paniculata*, water extract of *Syzygium polyanthum* and its combination, and also standard drugs (metformin, insulin) with 10 rats in each group. The extracts and the standard were given orally for 7 days. At day 7 blood sample was collected, serum separated and glycemic level, total cholesterol, LDL, HDL, triglyceride, and alkaline phosphatase were determined. The body weight was recorded on day 1 and day 7.

Results and discussion: The data obtained revealed that the water extract of *Andrographis paniculata* (WAP), water extract of *Syzygium polyanthum* (WSP) and its combination (1 WAP: 6 WSP; 6 WAP:1 WSP) reduced the glucose level. The treatment with these extracts also reduced the LDL and triglyceride level significantly and alkaline phosphatase level as compared with normal and diabetic groups.

Conclusions: The water extract of *Andrographis paniculata*, water extract of *Syzygium polyanthum* and its combination possess significant antidiabetic activity along with antihypercholesterolemic activity.

**Key words:** *Andrographis paniculata*, *Syzygium polyanthum*, water extracts, antidiabetic, antihypercholesterolemic

### 1. Introduction

Diabetes mellitus is one of the common metabolic disorders with micro and macrovascular complications that results in significant morbidity and mortality. It is considered as one of the five leading causes of death in the world<sup>1-2</sup>. Hence, the global number of people with diabetes is estimated to rise up to 266 millions in 2030<sup>3-4</sup>.

In modern medicine no satisfactory effective therapy is still available to cure diabetes mellitus<sup>5</sup>. There is increasing demand by patients to use natural products with antidiabetic activity due to side effects associated with the use of insulin and oral hypoglycemic agents<sup>6-8</sup>.

World is endowed with a rich wealth of medicinal plants. These plants have made a good contribution to the development of ancient materia medica. More than 400 medicinal plants are present worldwide for the treatment of diabetes mellitus, while only few of them have

been subjected to scientific authentication as anti-diabetic agents<sup>9-10</sup>.

*Andrographis paniculata* (Acanthaceae), commonly called as king of bitter is a traditional

medicinal plant common in South East Asia and found from India to Indo-China. According to the study conducted by Widjajakusuma et al.<sup>11</sup> pretreatment with *Andrographis paniculata* aqueous extract with dose 200 mg/Kg BW demonstrated significant antidiabetic activity in alloxan induced diabetic rats. Combination of water extract of *Andrographis paniculata* and water extract of *Syzygium polyanthum* (6:1) have been reported to reduce blood glucose in normal and alloxan rats. This combination also gave protection effects on kidney, liver and pancreas animals tested from alloxan toxicity<sup>11</sup>.

In this study the prolonged effect (up to 7 day) of the water extract of *Andrographis paniculata*, water extract of *Syzygium polyanthum* and its combination in fasting blood glucose (FBG) and biochemical parameters such as serum total cholesterol (TC), LDL, HDL, triglyceride, and

alkaline phosphatase were studied in alloxan induced diabetic rats.

## 2. Materials and Methods

### 2.1. Plant Material

The plant material of *Andrographis paniculata* and *Syzygium polyanthum* were taken from Natura Laboratoria Prima on dried powder form. Natura Laboratoria Prima is a nature-based health Products Company in Indonesia, which produce nature-based health products, one of them is quality dried extracts from plant materials.

16

### 2.2. Animals

Male wistar albino rats (8–10 weeks) were obtained from the animal use of Widya Mandala Catholic University. Before and during the experiment, rats were fed with standard diet. After randomization into various groups and before initiation of experiment, the rats were acclimatized for a period of 7 days under standard environmental conditions of temperature, relative humidity, and dark/light cycle. Animals described as fasting were deprived of food and water for 12 hours ad libitum.

### 2.3. Experimental Design

Eight groups of rats, ten in each received the following treatment schedule:

Group I: Normal control (CMC Na 0.5%).

Group II: Alloxan treated control (100 mg/kg.im).

Group III: Alloxan (100 mg/kg.im) + the water extract of *Andrographis paniculata* (200 mg/kg, p.o),

Group IV: Alloxan (100 mg/kg.im) + the water extract of *Syzygium polyanthum* (200mg/kg, p.o),

Group V: Alloxan (100 mg/kg.im) + Combination of water extract of *Andrographis paniculata* and water extract of *Syzygium polyanthum* (1:6; 200mg/kg, p.o),

Group VI: Alloxan (100 mg/kg.im) + Combination of water extract of *Andrographis paniculata* and water extract of *Syzygium polyanthum* (1:6; 200mg/kg, p.o),

Group VII: Alloxan (100 mg/kg.im) + Standard drug, Metformin (63 mg/kg, p.o)

Group VIII: Alloxan (100 mg/kg.im) + Standard drug, insulin (0.1 unit/Kg/day, subcutan)

The extracts, standard drug, and CMC Na 0.5% were administered with the help of feeding cannula. Group I serve as normal control, which received CMC Na 0.5% for 7 days. Group II to Group VIII are diabetic control rats. Group III to Group VIII (which previously received alloxan) are

treatment groups given a fixed dose for 7 consecutive days.

### 2.4. Induction of Diabetes in Experimental Animals

Rats were made diabetic by a single intraperitoneal injection of alloxan monohydrate (100 mg/kgBW)<sup>12</sup>. Alloxan was first weighed individually for each animal according to the body weight and then solubilized with 0.2 saline (154mM NaCl) just prior to injection. Two days after alloxan injection, rats with plasma glucose levels of >200 mg/dl were included in the study. Treatment with plant extracts was started 24 h after alloxan injection.

### 2.5. Collection of Blood Sample and Blood Glucose Determination

Pod samples were drawn from tail tip of rat. Fasting blood glucose estimation and body weight measurement were done on day 1 and 7 of the study. Blood glucose estimation can be done by one touch electronic glucometer using glucose test strips. On day 7, serum was separated and analyzed for serum cholesterol<sup>13</sup>, serum triglycerides by enzymatic DHBS colorimetric method<sup>14</sup> serum HDL<sup>15</sup>, serum LDL<sup>16</sup>, and serum alkaline phosphatase hydrolyzed phenol amino antipyrine method<sup>17</sup>.

### 2.6. Statistical Analysis.

All the values of body weight, fasting blood sugar, and biochemical estimations were expressed as mean ± standard error of mean (S.E.M.) and analyzed for ANOVA and post hoc Dunnet's t-test. Differences between groups were considered significant at P < .01 levels.

## 3. Results and Discussion

The results of antidiabetic activity of the water extract of *Andrographis paniculata*, the water extract of *Syzygium polyanthum* and its combination on fasting blood glucose, animal body weight and serum profile have been shown in Table 1-2. Vehicle control animals were found to be slightly decreased in their body weight but diabetic rats showed significant reduction in body weight during 7 days (Table 1). Alloxan caused body weight reduction, where the standard drugs and the treated groups cannot repair this condition.

Alloxan acts as a cytotoxin for beta-cells of the islet of langerhans, causes diabetes by inducing cell necrosis<sup>18-19</sup>. The Reactive Oxygen Species

mediates the cytotoxic action with the increase in cytosolic calcium concentration, leading to rapid beta-cells destruction<sup>20</sup>. This result into decreased insulin secretion and elevated blood glucose level<sup>21</sup>. This experimental study reveals that alloxan-treated rats received the water extract of *Andrographis paniculata*, water extract of *Syzygium polyanthum* and its combination showed lower blood glucose level as compared to the diabetic control group may be due to the possibility that few of beta cells are still surviving and stimulated by extract component (s), releasing insulin.

The extracts exhibited significant reduction of serum cholesterol level in alloxan-treated rats. The abnormal high concentration of serum lipids in the diabetic subject is mainly due to increase in the mobilization of free fatty acids from the peripheral fat depots, since insulin inhibits the hormone sensitive lipase whereas glucagon and epinephrine promote lipolysis<sup>18</sup>. This diabetic dyslipidemia is, therefore, regarded as a consequence of the unregulated actions on LPL on fat depot<sup>22</sup>. The results indicate that secondary metabolites in both extracts may exert their role in maintenance of the cholesterol profile, especially in LDL and HDL<sup>23-25</sup>.

Proteolysis, lipolysis and acute fluid loss during diabetes pave the way for weight loss<sup>26</sup>. The

weight gain in extract treated groups reflects the correction of body metabolism. Results reveal that the water extract of *Andrographis paniculata*, water extract of *Syzygium polyanthum* and its combination exhibits the antidiabetic activity in a dose dependent manner. In conclusion, this study indicates that the water extract of *Andrographis paniculata*, water extract of *Syzygium polyanthum* and its combination possess significant antidiabetic activity along with antihypercholesterolemic activity. The active principle (s) in the extracts may have better performance if isolated and purified form. Therefore, further investigation is necessary to determine the exact phytoconstituents (s) responsible for antidiabetic effect.

#### Acknowledgment

The authors sincerely thank to Indonesian Government for funding this research.

Table 1 Effect of various groups of water extract of *Andrographis paniculata*, water extract of *Syzygium polyanthum* and its combination on fasting blood glucose and animal body weight in alloxan (100 mg/kg, i.m.) induced diabetic albino rats after 7 days of treatment.

Groups	Animals Body Weight (gram)		Fasting blood glucose (mg/dl)	
	Day 1	Day 7	Day 1	Day 7
Normal control	253	242	88	74
Diabetic control	136	116	221	566
Alloxan + water extracts of <i>Syzygium polyanthum</i> leaves	169	142	390	75
Alloxan + water extracts of <i>Andrographis paniculata</i> aerial parts	188	146	367	62
Alloxan + combination 1 : 6 (water extracts of <i>Syzygium polyanthum</i> leaves : water extracts of <i>Andrographis paniculata</i> aerial parts)	196	151	343	98
Alloxan + combination (6 : 1 (water extracts of <i>Syzygium polyanthum</i> leaves : water extracts of <i>Andrographis paniculata</i> aerial parts)	178	143	488	75
Alloxan + metformin	172	134	485	59
Alloxan + insulin	228	181	323	134

Table 2 Effect of various groups of water extract of *Andrographis paniculata*, water extract of *Syzygium polyanthum* and its combination on serum profile in alloxan (100 mg/kg, i.m.) induced diabetic albino rats after 7

Groups	Cholesterol (mg/dl)	HDL (mg/dl)	LDL (mg/dl)	Triglyceride (mg/dl)	Alkaline phosphatase (mg/dl)
Normal control	32.33 ± 2.51	59.67 ± 2.31	27.11 ± 0.58	30.22 ± 3.21	232.56 ± 6.36
Diabetic control	40 ± 1.41	82.75 ± 2.12	32.5 ± 0.71	38.50 ± 7.07	226.33 ± 10.61
Alloxan + water extracts of <i>Syzygium polyanthum</i> leaves	54.60*	46.80*	22.50*	29.33*	227
Alloxan + water extracts of <i>Andrographis paniculata</i> aerial parts	30.67 ± 1.41*	34 ± 2.12*	24 ± 0.71*	28.57 ± 0.00*	287.29 ± 1.41*
Alloxan + combination 1 : 6 (water extracts of <i>Syzygium polyanthum</i> leaves : water extracts of <i>Andrographis paniculata</i> aerial parts)	43.25 ± 4.43*	36.50 ± 1.00*	23.33 ± 4.95*	38 ± 5.68	261.50 ± 2.12*
Alloxan + combination (6 : 1 (water extracts of <i>Syzygium polyanthum</i> leaves : water extracts of <i>Andrographis paniculata</i> aerial parts)	45.40 ± 7.79*	28.20 ± 2.12*	32.83 ± 0.70	28.17 ± 3.54*	284.83 ± 2.12*
Alloxan + metformin	39.17 ± 0.71	32.83 ± 2.12*	22.17 ± 5.65*	37.67 ± 3.46	211.83 ± 12.02*
Alloxan + insulin	37.20 ± 1.41*	51.80 ± 0.71	26.86 ± 0.71	30.29 ± 1.41*	406.86 ± 1.41*

3 Values are given as mean ± SEM for groups of ten animals each \*P < .01 (Dunnet t-test). Diabetic control was compared with the vehicle control and extract treated groups were compared with the diabetic control.

#### References

- Vats V, Yadav SP, Grover JK. Ethanolic extract of *Ocimum sanctum* leaves partially attenuates streptozotocin induced alterations in glycogen content and carbohydrate metabolism in rats. *J Ethnopharmacol* 2004; 90(1):155–160.
- Kumar GPS, Arulselvan P, Kumar DS, Subramanian SP. Anti-diabetic activity of fruits of *Terminalia chebula* on streptozotocin induced diabetic rats. *J Health Sci* 2006; 52 (3):283–291.
- Zimmet, Paul KGMM, Alberti, Shaw J. Global and societal implications of the diabetes epidemic. *Nature* 2001; 414: 782-787.
- Wild S, Roglic G, Green A, Sicree R, King H. Global Prevalence of Diabetes. *Diabetes Care* 2004; 27(5): 1047-1053.
- Sumana G, Suryawashi SA. Effect of vinca rosea extracts in treatment of alloxan diabetes in male albino rats. *Indian J Exp Biol* 2001; 39:748–758.
- Holman RR, Turner RC. Oral agents and insulin in the treatment of NIDDM. In: Pickup J,

- Williams G, editor. Text Book of Diabetes. UK: Blackwell; 1991. P. 467-469.
- Kameswara RB, Kesavulu MM, Apparao C. Antihyperglycemic activity of *Momordica cymbalaria* in alloxan diabetic rats. *J Ethnopharmacol* 2001; 78(1):67-71.
- Kameswarao B, Giri R, Kesavulu MM, Apparao C. Herbal medicine. In: Bajaj JS, editor. *The Management by Indigenous Resources, Diabetes Mellitus in Developing Countries India*: Interprint; 1997. P. 375-377.
- Sandhya B, Thomas S, Isabel W, Shenbagarathai R. Ethnomedicinal plants used by the Valaiyan community of piramalai hills (Reserved forest), Tamil Nadu, India -A pilot study. *African Ethnomedicines. Afr J Tradit Complem* 2006; 3:101-114.
- Prohp TP, Onoagbe IO, Onyebuagu PC, Omeni AA, Okoli RI, Obeto NP. Effect of *Triplochiton scleroxylon* on red blood cells and associated parameters in alloxan-induced diabetic rabbits. *Pak J Nutr* 2006; 5(5): 425-428.
- Widjakusuma EC, Sudjardhana A, Arianto. Antidiabetic Activity of *Andrographis paniculata*, *Syzygium polyanthum* and its combination in alloxan induced diabetic rats. Research Report 2011. Indonesia: Widya Mandala Catholic University.
- Aruna RV, Ramesh B, Kartha VNR. Effect of beta-carotene on protein glycosylation in alloxan induced diabetic rats. *Indian J Exp Biol* 1999; 37(4): 399-401.
- Roeschlau P, Bernt E, Gruber W. Enzymatic determination of total cholesterol in serum. *Zeitschrift für klinische Chemie und Klinische Biochemie* 1974; 12(5): 226.
- Mueller PH, Schmuelling RM, Liebich HM. A fully enzymatic triglyceride determination. *J Clin Chem Clin Biochem* 1977; 15(9): 457-464.
- Allain CC, Poon LS, Chan CSG. Enzymatic determination of total serum cholesterol. *Clin Chem* 1974; 20(4):470-475.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972; 18(6): 499-502.
- Sasaki M. A new ultramicro method for the determination of serum alkaline phosphatase. Use of Berthelot's reaction for the estimation of phenol released by enzymatic activity. *Igaku To Seibutsugaku* 1966; 70(4):208-214.
- Jorns AR, Munday MT, Lenzen S. Comparative toxicity of alloxan, N-alkyl-alloxans and ninhydrin to isolated pancreatic islets in vitro. *J Endocrinol* 1997; 155: 283-293.
- Ledoux SP, Woodley SE, Patton NJ, Wilson LG. Mechanism of nitrosourea-induced-cells damage-alterations in DNA. *Diabetes* 1986; 35: 866-872.
- Szkudelski T. The mechanism of Alloxan and Streptozotocin action in B cells of the rat pancreas. *Physiol Res* 2001; 50: 537-546.
- Deewanjee S, Bose SK, Sahu R, Mandal SC. Antidiabetic effect of matured fruits of *Diospyros peregrine* in alloxan-induced diabetic rats. *Int J Green Pharm* 2008; 2(2): 95-99.
- Hussain HEMA. Reversal of diabetic retinopathy in streptozotocin induced diabetic rats using traditional Indian antidiabetic plant *Acalypha indica* (L). *Indian J Clin Biochem* 2002; 17(2): 115-123.
- Setchell KDR. Phytoestrogens, The biochemistry, physiology and implications for human health of soy phytoestrogens. *Am J Clin Nutr* 1998; 68: 1333S-1348S.
- Sen S, Makkar HPS, Becker K. Alfalfa saponins and their implication in animal nutrition. *J Agr Food Chem* 1998; 46: 131-140.
- Francis G, Kerem Z, Makkar HPS, Becker K. The biological action of saponins in animal systems. *Brit J Nutr* 2002; 88: 587-605.
- Alberti KGMM, Zimmet PZ. Definition, Diagnosis and Classification of Diabetes Mellitus and its Complications. Part 1: Disguises and Classification of Diabetes Mellitus Provisional Report of WHO consultations. *Diabet Med* 1998; 15: 539-553.

# The Lipid Lowering Effect of Water Extract of Andrographis paniculata, Water Extract of Syzigium polyanthum and its Combination in Alloxan-Induced Diabetic Rats

## ORIGINALITY REPORT

% **19**  
SIMILARITY INDEX

% **15**  
INTERNET SOURCES

% **12**  
PUBLICATIONS

% **10**  
STUDENT PAPERS

## PRIMARY SOURCES

**1** [ijpjournal.com](http://ijpjournal.com) %**2**  
Internet Source

**2** [www.ijddr.in](http://www.ijddr.in) %**2**  
Internet Source

**3** Safiyeh Soleimani ., Fathallah Fathi Azarbaizani ., Vahid Nejati .. "The Effect of Equisetum arvense L. (Equisetaceae) in Histological Changes of Pancreatic  $\beta$ Cells in Streptozotocin-Induced Diabetic in Rats", Pakistan Journal of Biological Sciences, 2007 %**2**  
Publication

**4** [ijipls.com](http://ijipls.com) %**2**  
Internet Source

**5** [www.ugcfrp.ac.in](http://www.ugcfrp.ac.in) %**2**  
Internet Source

**6** Submitted to Jawaharlal Nehru Technological University %**2**  
Student Paper

7	<a href="http://etheses.saurashtrauniversity.edu">etheses.saurashtrauniversity.edu</a> Internet Source	% 1
8	<a href="http://www.wjpr.net">www.wjpr.net</a> Internet Source	% 1
9	<a href="http://shodhganga.inflibnet.ac.in">shodhganga.inflibnet.ac.in</a> Internet Source	% 1
10	"Abstracts of the IDF Congress in Paris 2003", <i>Diabetologia</i> , 2003 Publication	% 1
11	<a href="http://irjpas.com">irjpas.com</a> Internet Source	% 1
12	Submitted to North South University Student Paper	% 1
13	Fengjie Huang. "Antidiabetic effect of a new peptide from <i>Squalus mitsukurii</i> liver (S-8300) in streptozocin-induced diabetic mice", <i>Journal of Pharmacy and Pharmacology</i> , 12/2005 Publication	% 1
14	<a href="http://citeseerx.ist.psu.edu">citeseerx.ist.psu.edu</a> Internet Source	% 1
15	<a href="http://www.unisa.it">www.unisa.it</a> Internet Source	% 1
16	Submitted to University of Leicester Student Paper	% 1

17

El Ghouli, J., N. A. Boughattas, and M. Ben-Attia. "Antihyperglycemic and antihyperlipidemic activities of ethanolic extract of *Zygophyllum album* in streptozotocin-induced diabetic mice", *Toxicology and Industrial Health*, 2012.

Publication

% 1

18

[ijpba.info](http://ijpba.info)

Internet Source

<% 1

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 10 WORDS