

LAMPIRAN A

CONTOH PERHITUNGAN PERSENTASE HASIL SINTESIS

I. Perhitungan berat teoritis

a. Ammonium tiosianat (BM : 76,12 g/mol)

Penimbangan : 1,14 gram

$$\text{mmol ammonium tiosianat : } \frac{1,14}{76,12} \times 1000 = 14,98 \text{ mmol} \cong 15 \text{ mmol}$$

b. 2-klorobenzoil klorida (BM : 175,01 g/mol, berat jenis : 1,378 g/cm³)

Volume : 1,3 ml

$$\text{mmol 2-klorobenzoil klorida : } \frac{1,3 \times 1,378}{175,01} \times 1000 = 10,2 \text{ mmol} \cong 10 \text{ mmol}$$

c. Amonia (BM : 35,05 g/mol, berat jenis : 0,91 g/cm³)

Volume : 1,15 ml

$$\text{mmol amonia : } \frac{1,15 \times 0,91}{35,05} \times 1000 = 30 \text{ mmol}$$

II. Perhitungan persentase hasil sintesis berdasarkan mmol teoritis

Persentase hasil N-2-klorobenzoilthiourea :

ammonium tiosianat + 2-klorobenzoil klorida $\rightarrow$ 2-klorobenzo-ilisotiosianat + NH₄Cl

awal	15 mmol	10 mmol	0	0	
reaksi	10 mmol	10 mmol -	10 mmol	10 mmol	+
sisa	5 mmol	0	10 mmol	10 mmol	



awal	10 mmol	30 mmol	0	
reaksi	10 mmol	10 mmol -	10 mmol	+
sisa	0	20 mmol	10 mmol	

BM teoritis = 210,5

Massa teoritis = 10 mmol x 210,5 = 2,10 gram

Massa praktis = 1,22 gram

% hasil = $\frac{1,22}{2,10} \times 100\% = 58,09\% \approx 58\%$

LAMPIRAN B
UJI ANOVA 1

	Sum of Squares	Df	Mean Square	F	Sig.
Between Groups	82.667	2	41.333	74.400	.000
Within Groups	3.333	6	.556		
Total	86.000	8			

Post Hoc Tests

Multiple Comparisons

Dependent Variable: Hasil
Tukey HSD

(I) senyawa	(J) senyawa	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
		Lower Bound	Upper Bound	Lower Bound	Upper Bound	Lower Bound
1	2	-7.333(*)	.609	.000	-9.20	-5.47
	3	-4.667(*)	.609	.001	-6.53	-2.80
2	1	7.333(*)	.609	.000	5.47	9.20
	3	2.667(*)	.609	.011	.80	4.53
3	1	4.667(*)	.609	.001	2.80	6.53
	2	-2.667(*)	.609	.011	-4.53	-.80

* The mean difference is significant at the .05 level.

Homogeneous Subsets

Hasil

Tukey HSD


senyawa	N		Subset for alpha = .05		
	1		2	3	1
1	3		57.00		
3	3			61.67	
2	3				64.33
Sig.			1.000	1.000	1.000

Means for groups in homogeneous subsets are displayed.

a Uses Harmonic Mean Sample Size = 3.000.

LAMPIRAN C

SKEMA KERJA SINTESIS TURUNAN N-2-KLORO BENZOILTIOUREA




Keterangan *:

Sintesis 2-kloro-N-(morfolin-4-karbonotiol)benzamida : 2.59 ml (30 mmol)

Morfolin

Sintesis 2-kloro-N-(piperazin-1-karbonotiol)benzamida : 2.58 g (30 mmol)


Piperazin




LAMPIRAN D

ESTIMASI RMI-¹H SENYAWA N-2-KLOROBENZOILTIOUREA

ChemNMR ¹H Estimation



Estimation quality is indicated by color: **good**, **medium**, **rough**




Protocol of the H-1 NMR Prediction:

Node	Shift	Base + Inc.	Comment (ppm rel. to TMS)
NH	11.44	8.00	sec. amide
		3.44	general corrections
NH2	9.53	2.00	amine
		7.53	general corrections
CH	7.67	7.26	1-benzene
		0.01	1 -Cl
		0.18	1 -C(=O)N
		0.22	general corrections
CH	7.43	7.26	1-benzene
		-0.06	1 -Cl
		-0.69	1 -C(=O)N
		-0.46	general corrections
CH	7.64	7.26	1-benzene
		-0.06	1 -Cl
		0.25	1 -C(=O)N
		0.19	general corrections
CH	7.51	7.26	1-benzene
		-0.12	1 -Cl
		0.18	1 -C(=O)N
		0.19	general corrections


LAMPIRAN E

ESTIMASI RMI-¹H SENYAWA 2-KLORO-N-(MORFOLIN-4-KARBONOTIOL)BENZAMIDA

ChemNMR ¹H Estimation



Estimation quality is indicated by color: **good**, **medium**, **rough**




Protocol of the H-1 NMR Prediction:

Node	Shift	Base + Inc.	Comment (ppm rel. to TMS)
NH	10.57	8.00	sec. amide
		2.57	general corrections
CH2	3.65	3.67	tetrahydro-1,4-oxazine
		-0.02	general corrections
CH2	3.65	3.67	tetrahydro-1,4-oxazine
		-0.02	general corrections
CH2	3.15	2.87	tetrahydro-1,4-oxazine
		?	1 -R from N-CHx
		0.28	general corrections
CH2	3.15	2.87	tetrahydro-1,4-oxazine
		?	1 -R from N-CHx
		0.28	general corrections
CH	7.67	7.26	1-benzene
		0.01	1 -Cl
		0.18	1 -C(=O)N
		0.22	general corrections
CH	7.43	7.26	1-benzene
		-0.06	1 -Cl
		0.69	1 -C(=O)N
		-0.46	general corrections
CH	7.64	7.26	1-benzene
		-0.06	1 -Cl
		0.25	1 -C(=O)N
		0.19	general corrections
CH	7.51	7.26	1-benzene
		-0.12	1 -Cl
		0.18	1 -C(=O)N
		0.19	general corrections


LAMPIRAN F

ESTIMASI RMI-¹H SENYAWA 2-KLORO-N-(PIPERAZIN-1-KARBONOTIOL)BENZAMIDA

ChemNMR ¹H Estimation



Estimation quality is indicated by color: good, medium, rough



Protocol of the H-1 NMR Prediction:

Node	Shift	Base + Inc.	Comment (ppm rel. to TMS)
NH	2.0	2.00	amine
NH	10.57	8.00	sec. amide
		2.57	general corrections
CH2	3.71	1.37	methylene
		2.12	1 alpha -NC(=S)NR
		0.08	1 beta -N-C
		0.14	general corrections
CH2	3.71	1.37	methylene
		2.12	1 alpha -NC(=S)NR
		0.08	1 beta -N-C
		0.14	general corrections
CH2	2.67	1.37	methylene
		1.22	1 alpha -N-C
		0.08	1 beta -N-C
CH2	2.67	1.37	methylene
		1.22	1 alpha -N-C
		0.08	1 beta -N-C
CH	7.67	7.26	1-benzene
		0.01	1 -Cl
		0.18	1 -C(=O)N
		0.22	general corrections
CH	7.43	7.26	1-benzene
		-0.06	1 -Cl
		0.69	1 -C(=O)N
		-0.46	general corrections
CH	7.64	7.26	1-benzene
		-0.06	1 -Cl
		0.25	1 -C(=O)N
		0.19	general corrections
CH	7.51	7.26	1-benzene
		-0.12	1 -Cl
		0.18	1 -C(=O)N
		0.19	general corrections

