

Effect of butterfly pea (*Clitoria ternatea*) flower extract on qualities, sensory properties, and antioxidant activity of wet noodles with various composite flour proportions

by Paini Sri Widyawati

Submission date: 20-Aug-2024 04:14PM (UTC+0700)

Submission ID: 2434935397

File name: Effect_of_butterfly_pea_Clitoria_ternatea_flower_ectract_on.pdf (3.17M)

Word count: 12357

Character count: 63189

Effect of butterfly pea (*Clitoria ternatea*) flower extract on qualities, sensory properties, and antioxidant activity of wet noodles with various composite flour proportions

Paini Sri Widyawati^{1*}, Thomas Indarto Putut Suseno¹, Felicia Ivana¹, Evelyne Natania¹ and Sutee Wangtueai²

¹ Food Technology Study Program, Faculty of Agricultural Technology, Widya Mandala Surabaya Catholic University, Dinoyo Street Number 42-44, Surabaya, 60265, Indonesia

² College of Maritime Studies and Management, Chiang Mai University, Samut Sakhon 74000, Thailand

* Corresponding author, E-mail: paini@ukwms.ac.id

Abstract

Improvement of wet noodles' qualities, sensory, and functional properties was carried out using a composite flour base with the butterfly pea flower extract added. The composite flour consisted of wheat flour and stink lily flour, and κ -carrageenan at ratios of 80:20:0 (K0), 80:19:1 (K1), 80:18:2 (K2), and 80:17:3 (K3) (% w/w) was used with concentrations of butterfly pea extract of 0 (T1), 15 (T15), and 30 (T30) (% w/v). The research employed a randomized block design with two factors, namely the composite flour and the concentration of butterfly pea flower extract, and resulted in 12 treatment combinations (K0T0, K0T15, K0T30, K1T0, K1T15, K1T30, K2T0, K2T15, K2T30, K3T0, K3T15, K3T30). The interaction of the composite flour and butterfly pea flower extract significantly affected the color profile, sensory properties, bioactive compounds, and antioxidant activities of wet noodles. However, each factor also significantly influenced the physical properties of wet noodles, such as moisture content, water activity, tensile strength, swelling index, and cooking loss. The use of κ -carrageenan up to 3% (w/w) in the mixture increased moisture content, swelling index, and tensile strength but reduced water activity and cooking loss. K3T30 treatment with composite flour of wheat flour-stink lily flour- κ -carrageenan at a ratio of 80:17:3 (% w/w) had the highest consumer acceptance based on hedonic sensory score.

Citation: Widyawati PS, Suseno TIP, Ivana F, Natania E, Wangtueai S. 2024. Effect of butterfly pea (*Clitoria ternatea*) flower extract on qualities, sensory properties, and antioxidant activity of wet noodles with various composite flour proportions. *Beverage Plant Research* 4: e022 <https://doi.org/10.48130/bpr-0024-0011>

Introduction

The use of composite flour in wet noodles has been widely used to increase its functional value and several characteristics, including physical, chemical, and sensory properties. Siddeeg et al.^[1] used wheat-sorghum-guar flour and wheat-millet-guar flour to increase the acceptability of wet noodles. Efendi et al.^[2] stated that potato starch and tapioca starch in a ratio of 50:50 (% w/w) could increase the functional value of wet noodles. Dhull & Sandhu^[3] stated that noodles made from wheat flour mixed with up to 7% fenugreek flour produce good texture and high consumer acceptability. Park et al.^[4] utilized the mixed ratio of purple wheat bran to improve the quality of wet noodles and antioxidant activity.

A previous study used stinky lily flour or konjac flour (*Amorophallus muelleri*) composited with wheat flour to increase the functional value of noodles by increasing biological activity (anti-obesity, anti-hyperglycemic, anti-hyper cholesterol, and antioxidant) and extending gastric emptying time^[5,6]. Widyawati et al.^[7] explained that using composite flour consisting of wheat flour, stink lily flour, and κ -carrageenan can improve the swelling index, total phenolic content (TPC), total flavonoid content (TFC), and 2,2-diphenyl-1-picrylhydrazyl free radical scavenging activity (DPPH), which influences the effectiveness of bioactive compounds in the composite flour that serve as antioxidant sources of wet noodles. Therefore, other ingredients containing phenolic compounds can be added to increase

composite flour's functional values as a source of antioxidants. Czajkowska-González et al.^[8] mentioned that incorporating phenolic antioxidants from natural sources can improve the functional values of bread. Widyawati et al.^[7] added pluchea extract to increase the TPC, TFC, and DPPH of wet noodles, however, this resulted in an unattractive wet noodle color. Therefore, it is necessary to incorporate other ingredients to enhance the wet noodles' color profile and their functional properties, one of which is the butterfly pea flower.

Butterfly pea (*Clitoria ternatea*) is a herb plant from the Fabaceae family with various flower colors, such as purple, blue, pink, and white^[9]. This flower has phytochemical compounds that are antioxidant sources^[10,11], including anthocyanins, tannins, phenolics, flavonoids, flobatannins, saponins, triterpenoids, anthraquinones, sterols, alkaloids, and flavonol glycosides^[12,13]. Anthocyanins of the butterfly pea flower have been used as natural colorants in many food products^[14,15], one of them is wet noodles^[16,17]. The phytochemical compounds, especially phenolic compounds, can influence the interaction among gluten, amylose, and amylopectin, depending on partition coefficients, keto-groups, double bonds (in the side chains), and benzene rings^[18]. This interaction involves their formed covalent and non-covalent bonds, which influenced pH and determined hydrophilic-hydrophobic properties and protein digestibility^[19]. A previous study has proven that the use of phenolic compounds from plant extracts, such as pluchea leaf^[7,20], gendarussa leaf (*Justicia gendarussa*

Burm.F.)^[21], carrot and beetroot^[22], kelakai leaf^[23] contributes to the quality, bioactive compounds, antioxidant activity, and sensory properties of wet noodles. Shiau et al.^[17] utilized the natural color of butterfly pea flower extract to make wheat flour-based wet noodles, resulting in higher total anthocyanin, polyphenol, DPPH, and ferric reducing antioxidant power (FRAP) than the control samples. This extract also improved the color preference and reduced cooked noodles' cutting force, tensile strength, and extensibility. Until now, the application of water extract of butterfly pea flowers in wet noodles has been commercially produced, but the interactions among phytochemical compounds and ingredients of wet noodles base composite flour (stinky lily flour, wheat flour, and κ -carrageenan) have not been elucidated. Therefore, the current study aimed to determine the effect of composite flour and butterfly pea flower extract on wet noodles' quality, bioactive content, antioxidant activity, and sensory properties.

46

Materials and methods

Raw materials and preparation

Butterfly pea flowers were obtained from Penjaringan Sari Garden, Wonorejo, Rungkut, Surabaya, Indonesia. The flowers were sorted, washed, dried under open sunlight, powdered using a blender (Philips HR2116, PT Philips, Netherlands) for 3 min, and sieved using a sieve shaker with 45 mesh size (analytic sieve shaker OASS203, Decent, Qingdao Decent Group, China). The water extract of butterfly pea flower was obtained using a hot water extraction at 95 °C for 3 min based on the modified method of Widyawati et al.^[20] and Putri et al.^[24] to obtain three concentrations of butterfly pea extract: 0 (T1), 15 (T15), and 30 (T30) (% w/v). The three-composite flour proportions were prepared by mixing wheat flour (Cakra Kembar, PT Bogasari Sukses Makmur, Indonesia), stinky lily flour (PT Rezka Nayatama, Sekotong, Lombok Barat, Indonesia), and κ -carrageenan (Sigma-Aldrich, St. Louis, MO, USA) at ratios of 80:20:0 (K0), 80:19:1 (K1), 80:18:2 (K2), and 80:17:3 (K3) (% w/w).

Chemicals and reagents

Gallic acid, 2,2-diphenyl-1-picrylhydrazyl (DPPH), (+)-catechin, and sodium carbonate were purchased from Sigma-Aldrich (St. Louis, MO, USA). Methanol, aluminum chloride, Folin-Ciocalteu's phenol reagent, chloride acid, acetic acid, sodium acetic, sodium nitric, sodium hydroxide, sodium hydrogen phosphate, sodium dihydrogen phosphate, potassium ferricyanide, chloroacetic acid, and ferric chloride were purchased from Merck (Kenilworth, NJ, USA). Distilled water was purchased from a local market (PT Aqua Surabaya, Surabaya, Indonesia).

Wet noodle preparation

Wet noodles were prepared based on the modified formula of Panjaitan et al.^[25], as shown in Table 1. In brief, the composite flour was sieved with 45 mesh size, weighed, and mixed with butterfly pea flower extract at various concentrations. Salt, water, and fresh whole egg were then added and kneaded to form a dough using a mixer (Oxone Master Series 600 Standing Mixer OX 851, China) until the dough obtained was smooth. The dough was sheeted to get noodles about 0.15 cm thick and cut using rollers equipped with cutting blades (Oxone OX355AT, China) to obtain noodles about 0.1 cm wide. Raw wet noodle strains were sprinkled with tapioca flour (Rose Brand, PT Budi Starch & Sweetener, Tbk) (4% w/w) before being heated in

Table 1. Formula of wet noodles.

Treatment	Code	Ingredients				
		Salt (g)	Fresh whole egg (g)	Water (mL)	Butterfly pea extract solution (mL)	Composite flour (g)
1	K0T0	3	30	30	0	150
2	K0T15	3	30	0	30	150
3	K0T30	3	30	0	30	150
4	K1T0	3	30	30	0	150
5	K1T15	3	30	0	30	150
6	K1T30	3	30	0	30	150
7	K2T0	3	30	30	0	150
8	K2T15	3	30	0	30	150
9	K2T30	3	30	0	30	150
10	K3T0	3	30	30	0	150
11	K3T15	3	30	0	30	150
12	K3T30	3	30	0	30	150

K0 = wheat flour : stinky lily flour : κ -carrageenan = 80:20:0 (%w/w). K1 = wheat flour : stinky lily flour : κ -carrageenan = 80:19:1 (%w/w). K2 = wheat flour : stinky lily flour : κ -carrageenan = 80:18:2 (%w/w). K3 = wheat flour : stinky lily flour : κ -carrageenan = 80:17:3 (%w/w). T0 = concentration of the butterfly pea extract = 0%. T15 = concentration of the butterfly pea extract = 15%. T30 = concentration of the butterfly pea extract = 30%.

boiled water (100 °C) with a ratio of raw noodles : water at 1:4 w/v for 2 min. Cooked wet noodles were coated with palm oil (Sania, PT Wilmar Nabati, Indonesia) (5% w/w) before being subjected to quality and sensory properties measurements, whereas uncooked noodles without oil coating were used to analyze bioactive compounds and antioxidant activity.

Extraction of bioactive compounds of wet noodles

Wet noodles were extracted based on the method of Widyawati et al.^[7]. Raw noodles were dried in a cabinet dryer (Gas drying oven machine OVG-6 SS, PT Agrowindo, Indonesia) at 60 °C for 2 h. The dried noodles were ground using a chopper (Dry Mill Chopper Philips set HR 2116, PT Philips, Netherlands). About 20 g of sample was mixed with 50 mL of solvent mixture (1:1 v/v of methanol/water), stirred at 90 rpm in a shaking water bath at 35 °C for 1 h, and centrifuged at 5,000 rpm for 5 min to obtain supernatant. The obtained residue was re-extracted in an extraction time for three intervals. The supernatant was collected and separated from the residue and then evaporated using a rotary evaporator (Buchi-rotary evaporator R-210, Germany) at 70 rpm, 70 °C, and 200 mbar to generate a concentrated wet noodle extract. The obtained extract was used for further analysis.

Moisture content analysis

The water content of cooked wet noodles was analyzed using the thermogravimetric method^[26]. About 1 g of the sample was weighed in a weighing bottle and heated in a drying oven at 105–110 °C for 1 h. The processes were followed by weighing the sample and measuring moisture content after obtaining a constant sample weight. The moisture content was calculated based on the difference of initial and obtained constant sample weight divided by the initial sample weight, expressed as a percentage of wet base.

Water activity analysis

The water activity of cooked wet noodles was analyzed using an A_w -meter (Water Activity Hygropalm HP23 A_w a set 40 Rotronic, Swiss). Ten grams of the sample were weighed, put into an A_w meter chamber, and analyzed to obtain the sample's water activity^[27].

Tensile strength analysis

Tensile strength is an essential parameter that measures the extensibility of cooked wet noodles^[28]. About 20 cm of the sample was measured for its tensile strength using a texture analyzer equipped with a Texture Exponent Lite Program and a noodle tensile rig probe (TA-XT Plus, Stable Microsystem, UK). The noodle tensile rig was set to pre-set speed, test speed, and post-test speed at 1, 3, and 10 mm/s, respectively. Distance, time, and trigger force were set to 100 mm, 5 s, and 5 g, respectively.

Color analysis

Ten grams of cooked wet noodles were weighed in a chamber, and the color was analyzed using a color reader (Konica Minolta CR 20, Japan) based on the method of Harijati et al.^[29]. The parameters measured were lightness (L^*), redness (a^*), yellowness (b^*), hue ($^{\circ}h$), and chroma (C). L^* value ranged from 0-100 expresses brightness, and a^* value shows red color with an interval between -80 and +100. b^* value represents a yellow color with an interval of -70 to +70^[30]. C indicates the color intensity and $^{\circ}h$ states the color of samples^[31].

Swelling index analysis

The swelling index was determined using a modified method of Islamiya^[32]. Approximately 5 g of the raw wet noodles were weighed in a chamber and cooked in 150 mL boiled water (100 °C) for 5 min. The swelling index was measured to observe the capability of raw wet noodles to absorb water that increased the weight of raw wet noodles^[33]. The swelling index was measured from the difference in noodle weights before and after boiling.

Cooking loss analysis

The cooking loss of the raw wet noodles was analyzed using a modified method of Aditia et al.^[34]. The cooking loss expresses the weight loss of wet noodles during cooking, indicated by the cooking water that turns cloudy and thick^[35]. About 5 g of the raw wet noodles was weighed in a chamber and cooked in 150 mL boiled water (100 °C) for 5 min. Then, the sample was drained and dried in a drying oven at 105 °C until the weight of the sample was constant.

Total phenolic content analysis

The total phenolic content of the wet noodles was determined using Folin-Ciocalteu's phenol reagent based on the modified method by Ayele et al.^[36]. About 50 μ L of the extract was added with 1 mL of 10% Folin-Ciocalteu's phenol reagent in a 10 mL volumetric flask, homogenized, and incubated for 5 min. Then, 2 mL of 7.5% Na_2CO_3 was added, and the volume was adjusted to 10 mL with distilled water. The solution's absorbance was measured spectrophotometrically at λ_{760} nm (Spectrophotometer UV-Vis 1800, Shimadzu, Japan). The standard reference used was gallic acid ($y = 0.0004x + 0.0287$, $R^2 = 0.9877$), and the result was expressed as mg GAE (Gallic Acid Equivalent) per kg of dried noodles. TPC of samples (mg GAE/kg dried noodles) was calculated using the equation:

$$\text{TPC (mg GAE/kg dried noodles)} = \frac{As - 0.0287}{0.0004} \times \frac{2 \text{ mL}}{x \text{ g}} \times \frac{1 \text{ L}}{1000 \text{ mL}} \times \frac{1000 \text{ g}}{1 \text{ kg}}$$

where, As = absorbance of the samples, and x = weight of the dried noodles.

Total flavonoid content analysis

Total flavonoid content was analyzed using the modified method of Li et al.^[37]. The procedure began with mixing 0.3 mL of 5% NaNO_2 and 250 μ L of noodle extract in a 10 mL volumetric flask and incubating the mixture for 5 min. Afterward, 0.3 mL of 10% AlCl_3 was added to the volumetric flask. After 5 min, 2 mL of 1 M NaOH was added, and the volume was adjusted to 10 mL with distilled water. The sample was homogenized before analysis using a spectrophotometer (Spectrophotometer UV-Vis 1800, Shimadzu, Japan) at λ_{510} nm. The result was determined using a (+)-catechin standard reference ($y = 0.0008x + 0.0014$, $R^2 = 0.9999$) and expressed as mg CE (Catechin Equivalent) per kg of dried noodles. TFC of samples (mg CE/kg dried noodles) was calculated using the equation:

$$\text{TFC (mg CE/kg dried noodles)} = \frac{As - 0.0014}{0.0008} \times \frac{2 \text{ mL}}{x \text{ g}} \times \frac{1 \text{ L}}{1000 \text{ mL}} \times \frac{1000 \text{ g}}{1 \text{ kg}}$$

where, As = absorbance of the samples, and x = weight of the dried noodles.

Total anthocyanin content analysis

Total anthocyanin content was determined using the method of Giusti & Wrolstad^[38]. About 250 μ L of the sample was added with buffer solutions at pH 1 and pH 4.5 in different 10 mL test tubes. Then, each sample was mixed and incubated for 15 min and measured at λ_{543} and λ_{700} nm (Spectrophotometer UV-Vis 1800, Shimadzu, Japan). The absorbance (A) of samples was calculated with the formula: $A = (A_{\lambda_{543}} - A_{\lambda_{700}})_{\text{pH}1.0} - (A_{\lambda_{543}} - A_{\lambda_{700}})_{\text{pH}4.5}$. The total anthocyanin monomer content (TA) ($\text{mg} \cdot \text{mL}^{-1}$) was calculated with the formula: $\frac{A \times \text{MW} \times \text{DF} \times 1000}{\epsilon \times 1}$, where A was the absorbance of samples, MW was the molecular weight of delphinidin-3-glucoside (449.2 $\text{g} \cdot \text{mol}^{-1}$), DF was the factor of sample dilution, and ϵ was the absorptivity molar of delphinidin-3-glucoside (29,000 $\text{L} \cdot \text{cm}^{-1} \cdot \text{mol}^{-1}$).

TA monomer (mg delphinidine-3-glucoside/kg dried noodles)

$$= \text{TA (mg/L)} \times \frac{2 \text{ mL}}{x \text{ g}} \times \frac{1 \text{ L}}{1,000 \text{ mL}} \times \frac{1,000 \text{ g}}{1 \text{ kg}}$$

where, x = the weight of dried noodles.

2,2-Diphenyl-1-picrylhydrazyl free radical scavenging activity

DPPH analysis was measured based on the methods of Shirazi et al.^[39] and Widyawati et al.^[40]. Briefly, 10 μ L of the extract was added to a 10 mL test tube containing 3 mL of DPPH solution (4 mg DPPH in 100 mL methanol) and incubated for 15 min in a dark room. The solution was centrifuged at 5,000 rpm for 5 min, and the absorbance of samples was measured at λ_{517} nm (Spectrophotometer UV-Vis 1800, Shimadzu, Japan). The antioxidant activity of the samples was stated as an inhibition capacity with gallic acid as the standard reference ($y = 0.1405x + 2.4741$, $R^2 = 0.9974$) and expressed as mg GAE (Gallic Acid Equivalent) per kg of dried noodles. The percentage of DPPH free radical scavenging activity was calculated using the equation:

Inhibition of DPPH free radical scavenging activity

$$y (\%) = \frac{A_0 - A_s}{A_0} \times 100\%$$

where, A_0 = absorbance of the control and A_s = absorbance of the samples.

$$\text{DPPH free radical scavenging activity (mg GAE/kg dried noodles)} = \frac{y - 2.4741}{0.1405} \times \frac{2 \text{ mL}}{x \text{ g}} \times \frac{131 \text{ L}}{1000 \text{ mL}} \times \frac{1000 \text{ g}}{1 \text{ kg}}$$

where, x = the weight of dried noodles.

Ferric reducing antioxidant power

FRAP analysis was performed using the modified method of Al-Temimi & Choundhary^[41]. Approximately 50 μL of the extract in a test tube was added with 2.5 mL of phosphate buffer solution at pH 6.6 and 2.5 mL of 1% potassium ferric cyanide, shaken and incubated for 20 min at 50 °C. After incubation, the solution was added with 2.5 mL of 10% mono-chloroacetic acid and shaken until homogenized. Then, 2.5 mL of the supernatant was taken and added with 2.5 mL of bi-distilled water and 2.5 mL of 0.1% ferric chloride and incubated for 10 min. After incubation, samples were measured with absorbance at λ_{700} nm (Spectrophotometer UV-Vis 1800, Shimadzu, Japan). Gallic acid was used as the standard reference ($y = 2.2025x - 0.0144$, $R^2 = 0.9983$), and the results were expressed in mg GAE (Gallic Acid Equivalent) per kg of dried noodles. The reducing power of samples was calculated using the formula:

The reducing power (RP) (%) = $[(A_s - A_0)/A_s] \times 100\%$
Where, A_0 = absorbance of the control and A_s = absorbance of the samples.

$$\text{FRAP (mg GAE/kg dried noodles)} = \frac{\text{RP} + 0.0144}{2.2025} \times \frac{2 \text{ mL}}{x \text{ g}} \times \frac{131 \text{ L}}{1000 \text{ mL}} \times \frac{1000 \text{ g}}{1 \text{ kg}}$$

where, x = the weight of dried noodles.

Sensory evaluation

The sensory properties of cooked wet noodles were analyzed based on the methods of Nugroho et al.^[42] with modifications. The assessment used hedonic scale scoring with the parameters including color, aroma, taste, and texture attributes with 15 level, score 1 was stated as very much dislike, and 15 was very much like. This sensory analysis was performed by 100 untrained panelists between 17 and 25 years old who had previously gained knowledge of the measurement procedure. Each panelist was presented with 12 samples to be tested and given a questionnaire containing testing instructions and asked to give a score to each sample according to their level of liking. The hedonic scale used is a value of 1–15 given by panelists according to their level of liking for the product. A score of 1–3 indicates very much dislike, a score of 4–6 does not like, a score of 7–9 is neutral, a score of 10–12 likes it, and a score of 13–15 is very much like it. The best treatment was determined by the index effectiveness test^[43]. The best determination was based on sensory assay which included preferences for color, aroma, taste, and texture. The principle of testing was to give a weight of 0–1 on each parameter based on the level of importance of each parameter. The higher the weight value given means the parameter was increasingly prioritized. The treatment that has the highest value was determined as the best treatment. Procedure to determine the best treatment for wet noodles included:

(a) Calculation of the average of the weight parameters based on the results filled in by panelists

(b) Calculation of normal weight (BN)

BN = Variable weight/Total weight

(c) Calculation of effectiveness value (NE)

NE = Treatment value – worst value/Best value – worst value

(d) Calculation of yield value (NH)

NH = NE $\times$ normal weight

(d) Calculation of the total productivity value of all parameters

Total NH = NH of color + NH of texture + NH of taste + NH of aroma

(e) Determining the best treatment by choosing the appropriate treatment had the largest total NH

Design of experiment and statistical analysis

The design of the experiment used was a randomized block design (RBD) with two factors, i.e., the four ratios of the composite flour (wheat flour, stink lily flour, and κ -carrageenan) including 80:20:0 (K0); 80:19:1 (K1); 80:18:2 (K2), and 80:17:3 (K3) (% w/w), and the three-butterfly pea flower powder extracts, including 0 (T0), 15 (T1), 30 (T3) (% w/v). The experiment was performed in triplicate. The homogenous triplicate data were expressed as the mean $\pm$ SD. One-way analysis of variance (ANOVA) was carried out, and Duncan's New multiple range test (DMRT) was used to determine the differences between means ($p \leq 0.05$) using the statistical analysis applied SPSS 23.0 software (SPSS Inc., Chicago, IL, USA).

Results and discussions

Quality of wet noodles

The quality results of the wet noodles, including moisture content, water activity, tensile strength, swelling index, cooking loss, and color, are shown in Tables 2–5, and Fig. 1. Moisture content and water activity (A_w) of raw wet noodles were only significantly influenced by the various ratios of composite flour ($p \leq 0.05$) (Table 3). However, the interaction of the two factors, the ratios of composite flour and the concentrations of butterfly pea extract, or the concentrations of butterfly pea extract itself did not give any significant effects on the water content and A_w of wet noodles ($p \leq 0.05$) (Table 2). The K3 sample had the highest water content (70% wet base) compared to K0 (68% wet base), K1 (68% wet base), and K2 (69% wet base) because the sample had the highest ratio of κ -carrageenan. An increase of κ -carrageenan proportion influenced the amount of free and bound water in the wet noodle samples, which also increased the water content of the wet noodles. Water content resembles the amount of free and weakly bound water in the samples' pores, intermolecular, and intercellular space^[7,20]. Protein networking between gliadin and glutelin forms a three-dimensional networking structure of gluten involving water molecules^[44]. The glucomannan of stinky lily flour can form a secondary structure with sulfhydryl groups of the gluten network to stabilize it, increasing water binding capacity and retarding the migration of water molecules^[45]. κ -carrageenan can bind water molecules around 25–40 times^[46]. κ -carrageenan can cause a structure change in gluten protein through electrostatic interactions and hydrogen bonding^[47]. The interaction among the major proteins of wheat flour (gliadin and glutelin), glucomannan of stinky lily flour, and κ -carrageenan also changed the conformation of the three-dimensional network structure formation involving electrostatic forces, hydrogen bonds, and intra- and inter-molecular disulfide bonds that can establish water mobility in the dough of the wet noodles. The interaction of all components in the

Table 2. Quality properties of wet noodles at various ratios of composite flour and concentrations of butterfly pea flower extract.

Samples	Moisture content (% w/w)	Water activity	Swelling index (%)	Cooking loss (%)	Tensile strength (g)
K0T0	67.94 ± 0.11	0.975 ± 0.008	126.39 ± 2.06	18.91 ± 0.03	0.102 ± 0.008
K0T15	68.31 ± 0.07	0.976 ± 0.005	126.84 ± 1.69	19.02 ± 0.10	0.094 ± 0.003
K0T30	67.86 ± 0.66	0.978 ± 0.008	131.85 ± 2.97	19.76 ± 0.75	0.095 ± 0.003
K1T0	67.64 ± 0.27	0.971 ± 0.009	127.45 ± 7.15	18.71 ± 0.13	0.108 ± 0.007
K1T15	68.34 ± 0.44	0.973 ± 0.004	131.46 ± 0.93	18.77 ± 0.11	0.116 ± 0.011
K1T30	68.63 ± 1.08	0.969 ± 0.005	141.83 ± 8.15	19.32 ± 0.29	0.108 ± 0.008
K2T0	68.64 ± 0.52	0.974 ± 0.008	132.81 ± 3.77	18.26 ± 0.12	0.140 ± 0.002
K2T15	69.57 ± 0.59	0.973 ± 0.004	138.12 ± 1.18	18.43 ± 0.06	0.138 ± 0.006
K2T30	68.46 ± 0.68	0.962 ± 0.002	141.92 ± 8.23	18.76 ± 0.06	0.138 ± 0.013
K3T0	69.71 ± 0.95	0.969 ± 0.008	155.00 ± 4.16	17.54 ± 0.27	0.183 ± 0.002
K3T15	69.08 ± 0.38	0.973 ± 0.005	158.67 ± 7.28	18.03 ± 0.28	0.170 ± 0.011
K3T30	69.76 ± 0.80	0.970 ± 0.005	163.66 ± 7.52	18.33 ± 0.03	0.161 ± 0.002

No significant effect of interaction between composite flour and butterfly pea extract on quality properties of wet noodles. The results were presented as SD of means that were achieved in triplicate. All of the data showed that no interaction of the two parameters influenced the quality properties of wet noodles at $p \leq 0.05$.

Table 3. Effect of composite flour proportions on quality properties of wet noodles.

Samples	Moisture content (% w/w)	Water activity	Swelling index (%)	Cooking loss (%)	Tensile strength (g)
K0	68.04 ± 0.40 ^a	0.976 ± 0.01 ^b	128.36 ± 3.30 ^a	19.23 ± 0.55 ^d	0.097 ± 0.097 ^a
K1	68.20 ± 0.74 ^a	0.971 ± 0.01 ^a	133.58 ± 8.42 ^b	18.93 ± 0.34 ^c	0.112 ± 0.111 ^b
K2	68.89 ± 0.73 ^b	0.970 ± 0.01 ^a	137.62 ± 6.05 ^b	18.48 ± 0.23 ^b	0.141 ± 0.139 ^c
K3	69.52 ± 0.73 ^c	0.971 ± 0.01 ^a	159.11 ± 6.77 ^c	17.96 ± 0.40 ^a	0.173 ± 0.171 ^d

All of the data showed that there was a significant effect of composite flour on the quality properties of wet noodles at $p \leq 0.05$. The results were presented as SD of means that were achieved in triplicate. Means with different superscript letters in the same column are significantly different, $p \leq 0.05$.

Table 4. Effect of butterfly pea extract concentration on quality properties of wet noodles.


Samples	Moisture content (% w/w)	Water activity	Swelling index (%)	Cooking loss (%)	Tensile strength (g)
T0	68.48 ± 0.96	0.970 ± 0.010	135.41 ± 12.72 ^a	18.35 ± 0.57 ^a	0.134 ± 0.034 ^b
T15	68.67 ± 0.66	0.974 ± 0.000	138.77 ± 13.12 ^a	18.56 ± 0.41 ^a	0.130 ± 0.030 ^{a,b}
T30	68.83 ± 1.00	0.970 ± 0.010	144.82 ± 13.55 ^b	19.04 ± 0.67 ^b	0.129 ± 0.028 ^a

All of the data showed that there was a significant effect of butterfly pea extract concentration on the quality properties of wet noodles at $p \leq 0.05$. The results were presented as SD of means that were achieved in triplicate. Means with different superscript letters in the same column are significantly different, $p \leq 0.05$.

composite flour significantly influenced the amount of free water ($p \leq 0.05$) (Table 3). The addition of κ -carrageenan between 1%–3% in the wet noodle formulation reduced the A_w by about 0.005–0.006. The capability of κ -carrageenan to absorb water molecules reduces the water mobility in the wet noodles due to the involvement of hydroxyl, carbonyl, and

ester sulfate groups of them to form complex structures^[48]. The complexity of the reaction among components in the wet noodles to form a three-dimensional network influenced the amount of free water molecules that determined water activity values. The strength of the bonding among the components between wet noodles and water molecules also contributed to the value of the water activity.

Tensile strength, swelling index, and cooking loss of cooked wet noodles were significantly influenced by each factor of the ratios of composite flour or the concentrations of butterfly pea flower extract ($p \leq 0.05$) (Tables 3 & 4). However, the interaction between the two factors was not seen to influence the tensile strength, swelling index, and cooking loss of wet noodles ($p \leq 0.05$) (Table 2). An increase in the ratio of κ -carrageenan in the composite flour increased the tensile strength and swelling index and decreased the cooking loss of wet noodles. On the other hand, the increasing butterfly pea extract concentration decreased the tensile strength and increased the swelling index and cooking loss of wet noodles. Different ratios of the composite flour affected the tensile strength, which ranged between 0.197 and 0.171 g. At the same time, incorporating butterfly pea extract caused the tensile strength of wet noodles (T15 and T30) to significantly decrease from around 0.003 to 0.008 than control (K1). The highest and lowest swelling index

**Fig. 1** Color of wet noodles with various proportions of composite flour and concentrations of butterfly pea flower extract.

values were owned by K3 and K0 samples, respectively. The swelling index values of wet noodles ranged from 128% to 159%. The effect of the composite flour proportion of wet noodles showed that the K0 sample had the highest cooking loss, and the K3 sample possessed the lowest cooking loss. In contrast, the effect of the concentrations of butterfly pea extract resulted in the lowest cooking loss values of the T0 sample and the highest cooking loss values of the T30 sample. The cooking loss values of wet noodles ranged from 18% to 19%.

Tensile strength, cooking loss, and swelling index of wet noodles were significantly influenced by the interaction of components in dough formation, namely glutenin, gliadin, glucomannan, κ -carrageenan, and polyphenolic compounds, which resulted in a three-dimensional network structure that determined the capability of the noodle strands being resistant to breakage and gel formation. κ -carrageenan is a high molecular weight hydrophilic polysaccharide composed of a hydrophobic 3,6-anhydrous-D-galactose group and hydrophilic sulfate ester group linked by α -(1,3) and β -(1,4) glycosidic linkages^[49] that can bind water molecules to form a gel. Glucoman is a soluble fiber with the β -1,4 linkage main chain of D-glucose and D-mannose that can absorb water molecules around 200 times^[50] to form a strong gel that increases the viscosity and swelling index of the dough^[51]. Park & Baik^[52] stated that the gluten network formation affects the tensile strength of noodles. Huang et al.^[48] also reported that κ -carrageenan can increase the firmness and viscosity of samples because of this hydrocolloid's strong water-binding capacity. Cui et al.^[45] claimed that konjac glucomannan not only stabilizes the structure of gluten network but also reacts with free water molecules to form a more stable three-dimensional networking structure, thus maintaining dough's rheological and tensile properties.

The increased swelling index of dough is caused by the capability of glucomannan to reduce the pore size and increase the pore numbers with uniform size^[53]. The synergistic interaction between these hydrocolloids and gluten protein results in a stronger, more elastic, and stable gel because of the association and lining up of the mannan molecules into the junction zones of helices^[54]. The cross-linking and polymerization involving functional groups of gluten protein, κ -carrageenan, and glucomannan determined binding forces with each other.

The stronger attraction between molecules composed of cross-linking reduces the particles or molecules' loss during cooking^[54,55]. The stability of the network dimensional structure of the protein was influenced by the interaction of protein wheat, glucomannan, κ -carrageenan, and polyphenol compounds in the wet noodle dough that determined tensile strength, swelling index, and cooking loss of wet noodles. Schefer et al.^[19] & Widyawati et al.^[7] explained that phenolic compounds can disturb the interaction between the protein of wheat flour (glutelin and gliadin) and carbohydrate (amylose) to form a complex structure through many interactions, including hydrophobic, electrostatic, and Van der Waals interactions, hydrogen bonding, and π - π stacking. The phenolic compounds of butterfly pea extract interacted with κ -carrageenan, glucomannan, protein, or polysaccharide and influenced complex network structure. The phenolic compounds can disrupt the three-dimensional networking of interaction among gluten protein, κ -carrageenan, and glucomannan through aggregates or chemical breakdown of covalent and noncovalent bonds, and disruption of disulfide bridges to form thiols radicals^[55]. These compounds can form complexes with protein and hydrocolloids, leading to structural and functional changes and influencing gel formation through aggregation formation and disulfide bridge breakdown^[19,20,56].

The color of wet noodles (Table 5 & Fig. 1) was significantly influenced by the interaction between the composite flour and butterfly pea extract ($p \leq 0.05$). The L^* , a^* , b^* , C , and $^{\circ}h$ increased with increasing the composite flour ratio and the concentration of butterfly pea extract. Most of the color parameter values were lower than the control samples (K0T0, K1T0, K2T0, K3T0), except yellowness and chroma values of K2T0 and K3T0, whereas an increased amount of butterfly pea extract changed all color parameters. The L^* , a^* , b^* , C , and $^{\circ}h$ ranges were about 44 to 67, -13 to 1, -7 to 17, 10 to 16, and 86 to 211, respectively. Lightness, redness, and yellowness of wet noodles intensified with a higher κ -carrageenan proportion and diminished with increasing butterfly pea flower extract. The chroma and hue of wet noodles decreased with increasing κ -carrageenan proportion from T0 until T15 and then increased at T30. K2T30 treatment had the strongest blue color compared with the other treatments ($p \leq 0.05$). The presence of κ -carrageenan in composite flour also supported the water-holding capacity of wet noodles that influenced color. κ -carrageenan was

Table 5. Effect of interaction between composite flour and butterfly pea extract on wet noodle color.

Samples	L^*	a^*	b^*	C	$^{\circ}h$
K0T0	66.10 $\pm$ 0.30 ^f	0.90 $\pm$ 0.10 ^f	15.70 $\pm$ 0.10 ^f	15.70 $\pm$ 0.10 ^f	86.60 $\pm$ 0.20 ^a
K0T15	48.70 $\pm$ 0.20 ^c	-11.40 $\pm$ 0.30 ^{bc}	-3.50 $\pm$ 0.20 ^c	12.00 $\pm$ 0.30 ^c	197.00 $\pm$ 0.70 ^c
K0T30	44.00 $\pm$ 0.60 ^a	-12.80 $\pm$ 0.20 ^a	-6.50 $\pm$ 0.30 ^a	14.40 $\pm$ 0.20 ^e	206.90 $\pm$ 1.00 ^d
K1T0	67.10 $\pm$ 0.40 ^f	0.90 $\pm$ 0.20 ^f	15.80 $\pm$ 0.60 ^f	15.80 $\pm$ 0.60 ^f	86.60 $\pm$ 0.50 ^a
K1T15	51.50 $\pm$ 1.80 ^d	-10.80 $\pm$ 0.40 ^{cd}	-3.00 $\pm$ 0.20 ^{cd}	11.30 $\pm$ 0.40 ^{bc}	195.60 $\pm$ 0.60 ^c
K1T30	45.50 $\pm$ 0.20 ^b	-11.80 $\pm$ 0.80 ^b	-6.30 $\pm$ 0.30 ^a	13.40 $\pm$ 0.70 ^d	208.40 $\pm$ 2.30 ^d
K2T0	67.10 $\pm$ 0.20 ^f	1.00 $\pm$ 0.10 ^f	16.30 $\pm$ 0.10 ^g	16.30 $\pm$ 0.10 ^g	86.40 $\pm$ 0.10 ^a
K2T15	53.40 $\pm$ 0.30 ^e	-10.30 $\pm$ 0.80 ^{de}	-2.80 $\pm$ 0.10 ^d	10.70 $\pm$ 0.80 ^b	195.50 $\pm$ 1.30 ^c
K2T30	46.00 $\pm$ 0.40 ^b	-10.40 $\pm$ 0.20 ^{de}	-6.10 $\pm$ 0.40 ^a	12.10 $\pm$ 0.40 ^c	210.60 $\pm$ 1.30 ^e
K3T0	67.40 $\pm$ 0.30 ^f	1.20 $\pm$ 0.10 ^f	16.80 $\pm$ 0.70 ^g	16.90 $\pm$ 0.70 ^g	85.90 $\pm$ 0.20 ^a
K3T15	53.80 $\pm$ 1.30 ^e	-9.80 $\pm$ 0.70 ^e	-1.20 $\pm$ 0.20 ^e	9.90 $\pm$ 0.70 ^a	187.50 $\pm$ 1.10 ^b
K3T30	47.90 $\pm$ 0.70 ^c	-10.10 $\pm$ 0.40 ^{de}	-5.50 $\pm$ 0.30 ^b	11.60 $\pm$ 0.20 ^{bc}	208.40 $\pm$ 2.30 ^d

All of the data showed that there was a significant effect of interaction between composite flour and butterfly pea extract on the quality properties of wet noodles at $p \leq 0.05$. The results were presented as SD of means that were achieved in triplicate. Means with different superscript letters in the same column are significantly different, $p \leq 0.05$.

Composite flour-butterfly pea flower

synergized with glucomannan to produce a strong stable network that involved sulfhydryl groups. Tako & Konishi^[57] reported that κ -carrageenan can associate polymer structure that involves intra- and intermolecular interaction, such as ionic bonding and electrostatic forces. The mechanism of making a three-dimensional network structure that implicated all components of composite flour was exceptionally complicated due to the involved polar and non-polar functional groups and many kinds of interaction between them. These influenced the water content and water activity of the wet noodles, which impacted the wet noodle color. Another possible cause that affects wet noodles' color profile is anthocyanin pigment from the butterfly pea extract. Vidana Gamage et al.^[58] reported that the anthocyanin pigment of butterfly pea is delphinidin-3-glucoside and has a blue color. Increasing butterfly pea extract concentration lowered the lightness, redness, yellowness, and chroma and also changed the hue color from yellow to green-blue color.

The effect of composite flour and butterfly pea extract on color was observed in chroma and hue values. Glucomannan proportion of stinky lily flour intensified redness and yellowness, but butterfly pea extract reduced the two parameters. Thanh et al.^[59] also found similarities in their research. Anthocyanin pigment of butterfly pea extract can interact with the color of stinky lily and κ -carrageenan, impacting the color change of wet noodles. Thus, the sample T0 was yellow, T15 was green, and T30 was blue. Color intensity showed as chroma values of yellow values increased along with the higher proportion of κ -carrageenan at the same concentration of butterfly pea extract. However, the higher concentration of butterfly pea extract lessened the green and blue colors of wet noodles made using the same proportion of composite flour. Wet

noodle color is also estimated to be influenced by the phenolic compound content, which underwent polymerization or degradation during the heating process. Widyawati et al.^[20] reported that the bioactive compounds in pluchea extract could change the wet noodle color because of the discoloration of pigment during cooking. K2T30 was wet noodles exhibiting the strongest blue color due to different interactions between anthocyanin and hydrocolloid compounds, especially κ -carrageenan, that could reduce the intensity of blue color or chroma values.

Phenolic (TPC), total flavonoid (TFC), and total anthocyanin (TAC) content of wet noodles

The results of TPC, TFC, and TAC are shown in Table 6. The TPC and TFC of wet noodles were significantly influenced by the interaction between two parameters: the ratio of composite flour and the concentration of butterfly pea extracts ($p \leq 0.05$). The highest proportion of κ -carrageenan and butterfly pea extract resulted in the highest TPC and TFC. The K2T30 had the highest TPC and TFC of about ~207 mg GAE/kg dried noodles and ~57 mg CE/kg dried noodles, respectively. The TAC of wet noodles was only influenced by the concentration of butterfly pea extract, and the increase in extract addition led to an increase in TAC. The extract addition in T30 possessed a TAC of about 3.92 ± 0.18 mg delphinidin-3-glucoside/kg dried noodles. In addition, based on Pearson correlation assessment, there was a strong, positive correlation between the TPC of wet noodles and the TFC at T0 ($r = 0.955$), T15 ($r = 0.946$), and T30 treatments ($r = 0.765$). In contrast, a weak, positive correlation was observed between the TPC of samples and the TAC at T0 ($r = 0.153$) and T30 treatments ($r = 0.067$), except the T15 treatment, which had a correlation coefficient of -0.092 (Table 7).

Table 6. Effect of interaction between composite flour and butterfly pea extract on wet noodle's bioactive compounds and antioxidant activity.

Samples	TPC (mg GAE/kg dried noodles)	TFC (mg CE/kg dried noodles)	TAC (mg delphinidin-3-glucoside/kg dried noodles)	DPPH (mg GAE/kg dried noodles)	FRAP (mg GAE/kg dried noodles)
K0T0	126.07 $\pm$ 0.90 ^a	16.74 $\pm$ 6.26 ^a	0.00 $\pm$ 0.00 ^a	2.99 $\pm$ 0.16 ^a	0.009 $\pm$ 0.001 ^a
K0T15	172.57 $\pm$ 2.14 ^e	36.66 $\pm$ 2.84 ^d	2.67 $\pm$ 0.21 ^b	21.54 $\pm$ 1.71 ^d	0.023 $\pm$ 0.002 ^c
K0T30	178.07 $\pm$ 2.54 ^f	48.36 $\pm$ 3.29 ^f	3.94 $\pm$ 0.28 ^c	39.23 $\pm$ 0.91 ^f	0.027 $\pm$ 0.002 ^e
K1T0	137.07 $\pm$ 1.32 ^b	21.66 $\pm$ 3.67 ^b	0.00 $\pm$ 0.00 ^a	3.13 $\pm$ 0.19 ^a	0.011 $\pm$ 0.001 ^a
K1T15	178.48 $\pm$ 0.95 ^f	36.95 $\pm$ 3.05 ^d	2.74 $\pm$ 0.21 ^b	21.94 $\pm$ 0.68 ^d	0.023 $\pm$ 0.001 ^{cd}
K1T30	183.65 $\pm$ 1.67 ^g	52.28 $\pm$ 3.08 ^g	3.84 $\pm$ 0.19 ^c	41.42 $\pm$ 1.30 ^g	0.029 $\pm$ 0.001 ^f
K2T0	150.40 $\pm$ 0.52 ^d	27.49 $\pm$ 5.39 ^c	0.00 $\pm$ 0.00 ^a	7.45 $\pm$ 0.69 ^c	0.014 $\pm$ 0.001 ^b
K2T15	202.48 $\pm$ 0.63 ^j	48.28 $\pm$ 2.41 ^f	2.95 $\pm$ 0.57 ^b	24.70 $\pm$ 0.90 ^e	0.025 $\pm$ 0.001 ^d
K2T30	206.90 $\pm$ 2.43 ⁱ	56.99 $\pm$ 7.45 ^h	3.93 $\pm$ 0.42 ^c	47.55 $\pm$ 1.31 ⁱ	0.034 $\pm$ 0.002 ^g
K3T0	141.15 $\pm$ 1.28 ^c	25.37 $\pm$ 3.46 ^c	0.00 $\pm$ 0.00 ^a	5.45 $\pm$ 0.49 ^b	0.013 $\pm$ 0.001 ^b
K3T15	186.32 $\pm$ 1.15 ^h	43.57 $\pm$ 2.28 ^e	2.66 $\pm$ 0.21 ^b	22.45 $\pm$ 0.48 ^d	0.024 $\pm$ 0.001 ^{cd}
K3T30	189.90 $\pm$ 0.63 ^k	54.95 $\pm$ 3.72 ^{gh}	3.98 $\pm$ 0.37 ^c	44.93 $\pm$ 1.28 ^h	0.031 $\pm$ 0.001 ^f

All of the data showed that there was a significant effect of interaction between composite flour and butterfly pea extract to bioactive compounds and antioxidant activity of wet noodles at $p \leq 0.05$. The results were presented as SD of means that were achieved in triplicate. Means with different superscript letters in the same column are significantly different, $p \leq 0.05$.

Table 7. Pearson correlation coefficients between bioactive contents (TPC, TFC, and TAC) and antioxidant activity (DPPH and FRAP).

Parameter	TPC			TFC			TAC			DPPH		
	T0	T15	T30	T0	T15	T30	T0	T15	T30	T0	T15	T30
TPC	1	1	1									
TFC	0.955	0.946	0.765	1	1	1						
TAC	0.153	-0.092	0.067	0.028	-0.239	-0.020	1	1	1			
DPPH	0.893	0.815	0.883	0.739	0.753	0.753	0.123	0.127	0.194	1	1	1
FRAP	0.884	0.425	0.859	0.902	0.464	0.742	0.056	-0.122	-0.131	0.881	0.321	0.847

Correlation significant at the 0.05 level (2-tailed).

The bioactive compounds of wet noodles were correlated with their quality properties and antioxidant activity (AOA). The dominant anthocyanin pigment from butterfly pea extract is delphinidin^[60] around 2.41 mg/g samples^[61] that has more free acyl groups and aglycone structure^[62] that can be used as a natural pigment. The addition of butterfly pea extract influenced the color of wet noodles. Anthocyanin is a potential antioxidant agent through the free-radical scavenging pathway, cyclooxygenase pathway, nitrogen-activated protein kinase pathway, and inflammatory cytokines signaling^[63]. Nevertheless, butterfly pea extract is also composed of tannins, phenolics, flavonoids, phlobatannins, saponins, essential oils, triterpenoids, anthraquinones, phytosterols (campesterol, stigmasterol, β -sitosterol, and sitostanol), alkaloids, and flavonol glycosides (kaempferol, quercetin, myricetin, 6-malonylstragalgin, phenylalanine, coumaroyl sucrose, tryptophan, and coumaroyl glucose)^[12,13], chlorogenic, gallic, p-coumaric caffeic, ferulic, protocatechuic, p-hydroxy benzoic, vanillic, and syringic acids^[64], ternatin anthocyanins, fatty acids, tocopherols, mome inositol, pentanal, cyclohexene, 1-methyl-4(1-methylethylidene), and hirsutene^[64], that contribute to the antioxidant activity^[10,64]. *Clitoria ternatea* exhibits antioxidant activity based on the antioxidant assays, such as 2,2-diphenyl-1-picrylhydrazyl radical (DPPH) radical scavenging, ferric reducing antioxidant power (FRAP), hydroxyl radical scavenging activity (HRSA), hydrogen peroxide scavenging, oxygen radical absorbance capacity (ORAC), superoxide radical scavenging activity (SRSA), ferrous ion chelating power, 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) radical scavenging, and Cu^{2+} reducing power assays^[64]. The TPC and TFC of wet noodles increased along with the higher proportion of glucomannan in the composite flour and the higher concentration of butterfly pea extract. Zhou et al.^[65] claimed that glucomannan contained in stinky lily has hydroxyl groups that can react with Folin Ciocalteu's phenol reagent. Devaraj et al.^[66] reported that 3,5-acetylalbulin is a flavonoid compound in glucomannan that can form a complex with AlCl_3 .

Antioxidant activity of wet noodles

The antioxidant activity (AOA) of wet noodles was determined using DPPH radical scavenging activity (DPPH) and ferric reducing antioxidant power (FRAP), as shown in Table 6. The proportion of composite flour and the concentration of butterfly pea extracts significantly affected the DPPH results ($p \leq 0.05$). The noodles exhibited DPPH values ranging from 3 to 48 mg GAE/kg dried noodles. Several wet noodle samples, including the composite flour of K0 and K1 and without butterfly pea extracts (K0T0 and K1T0), had the lowest DPPH, while the samples containing composite flour K2 with butterfly pea extracts 30% (K2T30) had the highest DPPH. Pearson correlation showed that the TPC and TFC were strongly and positively correlated with the DPPH (Table 7). The correlated coefficient values (r) between TPC and AOA at T0, T15, and T30 treatments were 0.893, 0.815, and 0.883, respectively. Meanwhile, the r values between TFC and DPPH at T0, T15, and T30 treatments were 0.883, 0.739, and 0.753, respectively. However, the correlation coefficient values between TAC and AOA at T0, T15, and T30 treatments were 0.123, 0.127, and 0.194, respectively. The interaction among glucomannan, phenolic compounds, amylose, gliadin, and glutenin in the dough of wet noodles determined the number and position of free hydroxyl groups

that influenced the TPC, TFC, and DPPH. Widyawati et al.^[40] stated that free radical inhibition activity and chelating agent of phenolic compounds depends on the position of hydroxyl groups and the conjugated double bond of phenolic structures. The values of TPC, TFC, and DPPH significantly increased with higher levels of stinky lily flour and κ -carrageenan proportion and butterfly pea extract for up to 18% and 2% (w/w) of stinky lily flour and κ -carrageenan and 15% (w/w) of extract. However, the use of 17% and 3% (w/w) of stinky lily flour and κ -carrageenan and 30% (w/w) of the extract showed a significant decrease. The results show that the use of stinky lily flour and κ -carrageenan with a ratio of 17%:3% (w/w) was able to reduce free hydroxyl groups, which had the potential as electron or hydrogen donors in testing TPC, TFC, and DPPH.

FRAP of wet noodles was significantly influenced by the interaction of two parameters of the proportion of composite flour and the concentration of butterfly pea extracts ($p \leq 0.05$). FRAP was used to measure the capability of antioxidant compounds to reduce Fe^{3+} ions to Fe^{2+} ions. The FRAP capability of wet noodles was lower than DPPH, which ranged from 0.01 to 0.03 mg GAE/kg dried noodles. The noodles without κ -carrageenan and butterfly pea extracts (K0T0) had the lowest FRAP, while the samples containing composite flour K2 with 30% of butterfly pea extract (K2T30) had the highest FRAP. The Pearson correlation values showed that TPC and TFC at T0 and T30 treatments had strong and positive correlations to FRAP activity, but T15 treatment possessed a weak and positive correlation (Table 7). The correlation coefficient (r) values of TAC at the 0 treatment was weak with a positive correlation to FRAP samples, but the r values at T15 and T30 treatments showed weak negative correlations (Table 7). The obtained correlation between DPPH and FRAP activities elucidates that the DPPH method was highly correlated with the FRAP method at the T0 and T30 treatments and weakly correlated at the T15 treatment (Table 7). The DPPH and FRAP methods showed the capability of wet noodles to scavenge free radicals was higher than their ability to reduce ferric ion. It proved that the bioactive compounds of wet noodles have more potential as free radical scavengers or hydrogen donors than as electron donors. Compounds that have reducing power can act as primary and secondary antioxidants^[67]. Poli et al.^[68] stated that bioactive compounds with DPPH free radical scavenging activity are grouped as a primary antioxidant. Nevertheless, Suhendy et al.^[69] claimed that a secondary antioxidant is a natural antioxidant that can reduce ferric ion (FRAP). Based on AOA assay results, phenolic compounds indicated a strong and positive correlation with flavonoid compounds, as flavonoids are the major phenolic compounds potential as antioxidant agents through their ability to scavenge various free radicals. The effectivity of flavonoid compounds in inhibiting free radicals and chelating agents is influenced by the number and position of hydrogen groups and conjugated diene at A, B, and C rings^[70]. Previous studies have proven that TPC and TFC significantly contribute to scavenge free radicals^[71]. However, TAC showed a weak correlation with TFC, TPC, or AOA, although Choi et al.^[71] stated that TPC and anthocyanins have a significant and positive correlation with AOA, but anthocyanins insignificantly correlate with AOA. Different structure of anthocyanins in samples determines AOA. Moreover, the polymerization or complexation of anthocyanins with other molecules also determines their capability as electron or hydrogen donors.

Table 8. Effect of interaction between composite flour and butterfly pea extract to sensory properties of wet noodles and the best treatment of wet noodles based on index effectiveness test.

Samples	Color	Aroma	Taste	Texture	Index effectiveness test
K0T0	8.69 ± 3.31 ^a	7.41 ± 3.80 ^a	8.71 ± 3.16 ^a	10.78 ± 2.86 ^{abcde}	0.1597
K0T15	8.96 ± 3.38 ^b	7.75 ± 3.89 ^b	9.35 ± 3.36 ^{cde}	11.19 ± 3.10 ^{abcd}	0.6219
K0T30	8.93 ± 3.50 ^{bc}	7.71 ± 3.76 ^c	9.26 ± 3.17 ^{bcd}	11.13 ± 3.09 ^a	0.6691
K1T0	8.74 ± 3.62 ^a	8.13 ± 3.56 ^{ab}	9.58 ± 3.13 ^{ab}	11.33 ± 3.12 ^{de}	0.4339
K1T15	9.98 ± 3.06 ^{bc}	8.40 ± 3.28 ^c	10.16 ± 2.59 ^{def}	10.61 ± 2.82 ^{ab}	0.7086
K1T30	10.08 ± 3.28 ^{bc}	9.10 ± 3.08 ^c	10.44 ± 2.32 ^{bcd}	10.36 ± 2.81 ^{ab}	0.7389
K2T0	10.41 ± 3.01 ^a	9.39 ± 3.27 ^{ab}	11.04 ± 2.44 ^{ab}	10.55 ± 2.60 ^{cde}	0.3969
K2T15	10.8 ± 2.85 ^{bc}	9.26 ± 3.10 ^c	10.11 ± 2.76 ^f	10.89 ± 2.65 ^{abcd}	0.9219
K2T30	10.73 ± 3.02 ^c	9.10 ± 3.46 ^c	9.85 ± 2.99 ^{def}	10.16 ± 2.74 ^{abc}	0.9112
K3T0	10.73 ± 3.42 ^a	9.19 ± 3.38 ^b	9.93 ± 2.50 ^{bc}	10.34 ± 2.84 ^e	0.5249
K3T15	10.91 ± 3.23 ^{bc}	9.48 ± 3.56 ^c	10.45 ± 2.82 ^{cde}	10.49 ± 2.68 ^{bcde}	0.9235
K3T30	10.88 ± 3.14 ^c	9.49 ± 3.59 ^c	10.81 ± 2.74 ^{ef}	10.86 ± 2.60 ^{bcde}	1.0504

All of the data showed that there was a significant effect of interaction between composite flour and butterfly pea extract on the sensory properties of wet noodles at $p \leq 0.05$. The results were presented as SD of means that were achieved in triplicate. Means with different superscript letters in the same column are significantly different, $p \leq 0.05$.

Martin et al.^[72] informed that anthocyanins are the major groups of phenolic pigments where their antioxidant activity greatly depends on the steric hindrance of their chemical structure, such as the number and position of hydroxyl groups and the conjugated double bonds, as well as the presence of electrons in the structural ring. However, TPC and TFC at T0 and T30 treatments were highly and positively correlated with FRAP assay due to the role of phenolic compounds as reducing power agents that contributed to donating electrons. Paddayappa et al.^[67] reported that the phenolic compounds are capable of embroiling redox activities with an action as hydrogen donor and reducing agent. The weak relationship between TPC, TFC, or DPPH, and FRAP in the T15 treatment suggested that there was an interaction between the functional groups in the benzene ring in phenolic and flavonoid compounds and the functional groups in components in composite flour, thereby reducing the ability of phenolic and flavonoid compounds to donate electrons.

Sensory evaluation

Sensory properties of wet noodles based on the hedonic test results showed that composite flour and butterfly pea extract addition significantly influenced color, aroma, taste, and texture preferences ($p \leq 0.05$) (Table 8). The preference values of color, aroma, taste, and texture attributes of wet noodles ranged from 5 to 6, 6 to 7, 6 to 7, and 5 to 7, respectively. Incorporating butterfly pea extracts decreased preference values of color, aroma, taste, and texture attributes of wet noodles. Anthocyanin of butterfly pea extract gave different intensities of wet noodle color that resulted in color degradation from yellow, green, to blue color, impacting the color preference of wet noodles. Nugroho et al.^[42] also reported that the addition of butterfly pea extracts elevated the preference of panelists for dried noodles. The aroma of wet noodles was also affected by two parameters of treatments, where the results showed that the higher proportion of stinky lily caused the wet noodles to have a stronger, musty smell. Utami et al.^[73] claimed that oxalic acid in stinky lily flour contributes to the odor of rice paper. Therefore, a high proportion of κ -carrageenan could reduce the proportion of stinky lily flour, thereby increasing the panelists' preference for wet noodle aroma. Sumartini & Putri^[74] noted that panelists preferred noodles substituted with a higher

κ -carrageenan. Widyawati et al.^[7] also proved that κ -carrageenan is an odorless material that does not affect the aroma of wet noodles. Neda et al.^[63] added that volatile compounds of butterfly pea extract can mask the musty smell of stinky lily flour, such as pentanal and mome inositol. In addition, Padmawati et al.^[75] revealed that butterfly pea extract could give a sweet and sharp aroma. The panelists' taste preference for wet noodles without butterfly pea extract addition was caused by alkaloid compounds, i.e., conisin^[76] due to Maillard reaction during stinky lily flour processing. Nevertheless, using butterfly pea extract at a higher concentration in wet noodles increased the bitter taste, which is contributed by tannin compounds in this flower, as has been found by Handayani & Kumalasari^[77]. The effect of composite flour proportion and butterfly pea extract addition also appeared to affect the texture preference of wet noodles. Panelists preferred wet noodles that did not break up easily, which was the K3T0 sample, as the treatment resulted in chewy and elastic wet noodles. The results were also affected by the tensile strength of wet noodles because of the different concentrations of butterfly pea extract added to the wet noodles. The addition of butterfly pea extract at a higher concentration resulted in sticky, easy-to-break, and less chewy wet noodles^[18,19,77] due to the competition among phenolic compounds, glutelin, gliadin, amylose, glucomannan, κ -carrageenan to interact with water molecules to form gel^[78]. Based on the index effectiveness test, the noodles made with composite flour of K3 and butterfly pea extract of 30% (K3T30) were the best treatment, with a total score of 1.0504.

Conclusions

Using composite flour containing wheat flour, stinky lily flour, and κ -carrageenan and butterfly pea extract in wet noodle influenced wet noodles' quality, bioactive compounds, antioxidant activity, and sensory properties. Interaction among glutelin, gliadin, amylose, glucomannan, κ -carrageenan, and phenolic compounds affected the three-dimensional network structure that impacted moisture content, water activity, tensile strength, color, cooking loss, swelling index, bioactive content, antioxidant activity, and sensory properties of wet noodles. The higher concentration of hydrocolloid addition caused increased

water content and swelling index and decreased water activity and cooking loss. In addition, incorporating butterfly pea extract improved color, bioactive content, and antioxidant activity and enhanced panelists preference for wet noodles. Glucomannan of stinky lily flour and bioactive compounds of butterfly pea extract increased the functional value of the resulting wet noodles.

Author contributions

The authors confirm contribution to the paper as follows: study conception and design, literature search, methodologies of the lab analyses design: Widyawati PS, Suseno TIP; Fieldwork implementation, data collection, analysis and interpretation of results: Widyawati PS, Ivana F, Natania E; draft manuscript preparation: Widyawati PS; Manuscript revision: Wangtueai S. All authors reviewed the results and approved the final version of the manuscript.

Data availability

Data reported in this study are contained within the article. The underlying raw data are available on request from the corresponding author.

Acknowledgments

The authors would like to thank the Food Nutrition Research Center, Research institutions and community service, Widya Mandala Surabaya Catholic University for the research grant (7554/WM01/N/2022).

Conflict of interest

The authors declare that they have no conflict of interest.

Dates

Received 24 September 2023; Revised 14 January 2024; Accepted 29 January 2024; Published online 19 June 2024

References

- Siddeeg A, Salih ZA, Ammar A, Almahi RAY, Ali AO. 2018. Production of noodles from composite flour and it is nutrition sensory characteristics. *Chinese Journal of Medical Research* 1(1):3–7
- Efendi Z, Surawan FED, Sulastri Y. 2016. Physical properties of wet noodles based on potato and tapioca composite flour. *Jurnal Agroindustri* 6:57–64
- Dhull SB, Sandhu KS. 2018. Wheat-fenugreek composite flour noodles: effect on functional, pasting, cooking and sensory properties. *Current Research in Nutrition and Food Science* 6:174–82
- Park G, Cho H, Kim K, Kweon M. 2022. Quality characteristics and antioxidant activity of fresh noodles formulated with flour-bran blends varied by particle size and blend ratio of purple-colored wheat bran. *Processes* 10:584
- Zalewski BM, Chmielewska A, Szajewska H. 2015. The effect of glucomannan on body weight in overweight or obese children and adults: a systematic review of randomized controlled trials. *Nutrition* 31:437–442. e2
- Mahirdini S, Afifah DN. 2016. Effect of substitution of wheat flour with porang flour (*Amorphophallus oncophyllus*) on protein content, dietary fiber, fat, and acceptance rate of biscuits. *The Indonesian Journal of Nutrition* 5(1):42–49
- Widyawati PS, Suseno TIP, Widjajaseputra AI, Widyastuti TEW, Moeljadi VW, et al. 2022. The effect of κ -carrageenan proportion and hot water extract of the *Pluchea indica* less leaf tea on the quality and sensory properties of stink lily (*Amorphophallus muel-leri*) wet noodles. *Molecules* 27:5062
- Czajkowska-González YA, Alvarez-Parrilla E, del Rocio Martínez-Ruiz N, Vázquez-Flores AA, Gaytán-Martínez M, et al. 2021. Addition of phenolic compounds to bread: antioxidant benefits and impact on food structure and sensory characteristics. *Food Production, Processing and Nutrition* 3:25
- Saleh AA, Gawish E, Mahmoud SF, Amber K, Awad W, et al. 2021. Effect of natural and chemical colorant supplementation on performance, egg-quality characteristics, yolk fatty-acid profile, and blood constituents in laying hens. *Sustainability* 13:4503
- Purwanto UMS, Aprilia K, Sulistiyan. 2022. Antioxidant activity of telang (*Clitoria tematea* L.) extract in inhibiting lipid peroxidation. *Current Biochemistry* 9(1):26–37
- Jeyaraj EJ, Lim YY, Choo WS. 2022. Antioxidant, cytotoxic, and antibacterial activities of *Clitoria ternatea* flower extracts and anthocyanin-rich fraction. *Scientific Reports* 12:14890
- Pakeerathan K, Buddika H, Dharmadasa RM, Arawwala LADM. 2021. Phytochemical properties of (*Fabaceae*) distinct flower morphometric *Clitoria tematea* L. plants available in Sri Lanka. *The 1st International Electronic Conference on Agronomy session Sustainable Management Practices for Soil Health and Food Security*, 3-17 May 2021. Basel, Switzerland: MDPI. <https://sciforum.net/event/IECAG2021/submissions/view/4>
- Nabila FS, Radhityaningtyas D, Yurisna VC, Listyaningrum F, Aini N. 2022. Potency of butterfly pea flower (*Clitoria ternatea* L.) as an antibacterial in food products. *Journal of JIIPARI* 7(1):68–77
- Thuy NM, Ben TC, Ngoc PTB, Van Tai N. 2022. Application of butterfly pea flower extract in processing some Vietnamese traditional foods. *Journal of Applied Biology & Biotechnology* 10:143–50
- Singh R, Yu CC, Chen GW, Chen CH, Sinaki NY, et al. 2022. Butterfly pea flower as a novel ingredient to produce antioxidant-enriched yellow pea-based breakfast cereals. *Foods* 11:3447
- Oguis GK, Gilding EK, Jackson MA, Craik DJ. 2019. Butterfly pea (*Clitoria ternatea*), a cyclotide-bearing plant with applications in agriculture and medicine. *Frontiers in Plant Science* 10:645
- Shiau SY, Yu Y, Li J, Huang W, Feng H. 2023. Phytochemical-rich colored noodles fortified with an aqueous extract of *Clitoria ternatea* flowers. *Foods* 12:1686
- Zhu F. 2015. Interactions between starch and phenolic compound. *Trends in Food Science & Technology* 43:129–43
- Schefer S, Oest M, Rohn S. 2021. Interactions between phenolic acids, proteins, and carbohydrates-influence on dough and bread properties. *Foods* 10:2798
- Widyawati PS, Yulian DDLM, Utomo AR, Salim PEA, Martalia DE, et al. 2023. The effect of hot water extract of *Pluchea indica* leaf powder on the physical, chemical and sensory properties of wet noodles. *Current Research in Nutrition and Food Science Journal* 11:276–93
- Yuliani, Sari RA, Emmawati A, Candra KP. 2020. The shelf life of wet noodle added by *Gendarussa* (*Justicia gendarussa* burm. F.) leaves extract. *Indonesian Journal of Agricultural Research* 3:23–30
- Hasni D, Nilda C, Amalia R. 2022. The study of wet noodles making the high fiber using Konjac flour substitution and natural coloring. *Journal of Agricultural Technology & Products* 27(1):41–34 (In Bahasa Indonesia)
- Gustiani V, Putri RV, Yulinda R. 2022. Study of physicochemical and organoleptic properties of wet noodle with the addition of kelakai leaves extract (*Stenochlaena palustris*). *Transpublika International Research in Exact Sciences (TIREs)* 1(2):9–18
- Putri IM, Fakhri TM, Suarantika F. 2024. Literature study of antioxidant assay on butterfly pea flower extract (*Clitoria ternatea* L.) using various tested methods. *Bandung Conference Series Pharmacy* 4(1):40–49 (In Bahasa Indonesia)

25. Panjaitan TWS, Rosida DA, Widodo R. 2017. Aspects of the quality and level of consumer preference for wet noodle products with porang flour substitution. *Heuristics* 14:1–16 (In Bahasa Indonesia) <http://jurnal.untag-sby.ac.id/index.php/HEURISTIC/article/view/1040>
26. Zambrano MV, Dutta B, Mercer DG, Maclean HL, Touchie MF. 2019. Assessment of moisture content measurement methods of dried food products in small-scale operations in developing countries: A review. *Trends in Food Science and Technology* 88:484–96
27. Ribeiro M, Sousa T, Poeta P, Bagulho AS, Igrejas G. 2020. Review of structural features and binding capacity of polyphenols to gluten proteins and peptides in vitro: relevance to celiac disease. *Antioxidants* 9:463
28. Cauvain SP. 2012. *Breadmaking: improving quality*. 2nd Edition. Cambridge, United Kingdom: Woodhead Publishing. www.science-direct.com/book/9780857090607
29. Harijati N, Indriyani S, Mastuti R. 2013. Temperature effect of extraction on physicochemical properties glukomanan origin *Amorphophallus muelleri* blume. *NATURAL B* 2(2):128–33 (In Bahasa Indonesia)
30. Indrayati F, Utami R, Nurhartadi E. 2013. The addition effect of white turmeric (*Kaempferia rotunda*) rhizome essential oil on edible coating on colour and pH stability patin fish fillet during frozen storage. *Jurnal Teknosains Pangan* 2(4):25–31 (In Bahasa Indonesia) <https://jurnal.uns.ac.id/teknosains-pangan/article/view/4462/3808>
31. Sukarman, Astuti DA, Utomo NBP. 2017. Evaluation of color quality of Clown fish *Amphiprion percula* Lacepede 1802 natural catches and cultivated products. *Journal of Aquaculture Research* 12(3):231–39 (In Bahasa Indonesia)
32. Islamiya TY. 2015. *Characteristics of wet noodles with substitution of yellow corn flour and Moringa leaf flour (Moringa oleifera) as a functional food*. Bachelor's Thesis (In Bahasa Indonesia). Jember: Jember University. <http://repository.unej.ac.id/bitstream/handle/123456789/65440/Tri%20Yuli%20Islamiya%20-%20101710101058.pdf?sequence=1&isAllowed=y>
33. Tan HZ, Li ZG, Tan B. 2009. Starch noodles: History, classification, materials, processing, structure, nutrition, quality evaluating and improving. *Food Research International* 42(5-6):551–76
34. Aditia RP, Munandar A, Surilayani D, Haryati S, Sumantri MH, et al. 2021. Characteristics of dry noodles with seaweed flour *Gracilaria* spp. substitution. *LEUIT Journal of Local Food Security* 2(1):83–90
35. Chen Z, Schols HA, Voragen AGJ. 2003. Starch granule size strongly determines starch noodle processing and noodle quality. *Journal of Food Science* 68:1584–89
36. Ayele DT, Akele ML, Melese AT. 2022. Analysis of total phenolic contents, flavonoids, antioxidant and antibacterial activities of *Croton macrostachyus* root extracts. *BMC Chemistry* 16:30
37. Li G, Yu S, Zhou YH, Chen QF. 2013. Spectrophotometric determination of flavonoids content in leaves of *Fagopyrum cymosum* complex. *Asian Journal of Chemistry* 25(13):7575–78
38. Giusti MM, Wrolstad RE. 2001. Characterization and measurement of anthocyanins by UV-Visible spectroscopy. *Current Protocols in Food Analytical Chemistry* 00(1):F1.2.1–F1.2.13
39. Shirazi OU, Khattak MMAK, Shukri NAM, Anuar MNN. 2014. Determination of total phenolic, flavonoid content, and free radical scavenging activities of common herbs and spices. *Journal of Pharmacognosy and Phytochemistry* 3(3):104–8 www.phytojournal.com/archives/2014/vol3issue3/PartB/29.1.pdf
40. Widyawati PS, Budianta TDW, Kusuma FA, Wijaya EL. 2014. Difference of solvent polarity to phytochemical content and antioxidant activity of *Pluchea indicia* Less leaves extracts. *International Journal of Pharmacy and Pharmaceutical Research* 6(4):850–55 http://impactfactor.org/PDF/IJPPR/6/IJPPR_Vol6_Issue4_Article_29.pdf
41. Al-Temimi A, Choundary R. 2013. Determination of antioxidant activity in different kinds of plants *in vivo* and *in vitro* by using diverse technical methods. *Journal of Nutrition & Food Sciences* 3:184
42. Nugroho WT, Kurnianto MF, Wibowo MJ, Brilliantina A, Hariono B. Chemical and sensory characteristics of dried noodles with addition of telang flower extract (*Clitoria ternatea* L). *The 3rd International Conference on Food and Agriculture*. 3(1):96-102. <https://proceedings.polije.ac.id/index.php/food-science/article/view/182>
43. DeGarmo EP, Sullivan WG, Candra CR. 1984. *Engineering Economy*. 7th Edition. London: Mc Millan Publ. Co
44. Ooms N, Delcour JA. 2019. How to impact gluten protein network formation during wheat flour dough making. *Current Opinion in Food Science* 25:88–97
45. Cui T, Wu T, Liu R, Sui W, Wang S, et al. 2019. Effect of degree of konjac glucomannan enzymatic hydrolysis on the physicochemical characteristic of gluten and dough. *ACS Omega* 4:9654–63
46. Schilling MW. 2019. Emulsifier applications in meat products. In *Food emulsifiers and their applications*, eds. Hasenhuettl G, Hartel R. Cham: Springer. pp. 347–77. https://doi.org/10.1007/978-3-030-29187-7_12
47. Li J, Yadav MP, Li J. 2019. Effect of different hydrocolloids on gluten proteins, starch and dough microstructure. *Journal of Cereal Science* 87:85–90
48. Huang M, Theng AHP, Yang D, Yang H. 2021. Influence of κ -carrageenan on the rheological behaviour of a model cake flour system. *LWT* 136:110324
49. Geonzon LC, Bacabac RG, Matsukawa S. 2019. Network structure and gelation mechanism of kappa and iota carrageenan elucidated by multiple particle tracking. *Food Hydrocolloids* 92:173–80
50. Supriyati Y. 2016. The diversity of iles-iles (*Amorphophallus* spp.) and its potential for the functional food, cosmetics, and bioethanol industries. *Agricultural Litbang Journal* 35(2):69–80
51. Dewi PS, Ulandari D, Susanto NS. 2021. Effect of glucomannan addition on physical and sensory characteristic of gluten-free muffin from modified cassava flour and maize flour. *IOP Conference Series: Earth and Environmental Science* 733:012085
52. Park SJ, Baik BK. 2009. Quantitative and qualitative role of added gluten on white salted noodles. *Cereal Chemistry* 86:646–52
53. Sutrisno A, Yuwono SS, Ikarini I. 2021. Effect of glucomannan and xanthan gum proportion on the physical and sensory characteristic of gluten-free bread. *IOP Conference Series: Earth and Environmental Science* 924:012028
54. Chen Y, Song C, Lv Y, Qian X. 2019. Konjac glucomannan/kappa carrageenan interpenetrating network hydrogels with enhanced mechanical strength and excellent self-healing capability. *Polymer* 184:121913
55. Labat E, Morel MH, Rouau X. 2000. Effects of laccase and ferulic acid on wheat flour doughs. *Cereal Chemistry* 77:823–28
56. Amoako D, Awika JM. 2016. Polyphenol interaction with food carbohydrates and consequences on availability of dietary glucose. *Current Opinion in Food Science* 8:14–18
57. Tako M, Konishi T. 2018. Discovery of κ -carrageenan-like agarose from a red seaweed, *Gracilaria coronopifolia*. *International Research Journal of Pure and Applied Chemistry* 17:1–11
58. Vidana Gamage GC, Lim YY, Choo WS. 2021. Anthocyanins from *Clitoria ternatea* flower: biosynthesis, extraction, stability, antioxidant activity, and applications. *Frontiers in Plant Science* 12:792303
59. Thanh VT, Tran NYT, Linh NTV, Vy TA, Truc TT. 2020. Application of anthocyanin natural colors from Butterfly Pea (*Clitoria ternatea* L.) extracts to cupcake. *IOP Conference Series: Materials Science and Engineering* 736:062014
60. Campbell SM, Pearson B, Marble SC. 2019. Butterfly pea (*Clitoria ternatea*) flower extract (BPFE) and its use as a pH-dependent natural colorant. *EDIS* 2019:1–5
61. Shen Y, Du L, Zeng H, Zhang X, Prinyawiwatkul W, et al. 2016. Butterfly pea (*Clitoria ternatea*) seed and petal extracts decreased HEP-2 carcinoma cell viability. *International Journal of Food Science & Technology* 51:1860–68

62. Marpaung AM. 2020. Review of the benefits of butterfly pea flowers (*Clitoria tematea* L.) for human health. *Journal of Functional Food and Nutraceutical* 1(2):47–69 (In Bahasa Indonesia)
63. Neda GD, Rabeta MS, Ong MT. 2013. Chemical composition and anti-proliferative properties of flowers of *Clitoria Ternatea*. *International Food Research Journal* 20(3):1229–34
64. Jeyaraj EJ, Lim YY, Choo WS. 2021. Extraction methods of butterfly pea (*Clitoria tematea*) flower and biological activities of its phytochemicals. *Journal of Food Science and Technology* 58:2054–2067
65. Zhou Y, Zhao D, Foster TJ, Liu Y, Wang Y, et al. 2014. Konjac glucomannan-induced changes in thiol/disulphide exchange and gluten conformation upon dough mixing. *Food Chemistry* 143:163–69
66. Devaraj RD, Reddy CK, Xu B. 2019. Health-promoting effects of konjac glucomannan and its practical applications: a critical review. *International Journal of Biological Macromolecules* 126:273–81
67. Padayappa M, Senthilkumar A, Moorthi M, Thangaraj A. 2021. Analysis of phytochemical properties, DPPH and FRAP assay of antioxidant activities of *Acalypha indica* L. *International Journal of Scientific Research* 9(12):31–34
68. Poli AR, Katja DG, Aritonang HF. 2022. Potency of antioxidant extract from matoa seed coat (*Pometia pinnata* J. R. & G. Forst). *Chemistry Progress* 15(1):25–30 (In Bahasa Indonesia) <https://ejournal.unsrat.ac.id/v3/index.php/chemprog/article/view/43151/37970>
69. Suhendy H, Alif A, Rahmiyanti I. 2022. Correlation of total phenolic and flavonoid contents to the antioxidant activity of several African leaf extracts (*Vernonia amygdalina* Delile) using the FRAP method (Ferric Reducing Antioxidant Power). *Medical Science: Pharmaceutical Scientific Journal* 7(2):225–36 (In Bahasa Indonesia)
70. Mutha RE, Tatiya AU, Surana SJ. 2021. Flavonoids as natural phenolic compounds and their role in therapeutics: an overview. *Future Journal of Pharmaceutical Sciences* 7:25
71. Choi SC, Kim JM, Lee YG, Kim C. 2019. Antioxidant activity and contents of total phenolic compounds and anthocyanins according to grain colour in several varieties of *Sorghum bicolor* (L.) Moench. *Cereal Research Communications* 47:228–38
72. Martin J, Kuskoski EM, Navas MJ, Asuero AG. 2017. Antioxidant Capacity of anthocyanin pigments. In *Flavonoids - From Biosynthesis to Human Health*, ed. Justino GC. London: Intech Open Limited. <https://doi.org/10.5772/67718>
73. Utami R, Setiawati L, Rahmawati R. 2021. Characteristics of rice paper formulated with suweg flour (*Amorphophallus campanulatus*). *Conversion Journal* 10(2):19–28 (In Bahasa Indonesia) <https://jurnal.umj.ac.id/index.php/konversi/article/view/11954>
74. Sumartini, Putri WR. 2022. Effect of additional carrageenan concentration on the characteristics of wetnoodles based on mangrove fruit flour variation. *Food Research* 6:174–83
75. Padmawati IGA, Pratiwi IDPK, Wiadnyani AAIS. 2022. Effect of adding butterfly pea flower extract (*Clitoria ternatea* Linn) on marshmallow characteristics. *Itepa:Journal of Food Science and Technology* 11(1):43–54 (In Bahasa Indonesia)
76. Yasin S, Padusung P, Mahrup M, Kusnara I, Sukartono S, Fahrudin F. 2021. Exploring the potential of porang plants as community forest cultivation plants (HKM) in North Lombok Regency. *Journal of Master of Science Education Service* 4(2):316–27 (In Bahasa Indonesia)
77. Handayani NE, Kumalasari ID. 2021. Microbiological and sensory analysis of wet noodles formulated using butterfly pea flower extract (*Clitoria ternatea* L.) as a natural preservative and antioxidant. *Agrointek* 16(2):153–63 (In Bahasa Indonesia)
78. Sievert D, Hosney RC, Delcours JA. 2007. Bread and Other Baked Products. In *Ullmann's Food and Feed*. Vol 1. New York: Wiley. pp 493. https://doi.org/10.1002/14356007.a04_331.pub2



Copyright: © 2024 by the author(s). Published by Maximum Academic Press, Fayetteville, GA. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.

Effect of butterfly pea (Clitoria ternatea) flower extract on qualities, sensory properties, and antioxidant activity of wet noodles with various composite flour proportions

ORIGINALITY REPORT

8%

SIMILARITY INDEX

5%

INTERNET SOURCES

6%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Adhi SUSILO, Eko Yuliastuti Endah SULISTYAWATI, Dini Nur HAKIKI, Hari HARIADI et al. "Physicochemical properties of snackbar canna edulis ker with additional butterfly pea extract as funtional food", Food Science and Technology, 2023

Publication

<1 %
- 2 Zheng-Hao Wan, Chun-Lei Wu, Si-Yuan Tong, Shi-Yu Qiu. "Fusing Deep Separable Networks for Emotion Recognition in Complex Environments", Springer Science and Business Media LLC, 2024

Publication

<1 %
- 3 A Nurfirdausi, N Aini, H Dwiyanti. "Characteristics of gluten-free cookies from konjac, jack beans and dragon fruit (Hylocereus undatus)", IOP Conference Series: Earth and Environmental Science, 2023

Publication

<1 %

4

Franziska M. Jürgens, Sara M. Robledo, Thomas J. Schmidt. "Evaluation of Pharmacokinetic and Toxicological Parameters of Arnica Tincture after Dermal Application In Vivo", Pharmaceutics, 2022

Publication

<1 %

5

M. Morales-de la Peña, L. Salvia-Trujillo, M.A. Rojas-Graü, O. Martín-Belloso. "Changes on phenolic and carotenoid composition of high intensity pulsed electric field and thermally treated fruit juice-soymilk beverages during refrigerated storage", Food Chemistry, 2011

Publication

<1 %

6

Submitted to Politeknik Negeri Medan

Student Paper

<1 %

7

www.ajol.info

Internet Source

<1 %

8

Tuti Sri Suhesti, Warsinah Warsinah, Pitra Wulandari. "Optimizing effervescent granules of butterfly pea (Clitoria ternatea L) flower ethanol extract as antioxidant", Acta Pharmaciae Indonesia : Acta Pharm Indo, 2022

Publication

<1 %

9

evols.library.manoa.hawaii.edu

Internet Source

<1 %

10	Dilip Ghosh, Pulok K. Mukherjee. "Natural Medicines - Clinical Efficacy, Safety and Quality", CRC Press, 2019 Publication	<1 %
11	library.wur.nl Internet Source	<1 %
12	Hubert, J.. "Effects of fermentation on the phytochemical composition and antioxidant properties of soy germ", Food Chemistry, 20080815 Publication	<1 %
13	back-edupack.santillana.es Internet Source	<1 %
14	publicatio.bibl.u-szeged.hu Internet Source	<1 %
15	www.coursehero.com Internet Source	<1 %
16	www.grafiati.com Internet Source	<1 %
17	Submitted to Management Development Institute Of Singapore Student Paper	<1 %
18	dspace.ucuenca.edu.ec Internet Source	<1 %
19	Submitted to Gitam University Student Paper	

<1 %

20

Submitted to National University of Singapore

Student Paper

<1 %

21

Skowyra, Monika, M. Isabel Calvo, M. Gabriela Gallego, Nurul A. M. Azman, and Maria P. Almajano. "Characterization of Phytochemicals in Petals of Different Colours From Viola × wittrockiana Gams. and Their Correlation With Antioxidant Activity", Journal of Agricultural Science, 2014.

Publication

<1 %

22

api.repository.cam.ac.uk

Internet Source

<1 %

23

Achmad Qodim Syafa'atullah, Arie Amira, Sonya Hidayati, Mahfud Mahfud.

"Anthocyanin from butterfly pea flowers (Clitoria ternatea) by ultrasonic-assisted extraction", AIP Publishing, 2020

Publication

<1 %

24

Megha Kumari, Prajya Arya, Sunil Khatkar, Pradyuman Kumar. "Development and characterization of apple pomace and finger millet-based pasta enriched with encapsulated micronutrient", Food Chemistry Advances, 2023

Publication

<1 %

25	Shuai Sun, Peichen Wu, Fuquan Gao, Xiaomin Yu, Ying Liu, Chao Zheng. "Genome-wide identification and expression analysis of phytochrome-interacting factor genes during abiotic stress responses and secondary metabolism in the tea plant", Plant Physiology and Biochemistry, 2024 Publication	<1 %
26	Wandee, Yuree, Dudsadee Uttapap, Santhanee Puncha-arnon, Chureerat Puttanlek, Vilai Rungsardthong, and Nuanchawee Wetprasit. "Quality assessment of noodles made from blends of rice flour and canna starch", Food Chemistry, 2015. Publication	<1 %
27	digilib.unimed.ac.id Internet Source	<1 %
28	journal2.uad.ac.id Internet Source	<1 %
29	public.pensoft.net Internet Source	<1 %
30	Yadav, B.S.. "Suitability of pigeon pea and rice starches and their blends for noodle making", LWT - Food Science and Technology, 201107 Publication	<1 %
31	thanasworldmarket.com Internet Source	<1 %

33 Dharam P. Abrol. "The Future Role of Dwarf Honeybees in Natural and Agricultural Systems", CRC Press, 2020

Publication

<1 %

34 J. K. Abhishek, C. S. Ravi, N. Sudharani, N. R. Nagaraja. "Genetic diversity analysis for nutraceuticals and antioxidant activity in selected Indian pennywort (*Centella asiatica* L.) accessions of hill zone of Karnataka", Genetic Resources and Crop Evolution, 2024

Publication

<1 %

35 Ke Xiong, Meng-meng Li, Yi-qiang Chen, Yumeng Hu, Wen Jin. "Formation and reduction of toxic compounds derived from the Maillard Reaction during the thermal processing of different food matrices", Journal of Food Protection, 2024

Publication

<1 %

36 Mohammad Afzal Hossain, Pappu Dey, Rahman Ishtiaque Joy. "Effect of Osmotic Pretreatment and Drying Temperature on Drying Kinetics, Antioxidant Activity, and Overall Quality of Taikor (*Garcinia pedunculata* Roxb.) Slices", Saudi Journal of Biological Sciences, 2021

Publication

<1 %

37	psasir.upm.edu.my Internet Source	<1 %
38	researchcommons.waikato.ac.nz Internet Source	<1 %
39	scidar.kg.ac.rs Internet Source	<1 %
40	tsiantific.wordpress.com Internet Source	<1 %
41	worldresearchersassociations.com Internet Source	<1 %
42	Anita Sønsteby, Arnfinn Nes, Finn Måge. "Effects of bark mulch and npk fertilizer on yield, leaf nutrient status and soil mineral nitrogen during three years of strawberry production", Acta Agriculturae Scandinavica, Section B - Soil & Plant Science, 2004 Publication	<1 %
43	Arshad Mehmood, Muhammad Ishaq, Lei Zhao, Sanabil Yaqoob, Bushra Safdar, Muhammad Nadeem, Masooma Munir, Chengtao Wang. "Impact of ultrasound and conventional extraction techniques on bioactive compounds and biological activities of blue butterfly pea flower (<i>Clitoria ternatea</i> L.)", Ultrasonics Sonochemistry, 2019 Publication	<1 %

44 Chirinos, R.. "Optimization of extraction conditions of antioxidant phenolic compounds from mashua (*Tropaeolum tuberosum* Ruiz & Pavon) tubers", Separation and Purification Technology, 20070615 $<1\%$

45 Cristina M. Rosell, Peter Koehler. "Chapter 3 Chemistry of Cereal Grains", Springer Science and Business Media LLC, 2023 $<1\%$

46 Nair, Vimal, Woo Young Bang, Elisa Schreckinger, Nuri Andarwulan, and Luis Cisneros-Zevallos. "The protective role of Ternatin Anthocyanins and Quercetin Glycosides from Butterfly Pea (*Clitoria ternatea* Leguminosae) Blue flower Petals against LPS-induced inflammation in Macrophage Cells", Journal of Agricultural and Food Chemistry $<1\%$

47 Yanwei Chai, Mingzhu Wang, Genyi Zhang. "Interaction between Amylose and Tea Polyphenols Modulates the Postprandial Glycemic Response to High-Amylose Maize Starch", Journal of Agricultural and Food Chemistry, 2013 $<1\%$

48

Internet Source

<1 %

49

discovery.researcher.life

Internet Source

<1 %

50

edoc.site

Internet Source

<1 %

51

efsa.onlinelibrary.wiley.com

Internet Source

<1 %

52

garj.org

Internet Source

<1 %

53

mafiadoc.com

Internet Source

<1 %

54

perpustakaan.poltekkes-malang.ac.id

Internet Source

<1 %

55

repository.ut.ac.id

Internet Source

<1 %

56

trepo.tuni.fi

Internet Source

<1 %

57

vdocument.in

Internet Source

<1 %

58

www.jd.id

Internet Source

<1 %

59

www.medicaljournals.se

Internet Source

<1 %

60

www.nutriweb.org.my

Internet Source

<1 %

61

www.repository.wima.ac.id

Internet Source

<1 %

62

www.spandidos-publications.com

Internet Source

<1 %

63

Se-Kwon Kim. "Seafood Science - Advances in Chemistry, Technology and Applications", CRC Press, 2019

Publication

<1 %

64

Anelise S. Nazari, Suziane A. Dias, Willian F. da Costa, Ciomar A. Bersani-Amado et al. " Anti-inflammatory and Antioxidant Activities of . and Major Constituents ", Pharmaceutical Biology, 2008

Publication

<1 %

65

Is Fatimah, Habibi Hidayat, Bambang Hernawan Nugroho, Saddam Husein. "Ultrasound-assisted biosynthesis of Silver and Gold Nanoparticles using Clitoria ternatea flower", South African Journal of Chemical Engineering, 2020

Publication

<1 %

66

Jose M. Alvarez-Suarez, Sara Tulipani, Daimy Díaz, Yadiley Estevez et al. "Antioxidant and antimicrobial capacity of several monofloral Cuban honeys and their correlation with color,

<1 %

polyphenol content and other chemical compounds", Food and Chemical Toxicology, 2010

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Effect of butterfly pea (Clitoria ternatea) flower extract on qualities, sensory properties, and antioxidant activity of wet noodles with various composite flour proportions

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12