- 1. Submitted to the Journal RSC Advances (9-2-2023)
- 2. First revision (major revision) (10-3-2023)
- 3. Revised version submitted (15-3-2023)
- 4. Paper accepted (17-3-2023)
- 5. Additional revision from editor (Live subject statement) (19-3-2023)
- 6. Revised version submitted (20-3-2023)
- 7. Permition to sign a license to publish (21-3-2023)
- 8. Sign accepted (21-3-2023)
- 9. Proof reading (22-3-2023)
- 10.Corrections received (24-3-2023)
- 11.Article published (28-3-2023)

1.Submitted to the Journal RSC Advances (9-2-2023)



Acknowledgement of your Submission to RSC Advances - RA-ART-02-2023-000868

1 message

RSC Advances <onbehalfof@manuscriptcentral.com> Reply-To: advances@rsc.org To: antonius.budiawan@ukwms.ac.id Cc: antonius.budiawan@ukwms.ac.id, levi.puradewa@ukwms.ac.id, erlien.dwi.cahyani@ukwms.ac.id Thu, Feb 9, 2023 at 1:19 PM

09-Feb-2023

Dear Mr Budiawan:

TITLE: Wound Healing Activity and Flavonoids Contents of Purslane (Portulaca grandiflora) Various Variety

Thank you for your submission to RSC Advances, published by the Royal Society of Chemistry. This is an automatic acknowledgement that you have uploaded your files to our online submission system. Your manuscript ID is: RA-ART-02-2023-000868

Your manuscript will be passed to an editor for initial assessment as soon as possible. If there are any problems with your submission we will contact you.

All RSC Advances articles are published under an open access model, and the appropriate article processing charge (APC) will apply. Details of the APC and discounted rates can be found at https://www.rsc.org/journals-books-databases/about-journals/rsc-advances/# charges-RSC-Advances

Please indicate the above manuscript ID when you contact us about this submission.

Do you have an ORCID iD? ORCID (Open Researcher and Contributor iD) is a unique researcher identifier that allows you to link your research output and other professional activities in a single record. We therefore encourage each researcher to sign up for their own ORCID account. Please edit your user account to link your ORCID iD or create a new one, ensuring that you have not linked your account to another researcher's ORCID iD. Please note that we are unable to do this on your behalf. If your article is accepted, you may choose to have your ORCID record updated automatically with details of the publication.

We already have the following information for authors of this manuscript: Budiawan, Antonius - https://orcid.org/0000-0003-3730-3913, Purwanto, Agus - No ORCID iD Available, Puradewa, Levi - No ORCID iD Available, Cahyani, Erlien Dwi - No ORCID iD Available, Purwaningsih, Christiana Endang - No ORCID iD Available If this is not how you want your name to appear on an Accepted Manuscript, please amend your ScholarOne account.

The Royal Society of Chemistry is a member of CrossCheck. Your submission may be compared against the CrossCheck database using the iThenticate plagiarism detection software. For further information, please see here: https://www.rsc.org/journals-books-databases/journal-authors-reviewers/processes-policies/#crosscheck

This journal has a policy of sharing reviewer reports. Please note that the Editor's decision and copies of the reports will be shared with all reviewers who provide a recommendation on your manuscript.

Please contact us if we can be of any assistance.

Yours sincerely, RSC Advances Editorial Office advances@rsc.org If you need to contact the journal, please use the email address advances@rsc.org

DISCLAIMER:

This communication is from The Royal Society of Chemistry, a company incorporated in England by Royal Charter (registered number RC000524) and a charity registered in England and Wales (charity number 207890). Registered office: Burlington House, Piccadilly, London W1J 0BA. Telephone: +44 (0) 20 7437 8656.

The content of this communication (including any attachments) is confidential, and may be privileged or contain copyright material. It may not be relied upon or disclosed to any person other than the intended recipient(s) without the consent of The Royal Society of Chemistry. If you are not the intended recipient(s), please (1) notify us immediately by replying to this email, (2) delete all copies from your system, and (3) note that disclosure, distribution, copying or use of this communication is strictly prohibited.

Any advice given by The Royal Society of Chemistry has been carefully formulated but is based on the information available to it. The Royal Society of Chemistry cannot be held responsible for accuracy or completeness of this communication or any attachment. Any

views or opinions presented in this email are solely those of the author and do not represent those of The Royal Society of Chemistry. The views expressed in this communication are personal to the sender and unless specifically stated, this e-mail does not constitute any part of an offer or contract. The Royal Society of Chemistry shall not be liable for any resulting damage or loss as a result of the use of this email and/or attachments, or for the consequences of any actions taken on the basis of the information provided. The Royal Society of Chemistry does not warrant that its emails or attachments are Virus-free; The Royal Society of Chemistry has taken reasonable precautions to ensure that no viruses are contained in this email, but does not accept any responsibility once this email has been transmitted. Please rely on your own screening of electronic communication.

More information on The Royal Society of Chemistry can be found on our website: www.rsc.org



Wound Healing Activity and Flavonoids Contents of Purslane (Portulaca grandiflora) Various Variety

Journal:	RSC Advances
Manuscript ID	Draft
Article Type:	Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Budiawan, Antonius; Widya Mandala Catholic University Surabaya, Pharmacy Diploma Purwanto, Agus; Widya Mandala Catholic University Surabaya, Biology Puradewa, Levi; Widya Mandala Catholic University Surabaya, Pharmacy Diploma Cahyani, Erlien Dwi; Widya Mandala Catholic University Surabaya, Pharmacy Diploma Purwaningsih, Christiana Endang; Widya Mandala Catholic University Surabaya, Biology
Subject area & keyword:	Natural products < Organic chemistry



ARTICLE

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Wound Healing Activity and Flavonoids Contents of Purslane (Portulaca grandiflora) Various Variety

Antonius Budiawan *^a , Agus Purwanto ^b , Levi Puradewa ^c, Erlien Dwi Cahyani ^d, and Christiana Endang Purwaningsih ^e

Purslane has various varieties with different active metabolite contents that need to be explored further to find each variety's activity in wound healing. Different purslane herbs showed different antioxidant activities, suggesting they will have different flavonoid content and wound healing activity. This research aimed to evaluate purslane's total flavonoid content and wound-healing activity. The wounds induced on the rabbit back skin were divided into 6 treatment groups such as negative control, positive control, 10 and 20% purslane herbs extract varieties A, 10 and 20% purslane herbs extract varieties C. Wounds were treated twice daily for 2 weeks, and measured on day 0, 7, 11, and 14. Total flavonoid content was measured with the AlCl3 colorimetric method. The wounds treated with 10 and 20% purslane herbs extract varieties a (*Portulaca grandiflora* magenta flower) have 0.32±0.55 and 1.63±1.96 mm wound diameters on day 7 and healed on day 11. The wounds treated with 10 and 20% purslane herbs extract varieties C (*Portulaca grandiflora* pink flower) showed 2.88±0.51 and 0.84±1.45 mm diameter and healed on day 11. The purslane herb A showed the highest wound healing activity, and purslane varieties A and C total flavonoid content were 0.55±0.02 and 1.58±0.02% W/W, respectively.

Introduction

A wound is an injury on the outside body part tissue that disturbs physical function^{1,2}. Approximately 1% of the world's population is believed to be affected by wounds³. Meanwhile, the prevalence of wounds in Indonesia based on the proportion of injuries is 20%⁴. Indonesia is an archipelagic country prone to natural disasters and has many remote areas. Wound care will be a big problem for people living in remote areas with limited wound treatment access. Wounds not appropriately treated will cause disability or at least scars that cause discomfort⁵.

The wound healing process at the molecular level is a complex process involving various components⁶. These components include mediator cells and extracellular matrix. The stages of wound healing itself are divided into four stages: hemostasis, inflammation, proliferation, and remodeling⁷. When skin gets injured, the hemostasis stage starts and follows by an increase in platelet aggregation and the production of various chemotactic factors such as TGF-ß1, TGF-ß2, and PDGF⁸.

Inflammatory cells such as neutrophils, lymphocytes, and macrophages at the inflammatory stage will go to the wound's center to prevent infection. In the next stage (proliferation and remodeling), keratinocytes, macrophages, platelets, and endothelial cells in the wound area release various growth factors so that new cells are formed in the wound area⁹.

Expensive and unaffordable treatment costs often hamper wound healing. So wound healing therapy at a low cost is needed at this time. Medicinal plants are sources that come from nature, so they have minimal side effects and are cheaper, so they can be a safe alternative to synthetic drugs for wound treatment¹⁰. Several medicinal plants have been investigated for their healing properties; one is purslane herbs.

Purslane herbs (*Portulaca grandiflora*) is a plant that grows well in a tropical climate. Society mainly considers it a problematic plant. In addition, purslane herbs are only used as decorative plants in the front yard. This fact encourages the researchers to find other potencies that could be utilized from purslane herbs. Several studies have been conducted to find purslane efficacies on diseases, and one of these efficacies is antibacterial activity¹¹. This antibacterial activity of purslane herbs can accelerate wound healing and have the potency to be developed as wound treatment topical products.

Purslane has various varieties with different active metabolite contents that need further exploration to find each variety's activity in wound healing¹². Based on several studies, total phenol and antioxidant activity from different varieties showed different results¹³. Flavonoid is a secondary metabolite widely known for its antioxidant properties¹⁴. Different purslane herbs showed different antioxidant activities, suggesting they will have different flavonoid content and wound healing activity.

^{a.} Pharmacy Diploma III Department, Widya Mandala Surabaya Catholic University; Manggis 15-17, Madiun City 63131, East Java, Indonesia.

^{b.} Biology Department, Widya Mandala Surabaya Catholic University; Manggis 15-17, Madiun City 63131, East Java, Indonesia.

^c Pharmacy Diploma III Department, Widya Mandala Surabaya Catholic University; Manggis 15-17, Madiun City 63131, East Java, Indonesia.

^{d.} Pharmacy Diploma III Department, Widya Mandala Surabaya Catholic University; Mangais 15-17, Madiun City 63131, East Java, Indonesia.

^{e.} Biology Department, Widya Mandala Surabaya Catholic University; Manggis 15-17, Madiun City 63131, East Java, Indonesia

⁺ Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

ARTICLE

Journal Name

Based on this background, research will be needed to determine purslane's various total flavonoid content and wound healing activity.

Result and Discussion

Purslane Herbs (Portulaca grandiflora) Determination Result.

The first step of the research was the purslane herbs determination process. This step was performed to ensure the sample was the correct purslane (*Portulaca grandiflora*). The determination process was performed on purslane herbs A and C. The determination process was performed by direct morphology observation based on determination keys and observation reports^{18–22}.

The observation result showed that two varieties of purslane (*Portulaca grandiflora*) have the characteristics as follows: magenta and pink colored rose-like flowers, succulent plant, 15-30 cm high stem, round and solid stem, redness brown colored



stem skin with slight hair on the surface, slippery surface, upright or fall growing direction, and branching is forming from the base point characteristic.

The leaf is single thick, and meaty, 2-5 cm long, shiny green colored, shipshape spread and spiral structure, linier-subulatus or cylindrical shape, with sharp (acutus) tip and blunt (obtusus) pole, flat (integer) side with 1-3 cm long. The flower is bisexual, single complex, and blooms on the end of the branch with a round (obovatus) rose-like shape, the diameter is 25-35 mm, and the stamens are orange-colored with more than 20-50 stamen count. The flower pole is surrounded by 8-10 leaves, 2 petals, and the largest flower could reach 4 cm in diameter. The morphological observation is similar to the previous researcher's report²². The different *Portulaca grandiflora* variety morphology that has been used in this research was on the flower color, which was magenta (variety A) and pink with white fade (variety C) color (Figure 1).



Figure 1. The branch, leaves, and flower morphology of purslane herbs (Portulaca grandiflora).

Extraction Result.

The purslane herbs extraction used a simple maceration method and fit with a thermolabile active compound. Ethanol was used as an extraction solvent because it could give a higher extract yield and total phenol¹³. Purslane herbs C showed a higher extract yield of 10.99% compared to purslane herbs A with 9.49%.

Wound Healing Activity Test Result.

The rabbit skin has a similar aging process, wound healing time, and response to drugs to humans so it can be used as a woundhealing activity animal test 16. In addition, rabbits have broad back skin, making it easier to observe wounds visually. The **Table 1. Wounds diameter on days 0, 7, 11, dan 14.** wound healing test was started with wound induction. The hair on the back part of the rabbit was shaved with shaving tools to simplify wound induction and wounds diameter observation. Lidocaine injection was used as anesthesia because of its good local activity during the wound induction process.

The wound-healing activity test result (Table 1) based on wound diameter size on day 7 showed that purslane herbs A extract 10% concentration and purslane herbs C extract 20% concentration groups have significantly different (p<0.05) from the negative control group. A10 also showed significant differences (p<0.05) with other groups except for A20 and C20. The A10 group showed the smallest wound diameter average with 0.32±0.55 mm size. The negative control group showed the largest wound diameter average with 6.90±1.03 mm size.

Groups	Wound diameter size on the day- (mm)						
Groups	0	7	11	14			
К(-)	8.00±0.00	6.90±1.03 ^{bcdef}	4.75±0.24 ^{bcdef}	0.94±0.01			
K(+)	8.00±0.00	3.03±0.63ª	0.00±0.00 ^a	-			
A10	8.00±0.00	0.32±0.55 ^{abe}	0.00±0.00 ^a	-			
A20	8.00±0.00	1.63±1.96ª	0.00±0.00 ^a	-			
C10	8.00±0.00	2.88±0.51 ^{ac}	0.00±0.00 ^a	-			
C20	8.00±0.00	0.84±1.45 ^{abd}	0.00±0.00 ^a	-			

This journal is © The Royal Society of Chemistry 20xx

Journal Name

The wound diameter observation on day 11 significantly differed between the negative control group and other groups. The negative control group showed the largest size wound diameter average with 4.75 ± 0.24 mm size. On this day, all the groups besides the negative control group showed complete wound recovery.

The wounds treated with distilled water were still not fully recovered on day 14. The wound diameter average was measured at 0.94±0.01 mm. The different results showed by betadine solution and purslane herbs extract-treated groups that were fully healed since day 11.

The wound healing activity was shown by betadine solution and extract-treated groups. The activity was shown by wound

diameter changes that were smaller daily and healed starting from day 11 to fully recover on day 14 (Figure 2).

The different healing process was shown by the negative control group that was treated with distilled water. The wounds showed slow healing progress. The healing progressed until the end of observation, and the wounds were still not fully closed or recovered on day 14. Unfortunately, the observation was not prolonged after day 14, so the real healing timing from the negative control group was never established. However, the wounds would be healed naturally on days 13-16, depending on the size¹⁶.



Figure 2. Wounds visualization on days 0, 7, 11, and 14.

The betadine solution-treated wounds showed better healing results compared to the untreated wounds. Betadine solution contains povidone-iodine 10%, an antibacterial compound widely used as first aid to treat wounds. Povidone-iodine is an

iodoform compound that penetrates inside bacteria cells and oxidizes key protein, nucleotide, and fat acid bacteria leading to bacteria death²⁴. Without bacteria around, the wound will be healed faster because the inflammation stage of the healing

ARTICLE

process will end faster. The proliferation and remodeling will be achieved even faster rather than untreated wounds. The inflammation process is needed to transfer neutrophils, monocytes, and lymphocytes to the wound site to kill invading bacteria. This process will be prolonged if bacteria are too many to handle, so antibacterial compounds such as povidone-iodine are needed to accelerate the healing process²⁵.

The wound diameter on all purslane herbs extracts groups showed a smaller size daily and fully recovered on day 14, similar to the betadine solution-treated group. The different concentration dosages on purslane herbs A and C extract groups showed different diameter sizes even though the differences were insignificant (p>0.05). It may be concluded that different concentration dosages did not give different wound healing activities. However, the dosage variation in this research is only 10% and 20%, and the effect of broader and higher dosage concentrations on wound healing activity remains unknown.

Qualitative Phytochemical Screening Result.

Phytochemical screening analysis showed that purslane herbs variety A and C extract showed similar compounds: alkaloids, flavonoids, saponins, and tannins. Alkaloid reaction test results showed brown sediment. Furthermore, on saponin reaction test showed persistent foam on HCl addition. The color-changing to dark green showed all extracts after FeCl₃ addition showed tannins content. Brown sediment was shown after chloroform, and H_2SO_4 concentrated addition showed terpenoid contains. The result was comparable with other studies, which showed that ethanolic purslane herbs extract contains flavonoids, alkaloids, terpenoids, saponins, and tannins²⁶ (Table 2).

Table Er i arbiane nerbs of fanous fancties extract priytoenernear ser cering result
--

Extract	Flavonoid	Alkaloid	Terpenoid	Saponin	Tannin
Purslane herbs A	+	+	+	+	+
Purslane herbs C	+	+	+	+	+

Total Flavonoid Content Analysis Result.

The calibration curve determination result gives the formula y=0,0233+0,0277x with a 0.9971 r value. The higher flavonoid content was shown by purslane herbs C extract with $1.58\pm0.02\%$. The purslane herbs A extract showed smaller flavonoid content with $0.55\pm0.02\%$. Those flavonoid contents were significantly different, with a p-value <0.05. This result showed that different varieties contain different metabolite content. This is comparable with other research that showed different flavonoid content in the various purslane varieties¹².

Discussion

The determination result showed that purslane herb A is *Portulaca grandiflora* with a magenta-colored flower variety. Moreover, purslane herb C is *Portulaca grandiflora* with a pink-colored flower variety. These differences are related to secondary metabolites compound content in the purslane herbs¹². Those purslane varieties have shown antibacterial activity on *Staphylococcus aureus*, *Escherichia coli*, and *Pseudomonas aeruginosa*¹⁵. Those bacteria are commonly found in wound infections, so this extract has the potency to be developed as a wound-healing compound.

A wound is an injury that could be acquired on every body part. The open wound on the skin will increase bacteria infection risk because the skin contains many bacteria. This open wound condition will increase bacteria's chance to enter and invade the deeper tissue¹. The body's inflammation mechanism response will follow this bacteria invasion. The pro-inflammation agents such as neutrophils, macrophages, and lymphocytes will migrate in to wound site and kill the bacteria that is invading with various mechanisms such as phagocytes and apoptosis²⁵. This inflammation effect will be prolonged if the invading bacteria is too much to handle, so an active compound with

antibacterial activity is necessary to shorten this stage. The inflammation stage will also be prolonged by oxidative stress caused by free radical excessive accumulation²⁷. This condition will cause further cell obstruction, prolonging the woundhealing process. That is why the active compound with antioxidant activity also accelerates the wound healing process. Based on phytochemical screening results, ethanolic purslane herbs extract contains various secondary metabolites such as flavonoids, alkaloids, saponin, tannins, and terpenoids that have taken part in wound healing activity with various mechanisms²⁸ that were shown by wound healing activity results. The wounds treated with purslane herbs extract A and C healed with no significantly different from betadine solutiontreated wounds, especially on days 11 and 14. At the end of observation, all purslane extract-treated wounds showed a similar healed stage to betadine solution-treated wounds. Different purslane species (Portulaca oleracea) grown in Jordania showed similar results with different animal tests²⁹. Flavonoid content in purslane herb extracts is known for its antiinflammation, antibacterial, and antioxidant activity^{30,31}. As an antiinflammation, flavonoids shorten the inflammation

an antiinflammation, flavonoids shorten the inflammation stage, leading to the proliferation and remodeling stage, shortening the healing process. The antibacterial activity of flavonoids supports this activity. The inflammation process ends faster in line with the bacteria number decline. Flavonoid antibacterial activity refers to its ability to complex forming with bacteria cell membrane protein³². Flavonoids work as an antioxidant by scavenging free radicals that lead to oxidative stress prevention²⁷. Free radical is a reactive molecule with antibacterial properties in a reactive oxygen species (ROS) form. Nevertheless, in a saturated condition, ROS could cause further tissue damage at the wound site^{33,34}. Thus condition prolonged the healing process, so the antioxidant activity of flavonoids in purslane herb extracts has the benefit that could prevent

Journal Name

oxidative stress from happening^{14,34,35}. Flavonoids could donate their electrons to ROS, preventing ROS from taking electrons from molecules to tissue formation, such as DNA, protein, and lipids³³.

Total flavonoid content analysis showed that purslane herb extracts of both varieties showed flavonoid content even though this result did not align with the wound healing activity test. Purslane herb variety A extract showed higher wound healing activity than variety C extract, even though its flavonoid content was lower than purslane herb variety C extract. This may be caused by differences in other metabolite constituents in both purslane herb varieties.

Another constituent in the purslane herb suspected of its role in wound healing is an alkaloid. Alkaloids may accelerate wound healing and tissue granulation by increasing collagen and fibroblast and lowering inflammation cell numbers³⁶. Fibroblast is a connective tissue that plays in collagen production³⁷. Collagen is a connective protein responsible for elasticity and skin structure strength^{38,39}. Alkaloids also play a role in SRC/MEK/ERK signaling pathway regulation that promotes the wound healing process⁴⁰

Saponin detected in purslane herb A, and C varieties is a secondary metabolite with antibacterial activity. This activity is related to its ability to increase bacteria cell membrane permeability, leading to hemolysis⁴¹. Other than that, saponin may also increase collagen production and accelerate cell epithelization in the wound⁴². Saponin also had an essential role in high glucose-induced wound healing by increasing endothelial cell proliferation⁴³.

Tannins may also play an essential role in the wound-healing process. Tannins are known for their antibacterial activity in rebuilding damaged tissue and accelerating fibrous tissue contraction in wound healing⁴⁴. Fibrous contraction by fibroblast will trigger scar formation that protects cell formation in the wound site. Tannin's antibacterial activity may be caused by its toxicity to the microorganism related to bacteria enzyme and substrate complex formation⁴⁵. Other than this mechanism, tannins are also known for their antioxidant with similar free radical scavenging mechanism activity to flavonoids^{46,47}.

Conclusions

Two varieties of purslane herbs (magenta and pink flower) showed different wound healing activities, especially on day 7. The wound healing activity test of purslane herbs A (*Portulaca grandiflora* magenta flower variety) 10 and 20% concentration showed 0.32±0.55 dan 1.63±1.96 mm, respectively, wound diameter size. The purslane herbs C (*Portulaca grandiflora* pink flower variety) 10 and 20% concentration showed 2.88±0.51 dan 0.84±1.45 mm, respectively, wound diameter size. The purslane herbs A (*Portulaca grandiflora* magenta flower variety) 10 and 20% concentration showed 1.63±1.45 mm, respectively, wound diameter size. The purslane herbs A (*Portulaca grandiflora* magenta flower variety) 10 and 20% concentrations showed higher activity in the wound healing process on day 7 with 0.32±0.55 dan 1.63±1.96 mm, respectively, which is the smallest wound diameter. Purslane herb varieties A and C total flavonoid contents were 0.55±0.02 and 1.58±0.02% W/W, respectively.

Author Contributions

Experimental work (AB, LP, AP, EDC), data analysis (AB and CEP), writiing manuscript (AB and AP).

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The acknowledgements come at the end of an article after the conclusions and before the notes and references.

References

- 1 Price, S. A. and L. M. Wilson, Patofisiologi: Konsep Klinis Proses-proses Penyakit 6th ed., 2006.
- 2 Naseri, E. and A. Ahmadi, European Polymer Journal, 2022, 173, 111293.
- 3 P. Grover, K. Khanna, A. Bhatnagar, and J. Purkayastha, Biomedicine and Pharmacotherapy, 2021, **140**, 111797
- 4 The Ministry of Health Republic of Indonesia, Laporan Nasional Riskesdas, 2018.
- L. Zhang, Y. Tai, X. Liu, Y. Liu, C. Yang, D. Kong, C. Qi, S. Wang, A. C. Midgley, Applied Materials Today, 2021, 25, 101186.
- 6 S. Patel, S. Srivastava, M. R. Singh, and D. Singh, Biomedicine and Pharmacotherapy, 2019, **112**, 108615.
- 7 J. S. Low, K. K. Mak, S. Zhang, M. R. Pichika, P. Marappan, K. Mohandas, M. K. Balijepali, Fitoterapia, 2021, **154**, 105026.
- D. H. Assar, N. Elhabashi, A. A. A. Mokhbatly, A. E. Ragab, Z. I. Elbialy, S. A. Rizk, A. E. Albalawi, N. A. Althobaiti, S. A. Jaouni, A. Atiba, Biomedicine & pharmacotherapy, 2021, 143, 112151.
- 9 M. E. Okur, I. D. Karantas, Z. Şenyiğit, O. Üstündağ, Neslihan and P. I. Siafaka, Asian Journal of Pharmaceutical Sciences, 2020, 15, 661.
- 10 Mssillou, Ibrahim, Bakour, Meryem, Slighoua, Meryem, Laaroussi, Hassan, H. Saghrouchni, F. E. Amrati, B. Lyoussi, E. Derwich, Journal of Ethnopharmacology, 2022, 298, 115663.
- 11 Fung, K. P., Han, Q. B., Ip, M., Yang, X. S., C. Bs. Lau, B. Cl. Chan, Hong Kong medical journal, 2017, **23**, 38.
- 12 Alam, Md Amirul, Juraimi, Abdul Shukor, Rafii, M. Y., Abdul Hamid, Azizah, F. Aslani, M. M. Hasan, M. A. M. Zainudin, Md. K. Uddin, BioMed Research International, 2014, January, 6.
- 13 C. K. Lim, W. N. Tiong, and J. L. Loo, International Journal of Phytopharmacy, 2014, 4, 1.
- 14 M. T. B. Carvalho, H. G. Araújo-Filho, A. S. Barreto, L. J. Quintans-Júnior, J. S. S. Quintans, R. S. S. Barreto, Phytomedicine, 2021, **90**, 153636.
- 15 A. Purwanto, Agri-Tek: Jurnal Ilmu Pertanian, Kehutanan dan Agroteknologi, 2021, **22**, 1.
- 16 A. Grada, J. Mervis, and V. Falanga, Journal of Investigative Dermatology, 2018, **138**, 2095.
- 17 The Ministry of Health Republic of Indonesia, Materia Medika Indonesia V, 1989.
- 18 C. G. G. J. Van Steenis, Flora, 2008.
- M.K. Uddin, A. S. Juraimi, Md. S. Hossain, M. A. U. Nahar, Md. E. Ali, M. M. Rahman, The Scientific World Journal, 2014, 2014, 951019.
- 20 H. Setyowati, Continuing Professional Development, 2017, 44, 818.
- 21 E. Djauhariya, and Hernani, Gulma Berkhasiat Obat, 2004.
- 22 F. I. D. Setiawan, S. I. Aisyah, and Krisantini, Journal of Tropical Crop Science, 2016, **3**, 67.

Journal Name

- 23 T. Setyawati, S. Narulita, I. P. Bahri, and G. T. Raharjo, A Guide Book to Invasive Plant Species in Indonesia, 2015.
- 24 D. Lepelletier, J. Y. Maillard, B. Pozzetto, and A. Simon, Antimicrobial Agents and Chemotherapy, 2020, **64**, 1.
- 25 S. Guo, and L. A. Di Pietro, Journal of Dental Research, 2010, **89**, 219.
- 26 R. Londonkar, and H. Nayaka, J Pharm Res, 2011, 4, 3553.
- 27 S. D. Fitzmaurice, R. K. Sivamani, and R. R. Isseroff, Skin Pharmacology and Physiology, 2011, **24**, 113.
- 28 L. C. Adiele, R. C. Adiele, and J. C. Enye, Asian Pacific Journal of Tropical Medicine, 2014, 7, 620.
- 29 A. N. Rashed, F. U. Afifi, and A. M. Disi, Journal of Ethnopharmacology, 2003, **88**, 131.
- 30 Y. X. Zhou, H. L. Xin, K. Rahman, S. J. Wang, C. Peng, H. Zhang, BioMed Research International, 2015, **2015**, 925631.
- 31 M. H. Masoodi, B. Ahmad, S. R. Mir, B. A. Zargar, and N. Tabasum, Journal of Pharmacy Research, 2011, **44**, 3044.
- 32 A. Royani, M. Hanafi, H. Julistiono, and A. Manaf, Materials Today: Proceedings, 2022.
- 33 C. Dunnill, T. Patton, J. Brennan, J. Barrett, M. Dryden, J. Cooke, D. Leaper, N. T. Georgopoulos, International Wound Journal, 2017, 14, 89.
- 34 A. Priyadarshi, G. K. Keshri, and A. Gupta, Phytomedicine Plus, 2022, **2**, 100292.
- 35 E. Asante-Kwatia, A. Evelyn, J. Silas, G. Yakubu, G. A. H. Lord, G. Adjei-Hinneh, I. K. Amponsah, A. Y. Mensah, Heliyon, 2021, 7, e08261.
- 36 T. Nagappan, T. C. Segaran, M. E. A. Wahid, P. Ramasamy, and C. S. Vairappan, Molecules, 2012, **17**, 14449.
- 37 L. Bao, X. Cai, M. Zhang, Y. Xiao, J. Jin, T. Qin, Y. Li, Journal of Functional Foods, 2022, **90**, 104981.
- 38 P. Bainbridge, Journal of Wound Care, 2013, 22, 407.
- 39 S. Sharma, V. K. Rai, R. K. Narang, and T. S. Markandeywar, Life Sciences, 2022, **290**, 120096.
- 40 X.Q. Shi, G. Chen, J. Q. Tan, Z. Li, S. M. Chen, J. H. He, L. Zhang, H. X. Xu, Journal of Ethnopharmacology, 2022, **295**, 115396.
- 41 M. P. Wei, H. Yu, Y. H. Guo, Y. L. Cheng, Y. F. Xie, W. R. Yao, Food Control, 2021, **130**, 108337.
- 42 D. MacKay, and A. L. Miller, Alternative Medicine Review, 2003, 8, 359.
- 43 T. Lei, Y. Gao, Y. Duan, C. Cui, L. Zhang, M. Si, Environmental Toxicology, 2022, **37**, 35385194.
- 44 A. Santana, A. Barros, H. Oliveira, and I. Victor, Biomedicine & Pharmacotherapy, 2017, **96**, 182.
- 45 A. Scalbert, Phytochemistry, 1991, 30, 3875.
- 46 M. Heinonen, Molecular Nutrition and Food Research, 2007, **51**, 684.
- 47 C. M. González, E. Llorca, A. Quiles, I. Hernando, and G. Moraga, LWT Food Science and Technology, 2022, **155**.

February 9th, 2023

Dear Editor of RSC Advances

Please find our article titled **Wound Healing Activity and Flavonoids Contents of Purslane** (*Portulaca grandiflora*) Various Variety. We hope that this article and any part that has never been submitted or published in any journal will be considered for publication in RSC Advances.

In this article, we determine various purslane (*Portulaca grandiflora*) herb varieties wound healing activity and their total flavonoid content. *Portulaca grandiflora* has various varieties with different active metabolite content related to its wound-healing activity, especially flavonoids.

It is an honor if our original article fits RSC Advances' aims, scope, and standard quality and is to be considered for publication.

Sincerely yours,

Antonius Budiawan Department of Pharmacy Diploma III Widya Mandala Surabaya Catholic University Manggis 15-17, Madiun City 63131, East Java, Indonesia Email: <u>antonius.budiawan@ukwms.ac.id</u> Telephone +6281217767966



Track the progress of your RSC Advances submission

1 message

RSC Advances <noreply@rsc.org> To: Mr Antonius Budiawan <antonius.budiawan@ukwms.ac.id> Thu, Feb 9, 2023 at 1:19 PM

Dear Mr Antonius Budiawan

Thank you for submitting your manuscript to RSC Advances. We have safely received the files and information you have provided.

Our team are currently processing your submission and will be in touch shortly.

You can use the following link to track the status of your submission. You can also share this link with your co-authors so that they can keep track of the manuscript's progress too.

https://submissions.rsc.org/tracker/RA-ART-02-2023-000868?t=ZKN9CQL792Y8xdmpilgzYUtRBJE9szleFs4TCVkq8R6CWQ%3D%3D

Yours sincerely, RSC Advances Editorial Office advances@rsc.org

Please do not reply to this email.

2. First revision (major revision) (10-3-2023)



Decision on submission to RSC Advances - RA-ART-02-2023-000868

1 message

RSC Advances <onbehalfof@manuscriptcentral.com> Reply-To: advances@rsc.org To: antonius.budiawan@ukwms.ac.id Fri, Mar 10, 2023 at 1:24 PM

10-Mar-2023

Dear Mr Budiawan:

Manuscript ID: RA-ART-02-2023-000868

TITLE: Wound Healing Activity and Flavonoids Contents of Purslane (Portulaca grandiflora) Various Variety

Thank you for your submission to RSC Advances, published by the Royal Society of Chemistry. I sent your manuscript to reviewers and I have now received their reports which are copied below.

I have carefully evaluated your manuscript and the reviewers' reports, and the reports indicate that major revisions are necessary.

Please submit a revised manuscript which addresses all of the reviewers' comments. Further peer review of your revised manuscript may be needed. When you submit your revised manuscript please include a point by point response to the reviewers' comments and highlight the changes you have made. Full details of the files you need to submit are listed at the end of this email.

Please submit your revised manuscript as soon as possible using this link:

*** PLEASE NOTE: This is a two-step process. After clicking on the link, you will be directed to a webpage to confirm. ***

https://mc.manuscriptcentral.com/rscadv?URL_MASK=bcec58b70ff048e09f562ea5beda1145

(This link goes straight to your account, without the need to log on to the system. For your account security you should not share this link with others.)

Alternatively, you can log in to your account (https://mc.manuscriptcentral.com/rscadv) where you will need your case-sensitive user ID and password details.

You should submit your revised manuscript as soon as possible; please note you will receive a series of automatic reminders. If your revisions will take a significant length of time, please contact me.

All RSC Advances articles are published under an open access model, and the appropriate article processing charge (APC) will apply. Details of the APC and discounted rates can be found at https://www.rsc.org/journals-books-databases/about-journals/rsc-advances/# charges-RSC-Advances

RSC Advances strongly encourages authors of research articles to include an 'Author contributions' section in their manuscript, for publication in the final article. This should appear immediately above the 'Conflict of interest' and 'Acknowledgement' sections. I strongly recommend you use CRediT (the Contributor Roles Taxonomy, https://credit.niso.org/) for standardised contribution descriptions. All authors should have agreed to their individual contributions ahead of submission and these should accurately reflect contributions to the work. Please refer to our general author guidelines https://www.rsc.org/journals-books-databases/author-and-reviewer-hub/authors-information/responsibilities/ for more information.

The Royal Society of Chemistry requires the submitting author to provide their ORCID iD when they submit a revised manuscript. This is quick and easy to do as part of the revised manuscript submission process. We will publish this information with the article, and you may choose to have your ORCID record updated automatically with details of the publication.

Please also encourage your co-authors to sign up for their own ORCID account and associate it with their account on our manuscript submission system. Please note that we are unable to do this on behalf of your co-authors. For further information see: https://www.rsc.org/journals-books-databases/journal-authors-reviewers/processes-policies/#attribution-id.

I look forward to receiving your revised manuscript.

Yours sincerely, Professor Luigi Vaccaro Associate Editor, RSC Advances

REVIEWER REPORT(S): Referee: 1 Recommendation: Minor revisions

Comments:

The authors determined various purslane (Portulaca grandiflora) herb varieties wound healing

activity and their total flavonoid content. Some minor concerns should be addressed.

1. Abstract, AICI3 colorimetric method, change to AICI₃

2. Introduction, "The wound healing process at the molecular level is a complex process involving various components." Delete the first "process".

3. Table 1. Provide the meaning of superscript "abcde".

Additional Questions: Does the work significantly advance the understanding or development in this field?: Yes

Is this work of relevance to the chemistry community?: Yes

Are the conclusions of the work convincing and sufficiently supported by experimental evidence?: Yes

Does the data provided fulfil the journal's data requirements?

See Journal specific guidelines: https://www.rsc.org/journals-books-databases/about-journals/rsc-advances/#Characterisation-of-new-compounds: Yes

Is the experimental section sufficiently detailed to allow others to reproduce the work?: Yes

Are the reported claims adequately discussed in the context of the literature?: Yes

Are the number of tables and figures in the manuscript appropriate and clear?: Yes

Referee: 2

Recommendation: Major revisions

Comments:

In this manuscript, the activity of wound healing of Purslane (Portulaca grandiflora) was evaluated. The authors carefully examined wound healing effects in in vivo study. This study was well-designed and the target sample was unique. However, the following points are not appropriately described and needed to be revised.

Page 3 right part

"Based on several studies, total phenol and antioxidant activity from different varieties showed different results. Flavonoid is a secondary metabolite widely known for its antioxidant properties. Different purslane herbs showed different antioxidant activities, suggesting they will have different flavonoid content and wound healing activity." The authors should provide a more detailed explanation by breaking down the "different" part, such as giving a few more specific

The authors should provide a more detailed explanation by breaking down the "different" part, such as giving a few more specific examples."

Page 4

Are there any descriptions of materials and methods? The authors must show the experimental method including statistics and number of replicates for this experiment. Moreover, what product was used for negative control, just water?

Page 6 5th line from the top and the bottom line

"This process will be prolonged if bacteria are too many to handle,"

"This inflammation effect will be prolonged if the invading bacteria is too much to handle,"

The description should be correct and unified.

Additional Questions:

Does the work significantly advance the understanding or development in this field?: Yes

Is this work of relevance to the chemistry community?: No

Are the conclusions of the work convincing and sufficiently supported by experimental evidence?: Yes

Does the data provided fulfil the journal's data requirements?

See Journal specific guidelines: <a href="https://www.rsc.org/journals-books-databases/about-journals/rsc-advances/#Characterisation-

of-new-compounds" target="_new">https://www.rsc.org/journals-books-databases/about-journals/rsc-advances/#Characterisation-of-new-compounds: Yes

Is the experimental section sufficiently detailed to allow others to reproduce the work?: No

Are the reported claims adequately discussed in the context of the literature?: Yes

Are the number of tables and figures in the manuscript appropriate and clear?: Yes

FILES TO PROVIDE WITH YOUR REVISED MANUSCRIPT:

IMPORTANT: Your original files are available to you when you upload your revised manuscript. Please delete any redundant files before completing the submission. Please carefully check the spelling and format of all author names, affiliations and funding information. If your paper is accepted for publication, it is important this information is accurate to ensure your article is correctly indexed, which may affect citations and future funding evaluation. Please note that if you have selected Accepted Manuscript publication, the author list will appear as provided in the ScholarOne submission details until your Advance Article is published and this information is updated from your article.

- A point-by-point response to the comments made by the reviewer(s)
- Your revised manuscript with any changes clearly marked (.doc(x) or.pdf file)

• Your revised manuscript as a .doc(x) file including figures, without highlighting, track changes, etc. (If providing in TeX format instead, please also provide a final PDF version including figures). Please note that we cannot proceed with publication using a .pdf file only.

High quality images EITHER embedded in a doc(x) file OR

as separate numbered Figures, Schemes or Charts in .tif, .eps or .pdf format, with a resolution of 600 dpi or greater.

AND

A table of contents entry: graphic maximum size 8 cm x 4 cm and 1-2 sentence(s) of editable text, with a maximum of 250 characters (including spaces), highlighting the key findings of the work. It is recommended authors make use of the full space available for the graphic. See our Author Guidelines for more details: https://www.rsc.org/journals-books-databases/author-and-reviewer-hub/
 Your revised Electronic Supplementary Information (if any)

• Your revised CheckCIF reports (if any). Please ensure that any revised cif files have been deposited with the Cambridge Crystallographic Data Centre (CCDC) via https://deposit.ccdc.cam.ac.uk/ before you submit your revised manuscript.

For Review-type articles only:

• A photograph and biography of yourself and your co-authors. Separate photographs of each author may be supplied or if you prefer, a group photograph, saved as a .tif, .pdf or .jpeg file. The resolution of the photographs should be 600 dpi or higher. The dimensions of the photograph in the printed journal will be 4 cm wide x 5 cm high (individual photograph) or 8.3 cm wide x 5 cm high (group photograph). Individual photographs should be accompanied by a maximum of 100 words; a group photograph by a maximum of 200 words. There can be a maximum of 6 individual biographies per article.

• Copies of permissions required from other publishers to reproduce figures. Please ensure that necessary permissions are acknowledged in the figure captions in accordance with the publishers' instruction. Information on how to obtain permissions and rights that we require are given on our website at

https://www.rsc.org/journals-books-databases/journal-authors-reviewers/licences-copyright-permissions/

If you need to contact the journal, please use the email address advances@rsc.org

DISCLAIMER:

This communication is from The Royal Society of Chemistry, a company incorporated in England by Royal Charter (registered number RC000524) and a charity registered in England and Wales (charity number 207890). Registered office: Burlington House, Piccadilly, London W1J 0BA. Telephone: +44 (0) 20 7437 8656.

The content of this communication (including any attachments) is confidential, and may be privileged or contain copyright material. It may not be relied upon or disclosed to any person other than the intended recipient(s) without the consent of The Royal Society of Chemistry. If you are not the intended recipient(s), please (1) notify us immediately by replying to this email, (2) delete all copies from your system, and (3) note that disclosure, distribution, copying or use of this communication is strictly prohibited.

Any advice given by The Royal Society of Chemistry has been carefully formulated but is based on the information available to it. The Royal Society of Chemistry cannot be held responsible for accuracy or completeness of this communication or any attachment. Any views or opinions presented in this email are solely those of the author and do not represent those of The Royal Society of Chemistry. The views expressed in this communication are personal to the sender and unless specifically stated, this e-mail does not constitute any part of an offer or contract. The Royal Society of Chemistry shall not be liable for any resulting damage or loss as a result of the use of this email and/or attachments, or for the consequences of any actions taken on the basis of the information provided. The Royal Society

of Chemistry does not warrant that its emails or attachments are Virus-free; The Royal Society of Chemistry has taken reasonable precautions to ensure that no viruses are contained in this email, but does not accept any responsibility once this email has been transmitted. Please rely on your own screening of electronic communication.

More information on The Royal Society of Chemistry can be found on our website: www.rsc.org

Responses to reviewers' comments

Dear Editor-In-Chief

We are grateful for the editors' and reviewers' constructive comments and suggestions that we have implemented in our manuscript. The suggested changes are highlighted in our revised manuscript and we also provide point by point responses to the reviewer's comments as below.

Referee: 1

Recommendation: Minor revisions

Comments:

The authors determined various purslane (Portulaca grandiflora) herb varieties wound healing activity and their total flavonoid content. Some minor concerns should be addressed.

Response: Thanks for supporting this manuscript's publication and for constructive comments and suggestions to our manuscript.

1. Abstract, AlCl3 colorimetric method, change to AlCl₃

Response: We have revised it per the reviewer's suggestion; please refer to the abstract section.

2. Introduction, "The wound healing process at the molecular level is a complex process involving various components." Delete the first "process".

Response: We have revised it per the reviewer's suggestion; please refer to the introduction section

3. Table 1. Provide the meaning of superscript "abcde".

Response: We have provided the meaning of the superscript as suggested by the reviewer, please refer to Table 1.

Referee: 2

Recommendation: Major revisions

Comments:

In this manuscript, the activity of wound healing of Purslane (Portulaca grandiflora) was evaluated. The authors carefully examined wound healing effects in in vivo study. This study

was well-designed and the target sample was unique. However, the following points are not appropriately described and needed to be revised.

Response: Thanks for supporting this manuscript's publication and for constructive comments and suggestions to our manuscript.

Page 3 right part

"Based on several studies, total phenol and antioxidant activity from different varieties showed different results. Flavonoid is a secondary metabolite widely known for its antioxidant properties. Different purslane herbs showed different antioxidant activities, suggesting they will have different flavonoid content and wound healing activity."

The authors should provide a more detailed explanation by breaking down the "different" part, such as giving a few more specific examples."

Response: Thanks for the suggestion; we have provided more detailed explanation as suggested.

Page 4

Are there any descriptions of materials and methods?

The authors must show the experimental method including statistics and number of replicates for this experiment. Moreover, what product was used for negative control, just water?

Response: Thanks for the suggestion; we have provided the detail of the experimental method in our revised manuscript; please refer experimental section in our revised version of the manuscript.

Page 6 5th line from the top and the bottom line

"This process will be prolonged if bacteria are too many to handle,"

"This inflammation effect will be prolonged if the invading bacteria is too much to handle," The description should be correct and unified.

Response: We have corrected and unified the sentences as suggested by the reviewer.

Response to Publishing Assistant

Dear Publishing Assistant

Royal Society of Chemistry - RSC Advances

Thank you for the consideration and suggestion for our manuscript. We have implemented the live subject statement addition needed in our manuscript.

Yours sincerely

Antonius Budiawan

3. Revised version submitted (15-3-2023)



Acknowledgement of your revised manuscript submission to RSC Advances - RA-ART-02-2023-000868.R1

1 message

 RSC Advances <onbehalfof@manuscriptcentral.com>
 Wed, Mar 15, 2023 at 4:20 PM

 Reply-To: advances@rsc.org
 To: antonius.budiawan@ukwms.ac.id

 To: antonius.budiawan@ukwms.ac.id, agus.purwanto@ukwms.ac.id, lpuradewa@gmail.com, erlien.dwi.cahyani@ukwms.ac.id, endang.ch.purwaningsih@ukwms.ac.id

15-Mar-2023

Dear Mr Budiawan:

TITLE: Wound Healing Activity and Flavonoids Contents of Purslane (<i>Portulaca grandiflora</i>) Various Variety AUTHORS: Budiawan, Antonius; Purwanto, Agus; Puradewa, Levi; Cahyani, Erlien Dwi; Purwaningsih, Christiana Endang

Thank you for your revised submission to RSC Advances, published by the Royal Society of Chemistry. This is an automatic acknowledgement that you have uploaded your files to our online submission system. Your manuscript ID is: RA-ART-02-2023-000868.R1

Your manuscript will be passed to an editor for initial assessment as soon as possible. If there are any problems with your submission we will contact you.

All RSC Advances articles are published under an open access model, and the appropriate article processing charge (APC) will apply. Details of the APC and discounted rates can be found at https://www.rsc.org/journals-books-databases/about-journals/rsc-advances/# charges-RSC-Advances

Please indicate the above manuscript ID when you contact us about this submission.

If there have been any changes to the list of authors during this revision, please inform the Editorial Office of the reason for the change. All authors must confirm they are happy with the authorship change by emailing advances@rsc.org.

Do you have an ORCID iD? ORCID (Open Researcher and Contributor iD) is a unique researcher identifier that allows you to link your research output and other professional activities in a single record. We therefore encourage each researcher to sign up for their own ORCID account. Please edit your user account to link your ORCID iD or create a new one, ensuring that you have not linked your account to another researcher's ORCID iD. Please note that we are unable to do this on your behalf. If your article is accepted, you may choose to have your ORCID record updated automatically with details of the publication.

We already have the following information for authors of this manuscript: Budiawan, Antonius - https://orcid.org/0000-0003-3730-3913, Purwanto, Agus - https://orcid.org/0009-0009-3325-2291, Puradewa, Levi - https://orcid.org/0000-0003-1426-3103, Cahyani, Erlien Dwi - https://orcid.org/0009-0009-6325-0684, Purwaningsih, Christiana Endang - https://orcid.org/0009-0003-3078-6008 If this is not how you want your name to appear on an Accepted Manuscript, please amend your ScholarOne account.

This journal has a policy of sharing reviewer reports. Please note that the Editor's decision and copies of the reports will be shared with all reviewers who provide a recommendation on your manuscript.

Please contact us if we can be of any assistance.

Yours sincerely, RSC Advances Editorial Office advances@rsc.org If you need to contact the journal, please use the email address advances@rsc.org

DISCLAIMER:

This communication is from The Royal Society of Chemistry, a company incorporated in England by Royal Charter (registered number RC000524) and a charity registered in England and Wales (charity number 207890). Registered office: Burlington House, Piccadilly, London W1J 0BA. Telephone: +44 (0) 20 7437 8656.

The content of this communication (including any attachments) is confidential, and may be privileged or contain copyright material. It may not be relied upon or disclosed to any person other than the intended recipient(s) without the consent of The Royal Society of Chemistry. If you are not the intended recipient(s), please (1) notify us immediately by replying to this email, (2) delete all copies from your system, and (3) note that disclosure, distribution, copying or use of this communication is strictly prohibited.

Any advice given by The Royal Society of Chemistry has been carefully formulated but is based on the information available to it. The Royal Society of Chemistry cannot be held responsible for accuracy or completeness of this communication or any attachment. Any views or opinions presented in this email are solely those of the author and do not represent those of The Royal Society of Chemistry. The views expressed in this communication are personal to the sender and unless specifically stated, this e-mail does not constitute any part of an offer or contract. The Royal Society of Chemistry shall not be liable for any resulting damage or loss as a result of the use of this email and/or attachments, or for the consequences of any actions taken on the basis of the information provided. The Royal Society of Chemistry does not warrant that its emails or attachments are Virus-free; The Royal Society of Chemistry has taken reasonable precautions to ensure that no viruses are contained in this email, but does not accept any responsibility once this email has been transmitted. Please rely on your own screening of electronic communication.

More information on The Royal Society of Chemistry can be found on our website: www.rsc.org

4. Paper accepted (17-3-2023)



Decision on submission to RSC Advances - RA-ART-02-2023-000868.R1

1 message

RSC Advances <onbehalfof@manuscriptcentral.com> Reply-To: advances@rsc.org To: antonius.budiawan@ukwms.ac.id Fri, Mar 17, 2023 at 10:03 PM

17-Mar-2023

Dear Mr Budiawan:

Manuscript ID: RA-ART-02-2023-000868.R1

TITLE: Wound Healing Activity and Flavonoids Contents of Purslane (<i>Portulaca grandiflora</i>) Various Variety

Thank you for submitting your revised manuscript to RSC Advances. I am pleased to accept your manuscript for publication in its current form.

You will shortly receive a separate email from us requesting you to submit a licence to publish for your article, so that we can proceed with the preparation and publication of your manuscript. We will email you information on how to access your RSC Advances article proofs when they are ready.

All RSC Advances articles are published under an open access model, and the appropriate article processing charge (APC) will apply. Details of the APC and discounted rates can be found at https://www.rsc.org/journals-books-databases/about-journals/rsc-advances/# charges-RSC-Advances.

We are offering all corresponding authors who are not already members of the Royal Society of Chemistry one year's Affiliate membership as part of their APC. If you would like to find out more please email membership@rsc.org, including the promo code OA100 in your message. Learn all about our member benefits at https://www.rsc.org/membership-and-community/join/#benefit.

Promote your research, accelerate its impact – find out more about our article promotion services here: https://rsc.li/ promoteyourresearch.

If you would like us to promote your article on our Twitter account @RSCAdvances please fill out this form: https://form.jotform.com/ 212705294847058. Alternatively, if you would like your article to be considered for further promotion via our WeChat channels then please do let us know.

By publishing your article in RSC Advances, you are supporting the Royal Society of Chemistry to help the chemical science community make the world a better place.

With best wishes,

Professor Luigi Vaccaro Associate Editor, RSC Advances

If you need to contact the journal, please use the email address advances@rsc.org

DISCLAIMER:

This communication is from The Royal Society of Chemistry, a company incorporated in England by Royal Charter (registered number RC000524) and a charity registered in England and Wales (charity number 207890). Registered office: Burlington House, Piccadilly, London W1J 0BA. Telephone: +44 (0) 20 7437 8656.

The content of this communication (including any attachments) is confidential, and may be privileged or contain copyright material. It may not be relied upon or disclosed to any person other than the intended recipient(s) without the consent of The Royal Society of Chemistry. If you are not the intended recipient(s), please (1) notify us immediately by replying to this email, (2) delete all copies from your system, and (3) note that disclosure, distribution, copying or use of this communication is strictly prohibited.

Any advice given by The Royal Society of Chemistry has been carefully formulated but is based on the information available to it. The Royal Society of Chemistry cannot be held responsible for accuracy or completeness of this communication or any attachment. Any views or opinions presented in this email are solely those of the author and do not represent those of The Royal Society of Chemistry. The views expressed in this communication are personal to the sender and unless specifically stated, this e-mail does not constitute any part of an offer or contract. The Royal Society of Chemistry shall not be liable for any resulting damage or loss as a result of the use of

this email and/or attachments, or for the consequences of any actions taken on the basis of the information provided. The Royal Society of Chemistry does not warrant that its emails or attachments are Virus-free; The Royal Society of Chemistry has taken reasonable precautions to ensure that no viruses are contained in this email, but does not accept any responsibility once this email has been transmitted. Please rely on your own screening of electronic communication.

More information on The Royal Society of Chemistry can be found on our website: www.rsc.org

5.Additional revision from editor (Live subject statement) (19-3-2023)



RSC Advances: RA-ART-02-2023-000868.R1

1 message

RSC Advances <onbehalfof@manuscriptcentral.com> Reply-To: RSC.Editorial_office@straive.com To: antonius.budiawan@ukwms.ac.id Mon, Mar 20, 2023 at 5:31 AM

19-Mar-2023

Dear Mr Budiawan:

TITLE: Wound Healing Activity and Flavonoids Contents of Purslane (<i>Portulaca grandiflora</i>) Various Variety

I notice that your submission contains studies involving live subjects rabbit), in such cases our policy requires you to provide a statement containing the information described below.

LIVE SUBJECT STATEMENT:

- that all experiments were performed in compliance with relevant laws or guidelines;
- that all experiments followed institutional guidelines;
- a statement on the institutional committee(s) that approved the experiments;
- if human subjects have been used, a statement that informed consent was obtained must also be provided.

For reference only, please see the sample ethical statement below:

"All animal procedures were performed in accordance with the Guidelines for Care and Use of Laboratory Animals of "____" University and approved by the Animal Ethics Committee of "____".

Kindly provide the main article with a detailed ethical statement to RSC.Editorial_office@straive.com.

Please contact me if I can be of any assistance.

Yours sincerely, Dorina Monique T. Lucillo Publishing Assistant Royal Society of Chemistry - RSC Advances

If you need to contact the journal, please use the email address advances@rsc.org

DISCLAIMER:

This communication is from The Royal Society of Chemistry, a company incorporated in England by Royal Charter (registered number RC000524) and a charity registered in England and Wales (charity number 207890). Registered office: Burlington House, Piccadilly, London W1J 0BA. Telephone: +44 (0) 20 7437 8656.

The content of this communication (including any attachments) is confidential, and may be privileged or contain copyright material. It may not be relied upon or disclosed to any person other than the intended recipient(s) without the consent of The Royal Society of Chemistry. If you are not the intended recipient(s), please (1) notify us immediately by replying to this email, (2) delete all copies from your system, and (3) note that disclosure, distribution, copying or use of this communication is strictly prohibited.

Any advice given by The Royal Society of Chemistry has been carefully formulated but is based on the information available to it. The Royal Society of Chemistry cannot be held responsible for accuracy or completeness of this communication or any attachment. Any views or opinions presented in this email are solely those of the author and do not represent those of The Royal Society of Chemistry. The views expressed in this communication are personal to the sender and unless specifically stated, this e-mail does not constitute any part of an offer or contract. The Royal Society of Chemistry shall not be liable for any resulting damage or loss as a result of the use of this email and/or attachments, or for the consequences of any actions taken on the basis of the information provided. The Royal Society of Chemistry does not warrant that its emails or attachments are Virus-free; The Royal Society of Chemistry has taken reasonable precautions to ensure that no viruses are contained in this email, but does not accept any responsibility once this email has been transmitted. Please rely on your own screening of electronic communication.

More information on The Royal Society of Chemistry can be found on our website: www.rsc.org

ARTICLE

Wound Healing Activity and Flavonoids Contents of Purslane (Portulaca grandiflora) Various Variety

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Antonius Budiawan *^a , Agus Purwanto ^b , Levi Puradewa ^c, Erlien Dwi Cahyani ^d, and Christiana Endang Purwaningsih ^e

Purslane has various varieties with different active metabolite contents that need to be explored further to find each variety's activity in wound healing. Different purslane herbs showed different antioxidant activities, suggesting they will have different flavonoid content and wound healing activity. This research aimed to evaluate purslane's total flavonoid content and wound-healing activity. The wounds induced on the rabbit back skin were divided into 6 treatment groups such as negative control, positive control, 10 and 20% purslane herbs extract varieties A, 10 and 20% purslane herbs extract varieties C. Wounds were treated twice daily for 2 weeks, and measured on day 0, 7, 11, and 14. Total flavonoid content was measured with the AlCl₃ colorimetric method. The wounds treated with 10 and 20% purslane herbs extract varieties a (*Portulaca grandiflora* magenta flower) have 0.32±0.55 and 1.63±1.96 mm wound diameters on day 7 and healed on day 11. The wounds treated with 10 and 20% purslane herbs extract varieties C (*Portulaca grandiflora* pink flower) showed 2.88±0.51 and 0.84±1.45 mm diameter and healed on day 11. The purslane herb A showed the highest wound healing activity, and purslane varieties A and C total flavonoid content were 0.55±0.02 and 1.58±0.02% W/W, respectively.

Introduction

A wound is an injury on the outside body part tissue that disturbs physical function^{1,2}. Approximately 1% of the world's population is believed to be affected by wounds³. Meanwhile, the prevalence of wounds in Indonesia based on the proportion of injuries is 20%⁴. Indonesia is an archipelagic country prone to natural disasters and has many remote areas. Wound care will be a big problem for people living in remote areas with limited wound treatment access. Wounds not appropriately treated will cause disability or at least scars that cause discomfort⁵.

Wound healing at the molecular level is a complex process involving various components⁶. These components include mediator cells and extracellular matrix. The stages of wound healing itself are divided into four stages: hemostasis, inflammation, proliferation, and remodeling⁷. When skin gets injured, the hemostasis stage starts and follows by an increase in platelet aggregation and the production of various chemotactic factors such as TGF-ß1, TGF-ß2, and PDGF⁸. Inflammatory cells such as neutrophils, lymphocytes, and macrophages at the inflammatory stage will go to the wound's center to prevent infection. In the next stage (proliferation and remodeling), keratinocytes, macrophages, platelets, and endothelial cells in the wound area release various growth factors so that new cells are formed in the wound area⁹.

Expensive and unaffordable treatment costs often hamper wound healing. So wound healing therapy at a low cost is needed at this time. Medicinal plants are sources that come from nature, so they have minimal side effects and are cheaper, so they can be a safe alternative to synthetic drugs for wound treatment¹⁰. Several medicinal plants have been investigated for their healing properties; one is purslane herbs.

Purslane herbs (*Portulaca grandiflora*) is a plant that grows well in a tropical climate. Society mainly considers it a problematic plant. In addition, purslane herbs are only used as decorative plants in the front yard. This fact encourages the researchers to find other potencies that could be utilized from purslane herbs. Several studies have been conducted to find purslane efficacies on diseases, and one of these efficacies is antibacterial activity¹¹. This antibacterial activity of purslane herbs can accelerate wound healing and have the potency to be developed as wound treatment topical products.

Purslane has various varieties with different active metabolite contents that need further exploration to find each variety's activity in wound healing¹². Based on several studies, total phenol and antioxidant activity from different flower colour varieties showed different results¹³. Flavonoid is a secondary metabolite widely known for its antioxidant properties¹⁴. Different flower colour varieties of purslane herbs showed different antioxidant activities, suggesting they will have

^{a.} Pharmacy Diploma III Department, Widya Mandala Surabaya Catholic University; Manggis 15-17, Madiun City 63131, East Java, Indonesia.

^{b.} Biology Department, Widya Mandala Surabaya Catholic University; Manggis 15-17, Madiun City 63131, East Java, Indonesia.

^c Pharmacy Diploma III Department, Widya Mandala Surabaya Catholic University; Manggis 15-17, Madiun City 63131, East Java, Indonesia.

^{d.} Pharmacy Diploma III Department, Widya Mandala Surabaya Catholic University; Manggis 15-17, Madiun City 63131, East Java, Indonesia.

^{e.} Biology Department, Widya Mandala Surabaya Catholic University; Manggis 15-17, Madiun City 63131, East Java, Indonesia

⁺ Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

ARTICLE

different flavonoid content and wound healing activity. Based on this background, research will be needed to determine purslane's various total flavonoid content and wound healing activity.

Result and Discussion

Purslane Herbs (Portulaca grandiflora) Determination Result.

The first step of the research was the purslane herbs determination process. This step was performed to ensure the sample was the correct purslane (*Portulaca grandiflora*). The determination process was performed on purslane herbs A and C. The determination process was performed by direct morphology observation based on determination keys and observation reports^{18–22}.

The observation result showed that two varieties of purslane (*Portulaca grandiflora*) have the characteristics as follows: magenta and pink colored rose-like flowers, succulent plant, 15-



30 cm high stem, round and solid stem, redness brown colored stem skin with slight hair on the surface, slippery surface, upright or fall growing direction, and branching is forming from the base point characteristic.

The leaf is single thick, and meaty, 2-5 cm long, shiny green colored, shipshape spread and spiral structure, linier-subulatus or cylindrical shape, with sharp (acutus) tip and blunt (obtusus) pole, flat (integer) side with 1-3 cm long. The flower is bisexual, single complex, and blooms on the end of the branch with a round (obovatus) rose-like shape, the diameter is 25-35 mm, and the stamens are orange-colored with more than 20-50 stamen count. The flower pole is surrounded by 8-10 leaves, 2 petals, and the largest flower could reach 4 cm in diameter. The morphological observation is similar to the previous researcher's report²². The different *Portulaca grandiflora* variety morphology that has been used in this research was on the flower color, which was magenta (variety A) and pink with white fade (variety C) color (Figure 1).



Figure 1. The branch, leaves, and flower morphology of purslane herbs (Portulaca grandiflora).

Extraction Result.

The purslane herbs extraction used a simple maceration method and fit with a thermolabile active compound. Ethanol was used as an extraction solvent because it could give a higher extract yield and total phenol¹³. Purslane herbs C showed a higher extract yield of 10.99% compared to purslane herbs A with 9.49%.

Wound Healing Activity Test Result.

The rabbit skin has a similar aging process, wound healing time, and response to drugs to humans so it can be used as a wound-healing activity animal test¹⁶. In addition, rabbits have broad back skin, making it easier to observe wounds visually. The **Table 1. Wounds diameter on days 0, 7, 11, dan 14.**

wound healing test was started with wound induction. The hair on the back part of the rabbit was shaved with shaving tools to simplify wound induction and wounds diameter observation. Lidocaine injection was used as anesthesia because of its good local activity during the wound induction process.

The wound-healing activity test result (Table 1) based on wound diameter size on day 7 showed that purslane herbs A extract 10% concentration and purslane herbs C extract 20% concentration groups have significantly different (p<0.05) from the negative control group. A10 also showed significant differences (p<0.05) with other groups except for A20 and C20. The A10 group showed the smallest wound diameter average with 0.32±0.55 mm size. The negative control group showed the largest wound diameter average with 6.90±1.03 mm size.

	D '	1 - 1	0	
Α	к			

Crowns	Wound diameter size on the day- (mm)						
Groups	0	7	11	14			
К(-)	8.00±0.00	6.90±1.03 ^{bcdef}	4.75±0.24 ^{bcdef}	0.94±0.01			
К(+)	8.00±0.00	3.03±0.63ª	0.00±0.00ª	-			
A10	8.00±0.00	0.32±0.55 ^{abe}	0.00±0.00 ^a	-			
A20	8.00±0.00	1.63±1.96ª	0.00±0.00 ^a	-			
C10	8.00±0.00	2.88±0.51 ^{ac}	0.00±0.00 ^a	-			
C20	8.00±0.00	0.84±1.45 ^{abd}	0.00±0.00 ^a	-			

Note: a Significantly different from K(-) (negative control); b Significantly different from K(+) (positive control); c Significantly different from A10 (purslane herbs A extract 10% concentration); ^dSignificantly different from A20 (purslane herbs A extract 20% concentration); "Significantly different from C10 (purslane herbs C extract 10% concentration); "Significantly different from C20 (purslane herbs C extract 20% concentration).

The wound diameter observation on day 11 significantly differed between the negative control group and other groups. The negative control group showed the largest size wound diameter average with 4.75±0.24 mm size. On this day, all the groups besides the negative control group showed complete wound recovery.

The wounds treated with distilled water were still not fully recovered on day 14. The wound diameter average was measured at 0.94±0.01 mm. The different results showed by betadine solution and purslane herbs extract-treated groups that were fully healed since day 11.

The wound healing activity was shown by betadine solution and extract-treated groups. The activity was shown by wound

diameter changes that were smaller daily and healed starting from day 11 to fully recover on day 14 (Figure 2).

The different healing process was shown by the negative control group that was treated with distilled water. The wounds showed slow healing progress. The healing progressed until the end of observation, and the wounds were still not fully closed or recovered on day 14. Unfortunately, the observation was not prolonged after day 14, so the real healing timing from the negative control group was never established. However, the wounds would be healed naturally on days 13-16, depending on the size¹⁶.



ARTICLE

Journal Name



Figure 2. Wounds visualization on days 0, 7, 11, and 14.

The betadine solution-treated wounds showed better healing results compared to the untreated wounds. Betadine solution contains povidone-iodine 10%, an antibacterial compound widely used as first aid to treat wounds. Povidone-iodine is an iodoform compound that penetrates inside bacteria cells and oxidizes key protein, nucleotide, and fat acid bacteria leading to bacteria death²⁴. Without bacteria around, the wound will be healed faster because the inflammation stage of the healing process will end faster. The proliferation and remodeling will be achieved even faster rather than untreated wounds. The inflammation process is needed to transfer neutrophils, monocytes, and lymphocytes to the wound site to kill invading bacteria. This process will be prolonged if bacteria are too many to handle, so antibacterial compounds such as povidone-iodine are needed to accelerate the healing process²⁵.

The wound diameter on all purslane herbs extracts groups showed a smaller size daily and fully recovered on day 14, similar to the betadine solution-treated group. The different concentration dosages on purslane herbs A and C extract groups showed different diameter sizes even though the differences were insignificant (p>0.05). It may be concluded that different concentration dosages did not give different wound healing activities. However, the dosage variation in this research is only 10% and 20%, and the effect of broader and higher dosage concentrations on wound healing activity remains unknown.

Qualitative Phytochemical Screening Result.

Phytochemical screening analysis showed that purslane herbs variety A and C extract showed similar compounds: alkaloids, flavonoids, saponins, and tannins. Alkaloid reaction test results showed brown sediment. Furthermore, on saponin reaction test showed persistent foam on HCl addition. The color-changing to dark green showed all extracts after FeCl₃ addition showed tannins content. Brown sediment was shown after chloroform, and H_2SO_4 concentrated addition showed terpenoid contains. The result was comparable with other studies, which showed that ethanolic purslane herbs extract contains flavonoids, alkaloids, terpenoids, saponins, and tannins²⁶ (Table 2).

Table 2. Fulsiane nerbs of various varieties extract phytochemical screening resu	Table 2. Pu	rslane herbs o	of various varieties	extract ph	vtochemical	I screening resu	ılt
---	-------------	----------------	----------------------	------------	-------------	------------------	-----

Extract	Flavonoid	Alkaloid	Terpenoid	Saponin	Tannin
Purslane herbs A	+	+	+	+	+
Purslane herbs C	+	+	+	+	+

Total Flavonoid Content Analysis Result.

The calibration curve determination result gives the formula y=0,0233+0,0277x with a 0.9971 r value. The higher flavonoid content was shown by purslane herbs C extract with $1.58\pm0.02\%$. The purslane herbs A extract showed smaller flavonoid content with $0.55\pm0.02\%$. Those flavonoid contents were significantly different, with a p-value <0.05. This result showed that different varieties contain different metabolite content. This is comparable with other research that showed different flavonoid content in the various purslane varieties¹².

Discussion

The determination result showed that purslane herb A is *Portulaca grandiflora* with a magenta-colored flower variety. Moreover, purslane herb C is *Portulaca grandiflora* with a pink-colored flower variety. These differences are related to secondary metabolites compound content in the purslane herbs¹². Those purslane varieties have shown antibacterial activity on *Staphylococcus aureus*, *Escherichia coli*, and *Pseudomonas aeruginosa*¹⁵. Those bacteria are commonly

Journal Name

A wound is an injury that could be acquired on every body part. The open wound on the skin will increase bacteria infection risk because the skin contains many bacteria. This open wound condition will increase bacteria's chance to enter and invade the deeper tissue¹. The body's inflammation mechanism response will follow this bacteria invasion. The pro-inflammation agents such as neutrophils, macrophages, and lymphocytes will migrate into wound site and kill the bacteria that is invading with various mechanisms such as phagocytes and apoptosis²⁵. This process will be prolonged if bacteria are too many to handle, so an active compound with antibacterial activity is necessary to shorten this stage. The inflammation stage will also be prolonged by oxidative stress caused by free radical excessive accumulation²⁷. This condition will cause further cell obstruction, prolonging the wound-healing process. That is why the active compound with antioxidant activity also accelerates the wound healing process.

Based on phytochemical screening results, ethanolic purslane herbs extract contains various secondary metabolites such as flavonoids, alkaloids, saponin, tannins, and terpenoids that have taken part in wound healing activity with various mechanisms²⁸ that were shown by wound healing activity results. The wounds treated with purslane herbs extract A and C healed with no significantly different from betadine solutiontreated wounds, especially on days 11 and 14. At the end of observation, all purslane extract-treated wounds showed a similar healed stage to betadine solution-treated wounds. Different purslane species (*Portulaca oleracea*) grown in Jordania showed similar results with different animal tests²⁹.

Flavonoid content in purslane herb extracts is known for its antiinflammation, antibacterial, and antioxidant activity^{30,31}. As an antiinflammation, flavonoids shorten the inflammation stage, leading to the proliferation and remodeling stage, shortening the healing process. The antibacterial activity of flavonoids supports this activity. The inflammation process ends faster in line with the bacteria number decline. Flavonoid antibacterial activity refers to its ability to complex forming with bacteria cell membrane protein³². Flavonoids work as an antioxidant by scavenging free radicals that lead to oxidative stress prevention²⁷. Free radical is a reactive molecule with antibacterial properties in a reactive oxygen species (ROS) form. Nevertheless, in a saturated condition, ROS could cause further tissue damage at the wound site^{33,34}. Thus condition prolonged the healing process, so the antioxidant activity of flavonoids in purslane herb extracts has the benefit that could prevent oxidative stress from happening^{14,34,35}. Flavonoids could donate their electrons to ROS, preventing ROS from taking electrons from molecules to tissue formation, such as DNA, protein, and lipids³³.

Total flavonoid content analysis showed that purslane herb extracts of both varieties showed flavonoid content even though this result did not align with the wound healing activity test. Purslane herb variety A extract showed higher wound healing activity than variety C extract, even though its flavonoid content was lower than purslane herb variety C extract. This may be caused by differences in other metabolite constituents in both purslane herb varieties.

Another constituent in the purslane herb suspected of its role in wound healing is an alkaloid. Alkaloids may accelerate wound healing and tissue granulation by increasing collagen and fibroblast and lowering inflammation cell numbers³⁶. Fibroblast is a connective tissue that plays in collagen production³⁷. Collagen is a connective protein responsible for elasticity and skin structure strength^{38,39}. Alkaloids also play a role in SRC/MEK/ERK signaling pathway regulation that promotes the wound healing process⁴⁰

Saponin detected in purslane herb A, and C varieties is a secondary metabolite with antibacterial activity. This activity is related to its ability to increase bacteria cell membrane permeability, leading to hemolysis⁴¹. Other than that, saponin may also increase collagen production and accelerate cell epithelization in the wound⁴². Saponin also had an essential role in high glucose-induced wound healing by increasing endothelial cell proliferation⁴³.

Tannins may also play an essential role in the wound-healing process. Tannins are known for their antibacterial activity in rebuilding damaged tissue and accelerating fibrous tissue contraction in wound healing⁴⁴. Fibrous contraction by fibroblast will trigger scar formation that protects cell formation in the wound site. Tannin's antibacterial activity may be caused by its toxicity to the microorganism related to bacteria enzyme and substrate complex formation⁴⁵. Other than this mechanism, tannins are also known for their antioxidant with similar free radical scavenging mechanism activity to flavonoids^{46,47}.

Experimental Section

Materials.

Tools to conduct in vivo wound healing activity tests were scissors and razors to cut the rabbit hair, 1 ml disposable syringe, biopsy punch 8 mm diameter, anatomical forceps, scalpel blade, vernier calipers to measure wound diameter, and evaporator. Analytical vaccum rotarv balance. spectrophotometer UV-Vis, and glassware were used to perform phytochemical and total flavonoid content analysis. The sample of two varieties of purslane was obtained from Madiun City of Java Island in Indonesia. The solvent that was used in the extraction process was ethanol 70%. The New Zealand rabbit was used as an animal test to perform wound healing activity. Lidocaine injection was used as a local anesthetic on rabbit skin when wound excision was conducted. Betadine solution and distilled water were used as the positive and negative control, respectively. The materials used for phytochemical screening and total flavonoid content analysis were ethanol 96% pro analysis, AICl₃, potassium acetate, and distilled water.

Methods.

Purslane herbs characterization (Portulaca grandiflora). Two kinds of purslane herbs obtained from Madiun City were characterized to know their variety. Determination was

ARTICLE

conducted in Biology Laboratory, Biology Department, Widya Mandala Surabaya Catholic University.

Extraction process.

Two hundred grams of each dried purslane herbs variety were macerated with 500 ml ethanol 70% for 5 days. The solid residue from the maceration process was macerated once more with the same amount of solvent. The first and second macerates were mixed and evaporated with a vacuum rotary evaporator at 50°C until a thick extract was obtained. The thick extract was used for phytochemical screening, total flavonoid content analysis, and in vivo wound healing activity tests.

Wound healing activity test.

The rabbit was used as an animal test to conduct in vivo wound healing activity. The New Zealand rabbit used in this study was 3-5 months old and had a 1.5-2.5 kg weight. Ethical clearance was obtained from Health Research Ethics Committee Faculty of Medicine, Muhammadiyah Surakarta University, Indonesia.

Rabbit's hair on the back part was shaved with shaving tools. The rabbit skin was anesthetized with lidocaine with 0.0125-0.7600 ml/kg BW dose until showing unresponsive when stabbed with a needle. A biopsy punch of 8 mm diameter was used to make wound excision on rabbit skin. Three rabbits were induced with six wounds for each rabbit.

The wounds on the rabbit's back skin were grouped into 6 treatment groups such as negative control (distilled water), positive control (betadine solution), 10 and 20% purslane herbs extract varieties A, 10 and 20% purslane herbs extract varieties C.

The wounds were cleaned and disinfected with an antiseptic solution and waited until the blood stopped before the treatment was applied twice daily for two weeks. The macroscopic condition observation and wound diameter measurement was conducted on days 0, 7, 11, and 14.

Qualitative Phytochemical Screening.

The flavonoid identification was conducted by dissolving 0.5 grams of each various variety of purslane ethanolic extract in the distilled water and added with magnesium and 5 drops of HCl 2 N. The mixture was heated for 5-10 minutes. After filtration and cooling, the filtrate was added with amyl alcohol and shaken vigorously. The reaction result is positive if red color is formed in the amyl alcohol layer.

The alkaloid identification was conducted by 0.5 grams of each various variety of purslane ethanolic extract was added with 1 ml ammonia and then added with chloroform. The mixture was filtrated, and the filtrate was placed in the test tube and added with 2N HCl. The mixture was shaken and waited until separation occurred. The filtrate was divided into three parts in the different test tubes. One drop of Dragendorff reagent was added to the first filtrate. The reaction result is positive if brown sediment is formed. One drop of Mayer reagent was added to the second filtrate, and the reaction was positive if white sediment formed. The third filtrate is blank or negative control. The saponin identification was conducted by placing 0.5 grams of each various variety of purslane ethanolic extract into a test

tube and then dissolved with hot distilled water. After the mixture was cold, it was shaken for 10 seconds. The reaction is positive if 1-10 cm persistent foam is formed for 10 minutes, and stays still after adding 1-2 drops of 2N HCl.

The terpenoid identification was conducted by placing 0.5 grams of each various variety of purslane ethanolic extract into a test tube and then added with chloroform and H_2SO_4 concentrate. The reaction is positive if brown sediment is formed.

The tannin identification was conducted by placing 0.5 grams of each various variety of purslane ethanolic extract into a test tube and then added with 1% FeCl₃ solvent. The reaction is positive if green or dark blue color is formed.

The total flavonoid content of various varieties of purslane was measured with AlCl₃ colorimetric methods. Twenty-five mg of quercetin was dissolved with 25 ml ethanol in a 25 ml volumetric flask to obtain a 1000 µg/ml initial solution. Standard solutions of 10, 20, 40, 60, 80, and 100 ppm were made from the initial solution. Each standard solution was pipetted 0.5 ml and then was added with 1.5 ml ethanol 96%, 1 ml 10% AlCl₃, 1 ml potassium acetate 1 M, and 2.8 ml distilled water. The solution was incubated at 25°C for 30 minutes. The absorbance was measured with a spectrophotometer UV-Vis on a wavelength (λ) of 434.2 nm. The absorbance reading result was used to make a calibration curve of absorbance vs. quercetin concentration.

Five grams of each various variety of purslane ethanolic extract was dissolved with 25 ml ethanol and stirred at 200 rpm, and then filtrated. The filtrate was added to 25 ml ethanol. The blank solution for the flavonoid content assay was made from 0.5 ml ethanol that was added with 1.5 ml ethanol 96%, 1 ml 10% AlCl₃, 1 ml potassium acetate 1 M, and 2.8 ml distilled water. One ml of various purslane ethanolic extract was added with ethanol until the marking sign of the 10 ml volumetric flask. Half ml of the solution was pipetted and added with 1.5 ml ethanol 96%, 1 ml 10% AlCl3, 1 ml potassium acetate 1 M, and 2.8 ml distilled water. The mixture was incubated at 25°C for 30 minutes, and the absorbance was read with а spectrophotometer UV-Vis on λ 434,2 nm. The total flavonoid content was counted with the following equation:

$F = (c \times V \times f \times 10-6)/m \times 100\%$

Note:

- F = flavonoid content
- c = quercetin equality (g/ml)
- V = total extract volume
- f = dilution factor
- m = sample weigth (g)

Data Analysis.

The observed data were statistically analyzed with a 95% (p<0.05) confidence limit using the Kruskal-Wallis analysis test.

Live Subject Statement.

All animal procedures were performed under the guidelines for care and use of laboratory animals of Widya Mandala Surabaya

Catholic University, Indonesia, and approved by the Health Research Ethics Committee Faculty of Medicine, Muhammadiyah Surakarta University, Indonesia.

Conclusions

Two varieties of purslane herbs (magenta and pink flower) showed different wound healing activities, especially on day 7. The wound healing activity test of purslane herbs A (*Portulaca grandiflora* magenta flower variety) 10 and 20% concentration showed 0.32±0.55 dan 1.63±1.96 mm, respectively, wound diameter size. The purslane herbs C (*Portulaca grandiflora* pink flower variety) 10 and 20% concentration showed 2.88±0.51 dan 0.84±1.45 mm, respectively, wound diameter size. The purslane herbs A (*Portulaca grandiflora* magenta flower variety) 10 and 20% concentration showed 2.88±0.51 dan 0.84±1.45 mm, respectively, wound diameter size. The purslane herbs A (*Portulaca grandiflora* magenta flower variety) 10 and 20% concentrations showed higher activity in the wound healing process on day 7 with 0.32±0.55 dan 1.63±1.96 mm, respectively, which is the smallest wound diameter. Purslane herb varieties A and C total flavonoid contents were 0.55±0.02 and 1.58±0.02% W/W, respectively.

Author Contributions

Experimental work (AB, LP, AP, EDC), data analysis (AB and CEP), and writing the manuscript (AB and AP).

Conflicts of interest

There are no conflicts to declare.

Acknowledgments

The acknowledgments come at the end of an article after the conclusions and before the notes and references.

References

- 1 Price, S. A. and L. M. Wilson, Patofisiologi: Konsep Klinis Proses-proses Penyakit 6th ed., 2006.
- 2 Naseri, E. and A. Ahmadi, European Polymer Journal, 2022, 173, 111293.
- 3 P. Grover, K. Khanna, A. Bhatnagar, and J. Purkayastha, Biomedicine and Pharmacotherapy, 2021, **140**, 111797
- 4 The Ministry of Health Republic of Indonesia, Laporan Nasional Riskesdas, 2018.
- L. Zhang, Y. Tai, X. Liu, Y. Liu, C. Yang, D. Kong, C. Qi, S. Wang, A. C. Midgley, Applied Materials Today, 2021, 25, 101186.
- 6 S. Patel, S. Srivastava, M. R. Singh, and D. Singh, Biomedicine and Pharmacotherapy, 2019, **112**, 108615.
- 7 J. S. Low, K. K. Mak, S. Zhang, M. R. Pichika, P. Marappan, K. Mohandas, M. K. Balijepali, Fitoterapia, 2021, **154**, 105026.
- D. H. Assar, N. Elhabashi, A. A. A. Mokhbatly, A. E. Ragab, Z. I. Elbialy, S. A. Rizk, A. E. Albalawi, N. A. Althobaiti, S. A. Jaouni, A. Atiba, Biomedicine & pharmacotherapy, 2021, 143, 112151.
- 9 M. E. Okur, I. D. Karantas, Z. Şenyiğit, O. Üstündağ, Neslihan and P. I. Siafaka, Asian Journal of Pharmaceutical Sciences, 2020, 15, 661.

- 10 Mssillou, Ibrahim, Bakour, Meryem, Slighoua, Meryem, Laaroussi, Hassan, H. Saghrouchni, F. E. Amrati, B. Lyoussi, E. Derwich, Journal of Ethnopharmacology, 2022, 298, 115663.
- 11 Fung, K. P., Han, Q. B., Ip, M., Yang, X. S., C. Bs. Lau, B. Cl. Chan, Hong Kong medical journal, 2017, **23**, 38.
- 12 Alam, Md Amirul, Juraimi, Abdul Shukor, Rafii, M. Y., Abdul Hamid, Azizah, F. Aslani, M. M. Hasan, M. A. M. Zainudin, Md. K. Uddin, BioMed Research International, 2014, January, 6.
- 13 C. K. Lim, W. N. Tiong, and J. L. Loo, International Journal of Phytopharmacy, 2014, 4, 1.
- 14 M. T. B. Carvalho, H. G. Araújo-Filho, A. S. Barreto, L. J. Quintans-Júnior, J. S. S. Quintans, R. S. S. Barreto, Phytomedicine, 2021, **90**, 153636.
- 15 A. Purwanto, Agri-Tek: Jurnal Ilmu Pertanian, Kehutanan dan Agroteknologi, 2021, 22, 1.
- 16 A. Grada, J. Mervis, and V. Falanga, Journal of Investigative Dermatology, 2018, **138**, 2095.
- 17 The Ministry of Health Republic of Indonesia, Materia Medika Indonesia V, 1989.
- 18 C. G. G. J. Van Steenis, Flora, 2008.
- M.K. Uddin, A. S. Juraimi, Md. S. Hossain, M. A. U. Nahar, Md. E. Ali, M. M. Rahman, The Scientific World Journal, 2014, 2014, 951019.
- 20 H. Setyowati, Continuing Professional Development, 2017, 44, 818.
- 21 E. Djauhariya, and Hernani, Gulma Berkhasiat Obat, 2004.
- 22 F. I. D. Setiawan, S. I. Aisyah, and Krisantini, Journal of Tropical Crop Science, 2016, **3**, 67.
- 23 T. Setyawati, S. Narulita, I. P. Bahri, and G. T. Raharjo, A Guide Book to Invasive Plant Species in Indonesia, 2015.
- 24 D. Lepelletier, J. Y. Maillard, B. Pozzetto, and A. Simon, Antimicrobial Agents and Chemotherapy, 2020, **64**, 1.
- S. Guo, and L. A. Di Pietro, Journal of Dental Research, 2010, 89, 219.
- 26 R. Londonkar, and H. Nayaka, J Pharm Res, 2011, 4, 3553.
- 27 S. D. Fitzmaurice, R. K. Sivamani, and R. R. Isseroff, Skin Pharmacology and Physiology, 2011, **24**, 113.
- 28 L. C. Adiele, R. C. Adiele, and J. C. Enye, Asian Pacific Journal of Tropical Medicine, 2014, 7, 620.
- 29 A. N. Rashed, F. U. Afifi, and A. M. Disi, Journal of Ethnopharmacology, 2003, **88**, 131.
- 30 Y. X. Zhou, H. L. Xin, K. Rahman, S. J. Wang, C. Peng, H. Zhang, BioMed Research International, 2015, 2015, 925631.
- 31 M. H. Masoodi, B. Ahmad, S. R. Mir, B. A. Zargar, and N. Tabasum, Journal of Pharmacy Research, 2011, 44, 3044.
- 32 A. Royani, M. Hanafi, H. Julistiono, and A. Manaf, Materials Today: Proceedings, 2022.
- 33 C. Dunnill, T. Patton, J. Brennan, J. Barrett, M. Dryden, J. Cooke, D. Leaper, N. T. Georgopoulos, International Wound Journal, 2017, 14, 89.
- 34 A. Priyadarshi, G. K. Keshri, and A. Gupta, Phytomedicine Plus, 2022, **2**, 100292.
- E. Asante-Kwatia, A. Evelyn, J. Silas, G. Yakubu, G. A. H. Lord, G. Adjei-Hinneh, I. K. Amponsah, A. Y. Mensah, Heliyon, 2021, 7, e08261.
- 36 T. Nagappan, T. C. Segaran, M. E. A. Wahid, P. Ramasamy, and C. S. Vairappan, Molecules, 2012, 17, 14449.
- 37 L. Bao, X. Cai, M. Zhang, Y. Xiao, J. Jin, T. Qin, Y. Li, Journal of Functional Foods, 2022, **90**, 104981.
- 38 P. Bainbridge, Journal of Wound Care, 2013, 22, 407.
- 39 S. Sharma, V. K. Rai, R. K. Narang, and T. S. Markandeywar, Life Sciences, 2022, **290**, 120096.
- 40 X.Q. Shi, G. Chen, J. Q. Tan, Z. Li, S. M. Chen, J. H. He, L. Zhang, H. X. Xu, Journal of Ethnopharmacology, 2022, 295, 115396.
- 41 M. P. Wei, H. Yu, Y. H. Guo, Y. L. Cheng, Y. F. Xie, W. R. Yao, Food Control, 2021, **130**, 108337.
- 42 D. MacKay, and A. L. Miller, Alternative Medicine Review, 2003, 8, 359.

- 43 T. Lei, Y. Gao, Y. Duan, C. Cui, L. Zhang, M. Si, Environmental Toxicology, 2022, **37**, 35385194.
- 44 A. Santana, A. Barros, H. Oliveira, and I. Victor, Biomedicine & Pharmacotherapy, 2017, **96**, 182.
- 45 A. Scalbert, Phytochemistry, 1991, 30, 3875.
- 46 M. Heinonen, Molecular Nutrition and Food Research, 2007, **51**, 684.
- 47 C. M. González, E. Llorca, A. Quiles, I. Hernando, and G. Moraga, LWT Food Science and Technology, 2022, **155**.

6. Revised version submitted (20-3-2023)



Main Article RA-ART-02-2023-000868.R1

1 message

Antonius Budiawan M.Farm. Apt. <antonius.budiawan@ukwms.ac.id> To: RSC.Editorial_office@straive.com Mon, Mar 20, 2023 at 2:08 PM

Dear Publishing Assistant

Royal Society of Chemistry - RSC Advances

Thank you for the consideration and suggestion for our manuscript. We have implemented the live subject statement addition needed in our manuscript.

Yours sincerely,

Antonius Budiawan

RSC Advances_Portulaca grandiflora Wound Healing_REV.docx 3405K



Join the RSC Advances reviewer community

1 message

RSC Advances <onbehalfof@manuscriptcentral.com> Reply-To: advances@rsc.org To: antonius.budiawan@ukwms.ac.id Tue, Mar 21, 2023 at 7:01 AM

21-Mar-2023

Dear Mr Budiawan:

Thank you for choosing to publish in RSC Advances, published by the Royal Society of Chemistry. Your paper was accepted for publication thanks to the reviewer(s) who provided reports, and so I hope that you found their comments constructive.

We value the input of all of our reviewers to improve the quality of our publications and RSC Advances is continually seeking to expand its reviewer community.

If you do not currently act as a reviewer for RSC Advances, I invite you to become a reviewer. Visit https://rsc.li/reviewer to find out more and register your interest. Please also log into your account and add/update your current research interests so that our editors can match you with the most appropriate submitted manuscripts.

If you do currently review for us, please ensure your account information is up to date. The information you give regarding your research interests and areas of expertise is essential for editors to match submitted manuscripts with reviewers, and to ensure that you are asked to review the most appropriate manuscripts.

You can add/update your research interests by logging into your account (https://mc.manuscriptcentral.com/rscadv), clicking on your name at the top of the screen, selecting 'User ID & Password' from the list, and then completing the 'Research Interests' box (at end of page). You may also select a number of keywords describing your particular area(s) of expertise. Under the "Keywords" heading, in the "Search on this list" box you can search the list using the wildcard * for keywords relevant to your area of expertise, e.g. ENZYM* to find keywords related to enzymes.

Please also link your ORCID iD to your account. For a step-by-step guide see:

https://www.rsc.org/journals-books-databases/journal-authors-reviewers/processes-policies/#attribution-id

Please contact me if I can be of any assistance.

Yours sincerely, Dorina Monique T. Lucillo Publishing Assistant Royal Society of Chemistry - RSC Advances If you need to contact the journal, please use the email address advances@rsc.org

DISCLAIMER:

This communication is from The Royal Society of Chemistry, a company incorporated in England by Royal Charter (registered number RC000524) and a charity registered in England and Wales (charity number 207890). Registered office: Burlington House, Piccadilly, London W1J 0BA. Telephone: +44 (0) 20 7437 8656.

The content of this communication (including any attachments) is confidential, and may be privileged or contain copyright material. It may not be relied upon or disclosed to any person other than the intended recipient(s) without the consent of The Royal Society of Chemistry. If you are not the intended recipient(s), please (1) notify us immediately by replying to this email, (2) delete all copies from your system, and (3) note that disclosure, distribution, copying or use of this communication is strictly prohibited.

Any advice given by The Royal Society of Chemistry has been carefully formulated but is based on the information available to it. The Royal Society of Chemistry cannot be held responsible for accuracy or completeness of this communication or any attachment. Any views or opinions presented in this email are solely those of the author and do not represent those of The Royal Society of Chemistry. The views expressed in this communication are personal to the sender and unless specifically stated, this e-mail does not constitute any part of an offer or contract. The Royal Society of Chemistry shall not be liable for any resulting damage or loss as a result of the use of this email and/or attachments, or for the consequences of any actions taken on the basis of the information provided. The Royal Society of Chemistry does not warrant that its emails or attachments are Virus-free; The Royal Society of Chemistry has taken reasonable precautions to ensure that no viruses are contained in this email, but does not accept any responsibility once this email has been transmitted. Please rely on your own screening of electronic communication.

More information on The Royal Society of Chemistry can be found on our website: www.rsc.org

7.Permition to sign a license to publish (21-3-2023)



Required: Licence to Publish - D3RA00868A

1 message

RSC Advances <RSC1@rsc.org> To: antonius.budiawan@ukwms.ac.id Tue, Mar 21, 2023 at 7:23 AM

Tuesday, March 21, 2023

Dear Mr Budiawan,

TITLE: Wound Healing Activity and Flavonoids Contents of Purslane (Portulaca grandiflora) Various Variety MANUSCRIPT ID: D3RA00868A

Before we can publish your manuscript in RSC Advances, we need you to sign a licence to publish.

Follow this link to choose and sign your licence to publish:

https://licences.rsc.org/email-link?id=D3RA00868A&t=ctECCWCf4mA2qZeriVstY0tTD5c1%2FTo%3D

We are unable to publish your manuscript until a valid licence has been received, so please sign the licence as soon as possible to avoid delays. If you would like to publish your article open access, relevant licence options are also available through the above link. Please reply to this email if you have any questions.

Thank you for choosing to publish with RSC Advances, a journal published by the Royal Society of Chemistry.

Yours sincerely, RSC Advances Editorial Office

DISCLAIMER:

8.Sign accepted (21-3-2023)



Confirmation: Licence to Publish signed - D3RA00868A

1 message

RSC Advances <RSC1@rsc.org> To: antonius.budiawan@ukwms.ac.id Tue, Mar 21, 2023 at 10:35 AM

Tuesday, March 21, 2023

Dear Mr Budiawan,

TITLE: Wound Healing Activity and Flavonoids Contents of Purslane (Portulaca grandiflora) Various Variety MANUSCRIPT ID: D3RA00868A

We are pleased to confirm we have received your signed licence to publish for your article D3RA00868A.

Your article will be published gold open access.

You will receive a separate email explaining how to access your article proofs.

Yours sincerely, RSC Advances Editorial Office

DISCLAIMER:



Payment details required for your article - D3RA00868A

1 message

RSC Advances <RSC1@rsc.org> To: antonius.budiawan@ukwms.ac.id Tue, Mar 21, 2023 at 10:35 AM

Tuesday, March 21, 2023

Dear Mr Budiawan,

TITLE: Wound Healing Activity and Flavonoids Contents of Purslane (Portulaca grandiflora) Various Variety JOURNAL: RSC Advances MANUSCRIPT ID: D3RA00868A

You have requested to publish your article gold open access and are therefore required to pay an article processing charge(APC) of £1000.00 (+ tax).

Please note that membership or institutional discounts may apply. This will be shown to you when you make your payment following the link below.

In our hybrid journals we are unable to publish your article until we have received payment, so please provide payment information as soon as possible to avoid delays.

Follow this link to provide payment information:

https://licences.rsc.org/email-link?id=D3RA00868A&t=ctECCWCf4mA2qZeriVstY0tTD5c2%2FDw%3D

You may pass this link on to your institution or funder to submit payment details on your behalf. Please be aware that in doing so, the recipient of the email will be able to view the following information:

Manuscript title

List of Authors and their Institutions

Funding details

Award / grant numbers

You can make a payment by card by following the link, completing the form, and choosing the 'Pay with a Credit or Debit card' option.

To issue you with an invoice we require details of who will pay the APC, including full name, institution, email and street address. We also require a VAT number (UK and Europe) or GST number (India) if applicable.

Once the form has been completed, we will send an invoice by email within one week; this will contain all the details needed to make the payment. Card payments will also receive an invoice (marked as paid) for your reference.

If you have already made a card payment online, you can ignore this email; you will receive a separate email confirming payment.

Please reply to this email if you have any questions.

Yours sincerely, Customer Services team Royal Society of Chemistry RSC1@rsc.org

DISCLAIMER:

9. Proof reading (22-3-2023)



Your RSC Advances article (D3RA00868A) proof is ready for checking

1 message



Dear Mr Antonius Budiawan

Wound Healing Activity and Flavonoids Contents of Purslane (<i>Portulaca grandiflora</i>) Various Variety

The proof for your article is now ready for checking. Please review and correct your proof in Proof Central. This link gives you corresponding author access to the proof and should not be shared unless you wish to delegate the submission of the proof corrections to another author. If you would like to share the proof with your co-authors, please use the "Invite Collaborators" feature in Proof Central. This gives each author you invite a unique link so their changes are tracked, allowing the corresponding author to review and approve all changes before submission. Co-authors invited through the "Invite Collaborators" feature are not able to submit the final corrections.

Proof Central provides you with a user-friendly interface to help you manage your proofs quickly and efficiently. You can find out more about Proof Central here. You can find instructions on the submission of your corrections at Proof Central.

Please return your corrections within 48 hours. If you are unable to return your corrections within 48 hours, please let us know.

If you prefer to send your corrections by email, please download your proof using Proof Central and send your corrections to advances@rsc.org

The page numbers on the proof are not the final page numbers of your article, we will send these once your article is published in an issue of *RSC Advances*.

If you have a waiver request pending, please do not return your proof corrections; due to our fast publication times we will be unable to stop or delay the publication of your article. If your waiver is not granted you will be required to pay the Article Processing Charge according to the terms and conditions which you accepted when you submitted your article.

With best wishes, From the Editors of RSC Advances Royal Society of Chemistry advances@rsc.org

If you are unable to access your proofs using the links above, please paste the following link into your browser: https://rscjournals.proofcentral.com/en-us/landing-page.html?token= 9612bqd10eec6032e7aa2074fab296

www.rsc.org

Registered charity number: 207890

You have been sent this message because you have submitted an article to RSC Advances. If you feel that you have received this in error, please contact advances@rsc.org

The Royal Society of Chemistry, Thomas Graham House, Science Park, Cambridge CB4 0WF, United Kingdom. Registered charity number: 207890.

© Royal Society of Chemistry [[:Year]]. All rights reserved.

DISCLAIMER:

This communication is from The Royal Society of Chemistry, a company incorporated in England by Royal Charter (registered number RC000524) and a charity registered in England and Wales (charity number 207890). Registered office: Burlington House, Piccadilly, London W1J 0BA. Telephone: +44 (0) 20 7437 8656.

The content of this communication (including any attachments) is confidential, and may be privileged or contain copyright material. It may not be relied upon or disclosed to any person other than the intended recipient(s) without the consent of The Royal Society of Chemistry. If you are not the intended recipient(s), please (1) notify us immediately by replying to this email, (2) delete all copies from your system, and (3) note that disclosure, distribution, copying or use of this communication is strictly prohibited.

Any advice given by The Royal Society of Chemistry has been carefully formulated but is based on the information available to it. The Royal Society of Chemistry cannot be held responsible for accuracy or completeness of this communication or any attachment. Any views or opinions presented in this email are solely those of the author and do not represent those of The Royal Society of Chemistry. The views expressed in this communication are personal to the sender and unless specifically stated, this e-mail does not constitute any part of an offer or contract. The Royal Society of Chemistry shall not be liable for any resulting damage or loss as a result of the use of this email and/or attachments, or for the consequences of any actions taken on the basis of the information provided. The Royal Society of Chemistry does not warrant that its emails or attachments are Virus-free; The Royal Society of Chemistry has taken reasonable precautions to ensure that no viruses are contained in this email, but does not accept any responsibility once this email has been transmitted. Please rely on your own screening of electronic communication.

More information on The Royal Society of Chemistry can be found on our website: www.rsc.org

This communication is from The Royal Society of Chemistry, a company incorporated in England by Royal Charter (registered number RC000524) and a charity registered in England and Wales (charity number 207890). Registered office: Burlington House, Piccadilly, London W1J 0BA. Telephone: +44 (0) 20 7437 8656.

The content of this communication (including any attachments) is confidential, and may be privileged or contain copyright material. It may not be relied upon or disclosed to any person other than the intended recipient(s) without the consent of The Royal Society of Chemistry. If you are not the intended recipient(s), please (1) notify us immediately by replying to this email, (2) delete all copies from your system, and (3) note that disclosure, distribution, copying or use of this communication is strictly prohibited.

Any advice given by The Royal Society of Chemistry has been carefully formulated but is based on the information available to it. The Royal Society of Chemistry cannot be held responsible for accuracy or completeness of this communication or any attachment. Any views or opinions presented in this email are solely those of the author and do not represent those of The Royal Society of Chemistry. The views expressed in this communication are personal to the sender and unless specifically stated, this e-mail does not constitute any part of an offer or contract. The Royal Society of Chemistry shall not be liable for any resulting damage or loss as a result of the use of this email and/or attachments, or for the consequences of any actions taken on the basis of the information provided. The Royal Society of Chemistry does not warrant that its emails or attachments are Virus-free; The Royal Society of Chemistry has taken reasonable precautions to ensure that no viruses are contained in this email, but does not accept any responsibility once this email has been transmitted. Please rely on your own screening of electronic communication.

More information on The Royal Society of Chemistry can be found on our website: www.rsc.org

Wound healing activity and flavonoid contents of purslane (*Portulaca grandiflora*) of various varieties



Table of Contents Entry



Different flower colour varieties of purslane (*Portulaca grandiflora*) showed different total flavonoid content. Flavonoids have a main role in wound healing activity in rabbit skin as antibacterial, antiinflammation, and antioxidant agents.

Abstract

Purslane has various varieties with different active metabolite contents that need to be explored further to find each variety's activity in wound healing. Different purslane herbs showed different antioxidant activities, suggesting they will have different flavonoid content and wound healing activity. This research aimed to evaluate purslane's total flavonoid content and wound-healing activity. The wounds induced on the rabbit back skin were divided into 6 treatment groups such as negative control, positive control, 10 and 20% purslane herbs extract varieties A, and 10 and 20% purslane herbs extract varieties C. Wounds were treated twice daily for 2 weeks, and measured on day 0, 7, 11, and 14. Total flavonoid content was measured with the AlCl₃ colorimetric method. The wounds treated with 10 and 20% purslane herbs extract varieties A (*Portulaca grandiflora* magenta flower) have 0.32 ± 0.55 and 1.63 ± 1.96 mm wound diameters on day 7 and healed on day 11. The wounds treated with 10 and 20% purslane herbs extract varieties C (*Portulaca grandiflora* pink flower) showed 2.88 \pm 0.51 and 0.84 \pm 1.45 mm diameter and healed on day 11. The

purslane herb A showed the highest wound healing activity, and purslane varieties A and C total flavonoid contents were 0.55 ± 0.02 and $1.58 \pm 0.02\%$ w/w, respectively.

Introduction

A wound is an injury on the outside body part tissue that disturbs physical function.^{1,2} Approximately 1% of the world's population is believed to be affected by wounds.³ Meanwhile, the prevalence of wounds in Indonesia based on the proportion of injuries is 20%.⁴ Indonesia is an archipelagic country prone to natural disasters and has many remote areas. Wound care will be a big problem for people living in remote areas with limited wound treatment access. Wounds not appropriately treated will cause disability or at least scars that cause discomfort.⁵

Wound healing at the molecular level is a complex process involving various components.⁶ These components include mediator cells and extracellular matrix. The stages of wound healing itself are divided into four stages: hemostasis, inflammation, proliferation, and remodeling.⁷ When skin gets injured, the hemostasis stage starts and follows by an increase in platelet aggregation and the production of various chemotactic factors such as TGF-B1, TGF-B2, and PDGF.⁸ Inflammatory cells such as neutrophils, lymphocytes, and macrophages at the inflammatory stage will go to the wound's center to prevent infection. In the next stage (proliferation and remodeling), keratinocytes, macrophages, platelets, and endothelial cells in the wound area release various growth factors so that new cells are formed in the wound area.⁹

Expensive and unaffordable treatment costs often hamper wound healing. So wound healing therapy at a low cost is needed at this time. Medicinal plants are sources that come from nature, so they have minimal side effects and are cheaper, so they can be a safe alternative to synthetic drugs for wound treatment.¹⁰ Several medicinal plants have been investigated for their healing properties; one is purslane herbs.

Purslane herbs (*Portulaca grandiflora*) is a plant that grows well in a tropical climate. Society mainly considers it a problematic plant. In addition, purslane herbs are only used as decorative plants in the front yard. This fact encourages the researchers to find other potencies that could be utilized from purslane herbs. Several studies have been conducted to find purslane efficacies on diseases, and one of these efficacies is antibacterial activity.¹¹ This antibacterial activity of purslane herbs can accelerate wound healing and have the potency to be developed as wound treatment topical products.

Purslane has various varieties with different active metabolite contents that need further exploration to find each variety's activity in wound healing.¹² Based on several studies, total phenol and antioxidant activity from different flower colour varieties showed different results.¹³ Flavonoid is a secondary metabolite widely known for its antioxidant properties.¹⁴ Different flower colour varieties of purslane herbs showed different antioxidant activities, suggesting they will have different flavonoid content and wound healing activity. Based on this background, research will be needed to determine purslane's various total flavonoid content and wound healing activity.

Result and discussion

Purslane herbs (Portulaca grandiflora) determination result

The first step of the research was the purslane herbs determination process. This step was performed to ensure the sample was the correct purslane (*Portulaca grandiflora*). The determination process was performed on purslane herbs A and C. The determination process was performed by direct morphology observation based on determination keys and observation reports. 18–2218–23

The observation result showed that two varieties of purslane (*Portulaca grandiflora*) have the characteristics as follows: magenta and pink colored rose-like flowers, succulent plant, 15–30 cm high stem, round and solid stem, redness brown colored stem skin with slight hair on the surface, slippery surface, upright or fall growing direction, and branching is forming from the base point characteristic.

The leaf is single thick, and meaty, 2-5 cm long, shiny green colored, shipshape spread and spiral structure, liniersubulatus or cylindrical shape, with sharp (*acutus*) tip and blunt (*obtusus*) pole, flat (*integer*) side with 1–3 cm long. The flower is bisexual, single complex, and blooms on the end of the branch with a round (*obovatus*) rose-like shape, the diameter is 25–35 mm, and the stamens are orange-colored with more than 20–50 stamen count. The flower pole is surrounded by 8–10 leaves, 2 petals, and the largest flower could reach 4 cm in diameter. The morphological observation is similar to the previous researcher's report.²² The different *Portulaca grandiflora* variety morphology that has been used in this research was on the flower color, which was magenta (variety A) and pink with white fade (variety C) color (Fig. 1).



Extraction result

The purslane herbs extraction used a simple maceration method and fit with a thermolabile active compound. Ethanol was used as an extraction solvent because it could give a higher extract yield and total phenol.¹³ Purslane herbs C showed a higher extract yield of 10.99% compared to purslane herbs A with 9.49%.

Wound healing activity test result

The rabbit skin has a similar aging process, wound healing time, and response to drugs to humans so it can be used as a wound-healing activity animal test.¹⁶ In addition, rabbits have broad back skin, making it easier to observe wounds visually. The wound healing test was started with wound induction. The hair on the back part of the rabbit was shaved with shaving tools to simplify wound induction and wounds diameter observation. Lidocaine injection was used as anesthesia because of its good local activity during the wound induction process.

The wound-healing activity test result (Table 1) based on wound diameter size on day 7 showed that purslane herbs A extract 10% concentration and purslane herbs C extract 20% concentration groups have significantly different (p < 0.05) from the negative control group. A10 also showed significant differences (p < 0.05) with other groups except for A20 and C20. The A10 group showed the smallest wound diameter average with 0.32 ± 0.55 mm size. The negative control group showed the largest wound diameter average with 6.90 ± 1.03 mm size.

Table 1

Wounds diameter on days 0, 7, 11, and 14

Channa	Wound diameter size on the day – (mm)						
Groups	0	7	11	14			
K(-)	8.00 ± 0.00	$6.90 \pm 1.03^{b,c,d,e,f}$	$4.75 \pm 0.24^{b,c,d,e,f}$	0.94 ± 0.01			
K(+)	8.00 ± 0.00	3.03 ± 0.63^{a}	0.00 ± 0.00^{a}	_			
A10	8.00 ± 0.00	$0.32 \pm 0.55^{a,b,e}$	$0.00\pm0.00^{\mathrm{a}}$				

A20	8.00 ± 0.00	1.63 ± 1.96^{a}	0.00 ± 0.00^{a}	_
C10	8.00 ± 0.00	$2.88 \pm 0.51^{a,c}$	0.00 ± 0.00^{a}	_
C20	8.00 ± 0.00	$0.84 \pm 1.45^{a,b,d}$	0.00 ± 0.00^{a}	_

Table Footnotes

^{*a*}Significantly different from K(-) (negative control).

 b Significantly different from K(+) (positive control).

^cSignificantly different from A10 (purslane herbs A extract 10% concentration).

 $^d Significantly different from A20 (purslane herbs A extract 20% concentration).$

^eSignificantly different from C10 (purslane herbs C extract 10% concentration).

^fSignificantly different from C20 (purslane herbs C extract 20% concentration).

The wound diameter observation on day 11 significantly differed between the negative control group and other groups. The negative control group showed the largest size wound diameter average with 4.75 ± 0.24 mm size. On this day, all the groups besides the negative control group showed complete wound recovery.

The wounds treated with distilled water were still not fully recovered on day 14. The wound diameter average was measured at 0.94 ± 0.01 mm. The different results showed by betadine solution and purslane herbs extract-treated groups that were fully healed since day 11.

The wound healing activity was shown by betadine solution and extract-treated groups. The activity was shown by wound diameter changes that were smaller daily and healed starting from day 11 to fully recover on day 14 (Fig. 2).

Fig. 2



The different healing process was shown by the negative control group that was treated with distilled water. The wounds showed slow healing progress. The healing progressed until the end of observation, and the wounds were still not fully closed or recovered on day 14. Unfortunately, the observation was not prolonged after day 14, so the real healing timing from the negative control group was never established. However, the wounds would be healed naturally on days 13–16, depending on the size.¹⁶

The betadine solution-treated wounds showed better healing results compared to the untreated wounds. Betadine solution contains povidone-iodine 10%, an antibacterial compound widely used as first aid to treat wounds. Povidone-iodine is an iodoform compound that penetrates inside bacteria cells and oxidizes key protein, nucleotide, and fat acid bacteria leading to bacteria death.²⁴ Without bacteria around, the wound will be healed faster because the inflammation stage of the healing process will end faster. The proliferation and remodeling will be achieved even faster rather than untreated wounds. The inflammation process is needed to transfer neutrophils, monocytes, and lymphocytes to the wound site to kill invading bacteria. This process will be prolonged if bacteria are too many to handle, so antibacterial compounds such as povidone-iodine are needed to accelerate the healing process.²⁵

The wound diameter on all purslane herbs extracts groups showed a smaller size daily and fully recovered on day 14, similar to the betadine solution-treated group. The different concentration dosages on purslane herbs A and C extract groups showed different diameter sizes even though the differences were insignificant (p > 0.05). It may be concluded

that different concentration dosages did not give different wound healing activities. However, the dosage variation in this research is only 10% and 20%, and the effect of broader and higher dosage concentrations on wound healing activity remains unknown.

Qualitative phytochemical screening result

Phytochemical screening analysis showed that purslane herbs variety A and C extract showed similar compounds: alkaloids, flavonoids, saponins, and tannins. Alkaloid reaction test results showed brown sediment. Furthermore, on saponin reaction test showed persistent foam on HCl addition. The color-changing to dark green showed all extracts after FeCl₃ addition showed tannins content. Brown sediment was shown after chloroform, and H_2SO_4 concentrated addition showed terpenoid contains. The result was comparable with other studies, which showed that ethanolic purslane herbs extract contains flavonoids, alkaloids, terpenoids, saponins, and tannins²⁶ (Table 2).

Table 2

Purslane herbs of various varieties extract phytochemical screening result

Extract	Flavonoid	Alkaloid	Terpenoid	Saponin	Tannin
Purslane herbs A	+	+	+	+	+
Purslane herbs C	+	+	+	+	+

Total flavonoid content analysis result

The calibration curve determination result gives the formula $y = 00\,233 + 002\,77x$ with a 0.9971*r* value. The higher flavonoid content was shown by purslane herbs C extract with 1.58 \pm 0.02%. The purslane herbs A extract showed smaller flavonoid content with 0.55 \pm 0.02%. Those flavonoid contents were significantly different, with a *p*-value < 0.05. This result showed that different varieties contain different metabolite content. This is comparable with other research that showed different flavonoid content in the various purslane varieties.¹²

Discussion

The determination result showed that purslane herb A is *Portulaca grandiflora* with a magenta-colored flower variety. Moreover, purslane herb C is *Portulaca grandiflora* with a pink-colored flower variety. These differences are related to secondary metabolites compound content in the purslane herbs.¹² Those purslane varieties have shown antibacterial activity on *Staphylococcus aureus*, *Escherichia coli*, and *Pseudomonas aeruginosa*.¹⁵ Those bacteria are commonly found in wound infections, so this extract has the potency to be developed as a wound-healing compound.

A wound is an injury that could be acquired on every body part. The open wound on the skin will increase bacteria infection risk because the skin contains many bacteria. This open wound condition will increase bacteria's chance to enter and invade the deeper tissue.¹ The body's inflammation mechanism response will follow this bacteria invasion. The pro-inflammation agents such as neutrophils, macrophages, and lymphocytes will migrate into wound site and kill the bacteria that is invading with various mechanisms such as phagocytes and apoptosis.²⁵ This process will be prolonged if bacteria are too many to handle, so an active compound with antibacterial activity is necessary to shorten this stage. The inflammation stage will also be prolonged by oxidative stress caused by free radical excessive accumulation.²⁷ This condition will cause further cell obstruction, prolonging the wound-healing process. That is why the active compound with antioxidant activity also accelerates the wound healing process.

Based on phytochemical screening results, ethanolic purslane herbs extract contains various secondary metabolites such as flavonoids, alkaloids, saponin, tannins, and terpenoids that have taken part in wound healing activity with various mechanisms²⁸ that were shown by wound healing activity results. The wounds treated with purslane herbs extract A and C healed with no significantly different from betadine solution-treated wounds, especially on days 11 and 14. At the end of observation, all purslane extract-treated wounds showed a similar healed stage to betadine solution-treated

wounds. Different purslane species (*Portulaca oleracea*) grown in Jordania showed similar results with different animal tests.²⁹

Flavonoid content in purslane herb extracts is known for its antiinflammation, antibacterial, and antioxidant activity. ^{30,31} As an antiinflammation, flavonoids shorten the inflammation stage, leading to the proliferation and remodeling stage, shortening the healing process. The antibacterial activity of flavonoids supports this activity. The inflammation process ends faster in line with the bacteria number decline. Flavonoid antibacterial activity refers to its ability to complex forming with bacteria cell membrane protein.³² Flavonoids work as an antioxidant by scavenging free radicals that lead to oxidative stress prevention.²⁷ Free radical is a reactive molecule with antibacterial properties in a reactive oxygen species (ROS) form.

Nevertheless, in a saturated condition, ROS could cause further tissue damage at the wound site.^{33,34} Thus condition prolonged the healing process, so the antioxidant activity of flavonoids in purslane herb extracts has the benefit that could prevent oxidative stress from happening.^{14,34,35} Flavonoids could donate their electrons to ROS, preventing ROS from taking electrons from molecules to tissue formation, such as DNA, protein, and lipids.³³

Total flavonoid content analysis showed that purslane herb extracts of both varieties showed flavonoid content even though this result did not align with the wound healing activity test. Purslane herb variety A extract showed higher wound healing activity than variety C extract, even though its flavonoid content was lower than purslane herb variety C extract. This may be caused by differences in other metabolite constituents in both purslane herb varieties.

Another constituent in the purslane herb suspected of its role in wound healing is an alkaloid. Alkaloids may accelerate wound healing and tissue granulation by increasing collagen and fibroblast and lowering inflammation cell numbers.³⁶ Fibroblast is a connective tissue that plays in collagen production.³⁷ Collagen is a connective protein responsible for elasticity and skin structure strength.^{38,39} Alkaloids also play a role in SRC/MEK/ERK signaling pathway regulation that promotes the wound healing process.⁴⁰

Saponin detected in purslane herb A, and C varieties is a secondary metabolite with antibacterial activity. This activity is related to its ability to increase bacteria cell membrane permeability, leading to hemolysis.⁴¹ Other than that, saponin may also increase collagen production and accelerate cell epithelization in the wound.⁴² Saponin also had an essential role in high glucose-induced wound healing by increasing endothelial cell proliferation.⁴³

Tannins may also play an essential role in the wound-healing process. Tannins are known for their antibacterial activity in rebuilding damaged tissue and accelerating fibrous tissue contraction in wound healing.⁴⁴ Fibrous contraction by fibroblast will trigger scar formation that protects cell formation in the wound site. Tannin's antibacterial activity may be caused by its toxicity to the microorganism related to bacteria enzyme and substrate complex formation.⁴⁵ Other than this mechanism, tannins are also known for their antioxidant with similar free radical scavenging mechanism activity to flavonoids.^{46,47}

Experimental section

Materials

Tools to conduct *in vivo* wound healing activity tests were scissors and razors to cut the rabbit hair, 1 ml disposable syringe, biopsy punch 8 mm diameter, anatomical forceps, scalpel blade, vernier calipers to measure wound diameter, and vacuum rotary evaporator. Analytical balance, spectrophotometer UV-vis, and glassware were used to perform phytochemical and total flavonoid content analysis.

The sample of two varieties of purslane was obtained from Madiun City of Java Island in Indonesia. The solvent that was used in the extraction process was ethanol 70%. The New Zealand rabbit was used as an animal test to perform wound healing activity. Lidocaine injection was used as a local anesthetic on rabbit skin when wound excision was conducted. Betadine solution and distilled water were used as the positive and negative control, respectively. The

materials used for phytochemical screening and total flavonoid content analysis were ethanol 96% pro analysis, AlCl₃, potassium acetate, and distilled water.

Methods

Purslane herbs characterization (Portulaca grandiflora)

Two kinds of purslane herbs obtained from Madiun City were characterized to know their variety. Determination was conducted in Biology Laboratory, Biology Department, Widya Mandala Surabaya Catholic University.

Extraction process

Two hundred grams of each dried purslane herbs variety were macerated with 500 ml ethanol 70% for 5 days. The solid residue from the maceration process was macerated once more with the same amount of solvent. The first and second macerates were mixed and evaporated with a vacuum rotary evaporator at 50 °C until a thick extract was obtained. The thick extract was used for phytochemical screening, total flavonoid content analysis, and *in vivo* wound healing activity tests.

Wound healing activity test

The rabbit was used as an animal test to conduct *in vivo* wound healing activity. The New Zealand rabbit used in this study was 3–5 months old and had a 1.5–2.5 kg weight. Ethical clearance was obtained from Health Research Ethics Committee Faculty of Medicine, Muhammadiyah Surakarta University, Indonesia.

Rabbit's hair on the back part was shaved with shaving tools. The rabbit skin was anesthetized with lidocaine with 0.0125–0.7600 ml per kg BW dose until showing unresponsive when stabbed with a needle. A biopsy punch of 8 mm diameter was used to make wound excision on rabbit skin. Three rabbits were induced with six wounds for each rabbit.

The wounds on the rabbit's back skin were grouped into 6 treatment groups such as negative control (distilled water), positive control (betadine solution), 10 and 20% purslane herbs extract varieties A, 10 and 20% purslane herbs extract varieties C.

The wounds were cleaned and disinfected with an antiseptic solution and waited until the blood stopped before the treatment was applied twice daily for two weeks. The macroscopic condition observation and wound diameter measurement was conducted on days 0, 7, 11, and 14.

Qualitative phytochemical screening

The flavonoid identification was conducted by dissolving 0.5 grams of each various variety of purslane ethanolic extract in the distilled water and added with magnesium and 5 drops of HCl 2 N. The mixture was heated for 5–10 minutes. After filtration and cooling, the filtrate was added with amyl alcohol and shaken vigorously. The reaction result is positive if red color is formed in the amyl alcohol layer. $\frac{17}{12}$

The alkaloid identification was conducted by 0.5 grams of each various variety of purslane ethanolic extract was added with 1 ml ammonia and then added with chloroform. The mixture was filtrated, and the filtrate was placed in the test tube and added with 2 N HCl. The mixture was shaken and waited until separation occurred. The filtrate was divided into three parts in the different test tubes. One drop of Dragendorff reagent was added to the first filtrate. The reaction result is positive if brown sediment is formed. One drop of Mayer reagent was added to the second filtrate, and the reaction was positive if white sediment formed. The third filtrate is blank or negative control. $\frac{17}{2}$

The saponin identification was conducted by placing 0.5 grams of each various variety of purslane ethanolic extract into a test tube and then dissolved with hot distilled water. After the mixture was cold, it was shaken for 10 seconds. The reaction is positive if 1–10 cm persistent foam is formed for 10 minutes, and stays still after adding 1–2 drops of $2 \text{ N HCL} \frac{17}{2}$

The terpenoid identification was conducted by placing 0.5 grams of each various variety of purslane ethanolic extract into a test tube and then added with chloroform and H_2SO_4 concentrate. The reaction is positive if brown sediment is formed.

The tannin identification was conducted by placing 0.5 grams of each various variety of purslane ethanolic extract into a test tube and then added with 1% FeCl₃ solvent. The reaction is positive if green or dark blue color is formed.

The total flavonoid content of various varieties of purslane was measured with $AlCl_3$ colorimetric methods. Twentyfive mg of quercetin was dissolved with 25 ml ethanol in a 25 ml volumetric flask to obtain a 1000 µg ml⁻¹ initial solution. Standard solutions of 10, 20, 40, 60, 80, and 100 ppm were made from the initial solution. Each standard solution was pipetted 0.5 ml and then was added with 1.5 ml ethanol 96%, 1 ml 10% AlCl₃, 1 ml potassium acetate 1 M, and 2.8 ml distilled water. The solution was incubated at 25 °C for 30 minutes. The absorbance was measured with a spectrophotometer UV-vis on a wavelength (λ) of 434.2 nm. The absorbance reading result was used to make a calibration curve of absorbance *vs.* quercetin concentration.

Five grams of each various variety of purslane ethanolic extract was dissolved with 25 ml ethanol and stirred at 200 rpm, and then filtrated. The filtrate was added to 25 ml ethanol. The blank solution for the flavonoid content assay was made from 0.5 ml ethanol that was added with 1.5 ml ethanol 96%, 1 ml 10% AlCl₃, 1 ml potassium acetate 1 M, and 2.8 ml distilled water. One ml of various purslane ethanolic extract was added with ethanol until the marking sign of the 10 ml volumetric flask. Half ml of the solution was pipetted and added with 1.5 ml ethanol 96%, 1 ml 10% AlCl₃, 1 ml potassium acetate 1 M, and 2.8 ml distilled water. The mixture was incubated at 25 °C for 30 minutes, and the absorbance was read with a spectrophotometer UV-vis on λ 434.2 nm. The total flavonoid content was counted with the following equation:

 $F = (c \times V \times f \times 10 - 6)/m \times 100\%$

Note: F = flavonoid content, c = quercetin equality (g ml⁻¹), V = total extract volume, f = dilution factor, m = sample weight (g).

Data analysis

The observed data were statistically analyzed with a 95% (p < 0.05) confidence limit using the Kruskal–Wallis analysis test.

Live subject statement

All animal procedures were performed under the guidelines for care and use of laboratory animals of Widya Mandala Surabaya Catholic University, Indonesia, and approved by the Health Research Ethics Committee Faculty of Medicine, Muhammadiyah Surakarta University, Indonesia.

Conclusions

Two varieties of purslane herbs (magenta and pink flower) showed different wound healing activities, especially on day 7. The wound healing activity test of purslane herbs A (*Portulaca grandiflora* magenta flower variety) 10 and 20% concentration showed 0.32 ± 0.55 and 1.63 ± 1.96 mm, respectively, wound diameter size. The purslane herbs C (*Portulaca grandiflora* pink flower variety) 10 and 20% concentration showed 2.88 ± 0.51 and 0.84 ± 1.45 mm, respectively, wound diameter size. The purslane herbs A (*Portulaca grandiflora* magenta flower variety) 10 and 20% concentrations showed 1.63 ± 1.96 mm, respectively, wound diameter size. The purslane herbs A (*Portulaca grandiflora* magenta flower variety) 10 and 20% concentrations showed higher activity in the wound healing process on day 7 with 0.32 ± 0.55 and 1.63 ± 1.96 mm, respectively, which is the smallest wound diameter. Purslane herb varieties A and C total flavonoid contents were 0.55 ± 0.02 and $1.58 \pm 0.02\%$ w/w, respectively.

Author contributions

Experimental work (AB, LP, AP, EDC), data analysis (AB and CEP), and writing the manuscript (AB and AP).

Conflicts of interest

There are no conflicts to declare.

References

1 S. A. Price and L. M. Wilson, Patofisiologi: Konsep Klinis Proses-proses Penyakit, 6th edn, 2006

2 E. Naseri and A. Ahmadi, Eur. Polym. J., 2022, 173, 111293.

- 3 P. Grover, K. Khanna, A. Bhatnagar and J. Purkayastha, Biomed. Pharmacother., 2021, 140, 111797.
- 4 The Ministry of Health Republic of Indonesia, Laporan Nasional Riskesdas, 2018
- 5 L. Zhang, Y. Tai, X. Liu, Y. Liu, C. Yang, D. Kong, C. Qi, S. Wang and A. C. Midgley, *Appl. Mater. Today*, 2021, **25**, 101186.
- 6 S. Patel, S. Srivastava, M. R. Singh and D. Singh, Biomed. Pharmacother., 2019, 112, 108615.
- 7 J. S. Low, K. K. Mak, S. Zhang, M. R. Pichika, P. Marappan, K. Mohandas and M. K. Balijepali, *Fitoterapia*, 2021, **154**, 105026.
- 8 D. H. Assar, N. Elhabashi, A. A. A. Mokhbatly, A. E. Ragab, Z. I. Elbialy, S. A. Rizk, A. E. Albalawi, N. A. Althobaiti, S. A. Jaouni and A. Atiba, *Biomed. Pharmacother.*, 2021, **143**, 112151.
- 9 M. E. Okur, I. D. Karantas, Z. Şenyiğit, Ü. N. Okur and P. I. Siafaka, *Asian J. Pharm. Sci.*, 2020, **15**, 661.
- 10 I. Mssillou, M. Bakour, M. Slighoua, H. Laaroussi, H. Saghrouchni, F. E. Amrati, B. Lyoussi and E. Derwich, J. Ethnopharmacol., 2022, 298, 115663.
- 11 K. P. Fung, Q. B. Han, M. Ip, X. S. Yang, C. Bs. Lau and B. Cl. Chan, *Hong Kong Med. J.*, 2017, 23, 38.
- 12 Md A. Alam, A. S. Juraimi, M. Y. Rafii, A. Abdul Hamid, F. Aslani, M. M. Hasan, M. A. M. Zainudin and Md. K. Uddin, *BioMed Res. Int.*, 2014, <u>2014</u>, <u>6296063</u>.
- 13 C. K. Lim, W. N. Tiong and J. L. Loo, Int. J. Phytopharm., 2014, 4, 1.
- 14 M. T. B. Carvalho, H. G. Araújo-Filho, A. S. Barreto, L. J. Quintans-Júnior, J. S. S. Quintans and R. S. S. Barreto, *Phytomedicine*, 2021, 90, 153636.
- 15 A. Purwanto, Agri-Tek: Jurnal Ilmu Pertanian, Kehutanan dan Agroteknologi, 2021, 22, 1.
- 16 A. Grada, J. Mervis and V. Falanga, J. Invest. Dermatol., 2018, 138, 2095.
- 17 The Ministry of Health Republic of Indonesia, Materia Medika Indonesia V, 1989
- 18 C. G. G. J. Van Steenis, Flora, 2008
- 19 M. K. Uddin, A. S. Juraimi, Md. S. Hossain, M. A. U. Nahar, Md. E. Ali and M. M. Rahman, Sci. World J., 2014, 2014, 951019.

20 H. Setyowati, Cont. Prof. Dev., 2017, 44, 818.

21 E. Djauhariya and <u>H.</u>Hernani, <u>Gulma Berkhasiat Obat, Gulma Berkhasiat Obat</u>, 2004

22 F. I. D. Setiawan, S. I. Aisyah and K. Krisantini, Journal of Tropical Crop Science, 2016, 3, 67.

23

T. Setyawati, S. Narulita, I. P. Bahri and G. T. Raharjo, A Guide Book to Invasive Plant Species in Indonesia, 2015

24 D. Lepelletier, J. Y. Maillard, B. Pozzetto and A. Simon, Antimicrob. Agents Chemother, 2020, 64, 1.

25 S. Guo and L. A. Di Pietro, J. Dent. Res., 2010, 89, 219.

26 R. Londonkar and H. Nayaka, J. Pharma Res., 2011, 4, 3553.

- 27 S. D. Fitzmaurice, R. K. Sivamani and R. R. Isseroff, Skin Pharmacol. Physiol., 2011, 24, 113.
- 28 L. C. Adiele, R. C. Adiele and J. C. Enye, Asian Pac. J. Trop. Med., 2014, 7, 620.
- 29 A. N. Rashed, F. U. Afifi and A. M. Disi, J. Ethnopharmacol., 2003, 88, 131.
- 30 Y. X. Zhou, H. L. Xin, K. Rahman, S. J. Wang, C. Peng and H. Zhang, *BioMed Res. Int.*, 2015, 2015, 925631.
- 31 M. H. Masoodi, B. Ahmad, S. R. Mir, B. A. Zargar and N. Tabasum, J. Pharm. Res., 2011, 44, 3044.
- 32 A. Royani, M. Hanafi, H. Julistiono and A. Manaf, *Mater. Today: Proc.*, <u>2023</u>, <u>72</u>, <u>6</u>, <u>2022</u><u>2796</u>_ <u>2802</u>.
- 33 C. Dunnill, T. Patton, J. Brennan, J. Barrett, M. Dryden, J. Cooke, D. Leaper and N. T. Georgopoulos, *Int. Wound J.*, 2017, 14, 89.
- 34 A. Priyadarshi, G. K. Keshri and A. Gupta, *Phytomedicine Plus*, 2022, 2, 100292.
- 35 E. Asante-Kwatia, A. Evelyn, J. Silas, G. Yakubu, G. A. H. Lord, G. Adjei-Hinneh, I. K. Amponsah and A. Y. Mensah, *Heliyon*, 2021, 7, e08261.
- 36 T. Nagappan, T. C. Segaran, M. E. A. Wahid, P. Ramasamy and C. S. Vairappan, *Molecules*, 2012, 17, 14449.
- 37 L. Bao, X. Cai, M. Zhang, Y. Xiao, J. Jin, T. Qin and Y. Li, J. Funct. Foods, 2022, 90, 104981.
- 38 P. Bainbridge, J. Wound Care, 2013, 22, 407.
- 39 S. Sharma, V. K. Rai, R. K. Narang and T. S. Markandeywar, Life Sci., 2022, 290, 120096.
- 40 X. Q. Shi, G. Chen, J. Q. Tan, Z. Li, S. M. Chen, J. H. He, L. Zhang and H. X. Xu, J. *Ethnopharmacol.*, 2022, **295**, 115396.
- 41 M. P. Wei, H. Yu, Y. H. Guo, Y. L. Cheng, Y. F. Xie and W. R. Yao, *Food Control*, 2021, **130**, 108337.
- 42 D. MacKay and A. L. Miller, Alternative Med. Rev., 2003, 8, 359.
- 43 T. Lei, Y. Gao, Y. Duan, C. Cui, L. Zhang and M. Si, Environ. Toxicol., 2022, 37, 35385194.
- 44 A. Santana, A. Barros, H. Oliveira and I. Victor, Biomed. Pharmacother., 2017, 96, 182.
- 45 A. Scalbert, Phytochemistry, 1991, 30, 3875.
- 46 M. Heinonen, Mol. Nutr. Food Res., 2007, 51, 684.
- 47 C. M. González, E. Llorca, A. Quiles, I. Hernando and G. Moraga, *LWT–Food Sci. Technol.*, 2022, 155, <u>112961</u>.

Queries and Answers

Q1

Query: Have all of the funders of your work been fully and accurately acknowledged?

Answer: Yes

Q2

Query: Have all of the author names been spelled and formatted correctly? Names will be indexed and cited as shown on the proof, so these must be correct. No late corrections can be made. Answer: Yes

Q3

Query: Do you wish to add an e-mail address for the corresponding author? If so, please provide the relevant information. Answer: antonius.budiawan@ukwms.ac.id

Q4

Query: Please check that ref. 9 has been displayed correctly. **Answer:** Ref. 9 has been displayed correctly

Q5

Query: Ref. 12: Please provide the page (or article) number(s).

Answer: The volume and article number has been updated and added.

Md A. Alam, A. S. Juraimi, M. Y. Rafii, A. Abdul Hamid, F. Aslani, M. M. Hasan, M. A. M. Zainudin and Md. K. Uddin, *BioMed Res. Int.*, 2014, **2014**, **6**296063.

Q6

Query: Please indicate where ref. 17 should be cited in the text.

Answer: Ref. 17 cited in the section "Qualitative phytochemical screening" paragraphs 1-3.

Q7

Query: Ref. 21: Please provide the initial(s) for the 2nd author.

Answer: The 2nd author's initials have been added.

E. Djauhariya and H. Hernani, Gulma Berkhasiat Obat, Gulma Berkhasiat Obat, 2004

Q8

Query: Ref. 22: Please provide the initial(s) for the 3rd author.

Answer: The 3rd author's initials have been added.

F. I. D. Setiawan, S. I. Aisyah and K. Krisantini, Journal of Tropical Crop Science, 2016, 3, 67.

Q9

Query: Please indicate where ref. 23 should be cited in the text.

Answer: Ref. 23 cited in the section "Results of Determination of Purslane (Portulaca grandiflora)".

The determination process was performed by direct morphology observation based on determination keys and observation reports.^{18–23}

Q10

Query: Ref. 32: Please provide the page (or article) number(s).

Answer: Article numbers and pages have been added. The year of the article has also been updated

A. Royani, M. Hanafi, H. Julistiono and A. Manaf, Mater. Today: Proc., 2023, 72, 6, 20222796-2802.

Q11

Query: Ref. 47: Please provide the page (or article) number(s). **Answer:** The article number have been added. Please check ref. 47 10. Corrections received (24-3-2023)



Corrections received for article d3ra00868a

1 message

noreply@proof-central.com <noreply@proof-central.com> To: antonius.budiawan@ukwms.ac.id Fri, Mar 24, 2023 at 3:58 PM

This is an automatically generated message. Please do not reply because this mailbox is not monitored.

Dear Antonius Budiawan,

Thank you for using Proof Central for your RSC Advances article d3ra00868a "(Wound Healing Activity and Flavonoids Contents of Purslane (*Portulaca grandiflora*) Various Variety)". We have attached the PDF summary of your corrections for your reference.

Please note that now we have received your corrections, the final version of your article is being prepared and further changes are no longer possible.

All your corrections have been saved in our system. The PDF summary of your corrections, generated from Proof Central, can be downloaded from the following site for your reference: https://pcv3-rsc-live.s3.amazonaws.com/37c6c94c2fd54ab0bbf75d49eb92b3/RA_d3ra00868a_edit_report.pdf

With best wishes, From the Editors of RSC Advances Royal Society of Chemistry advances@rsc.org

This is an automatically generated message. Please do not reply because this mailbox is not monitored.

11. Article published (28-3-2023)



Your RSC Advances article page numbers are now available

1 message

Follow us and stay in touch	諧 in

Dear Mr Antonius Budiawan

Wound Healing Activity and Flavonoids Contents of Purslane (<i>Portulaca grandiflora</i>) Various Variety

We have published your RSC Advances article in an issue on pages 9871 - 9877.

It can now be cited as:

RSC Advances, 2023, 13, 9871 - 9877

By publishing your article in RSC Advances, you are supporting the Royal Society of Chemistry to help the chemical science community make the world a better place. All corresponding authors who are not already members of the Royal Society of Chemistry are entitled to one year's Affiliate membership free of charge. If you would like to find out more please email membership@rsc.org, including the promo code OA100 in your message. Learn all about our member benefits here.

Interested in publishing with us again? Why not check out our Calls for papers to submit to one of our open collections.

Access your PDF reprints here

Manuscript ID: D3RA00868A

Password: 992927 (this link will expire in 60 days)

You can now access the enhanced HTML version of your article here

Want to enhance the visibility of your article? Find out more about our article promotion services here.

If you'd like to learn more about the other post-publication author benefits available to you, such as grants and memberships, please visit our Author Hub.

We look forward to publishing your next article with RSC Advances or another Royal Society of Chemistry journal soon.

With best wishes, From the Editors of RSC Advances Royal Society of Chemistry advances@rsc.org

You have been sent this message because you have submitted an article to RSC Advances. If you feel that you have received this in error, please contact advances@rsc.org

The Royal Society of Chemistry, Thomas Graham House, Science Park, Cambridge CB4 0WF, United Kingdom. Registered charity number: 207890.

© Royal Society of Chemistry 2023. All rights reserved.

DISCLAIMER:

This communication is from The Royal Society of Chemistry, a company incorporated in England by Royal Charter (registered number RC000524) and a charity registered in England and Wales (charity number 207890). Registered office: Burlington House, Piccadilly, London W1J 0BA. Telephone: +44 (0) 20 7437 8656.

The content of this communication (including any attachments) is confidential, and may be privileged or contain copyright material. It may not be relied upon or disclosed to any person other than the intended recipient(s) without the consent of The Royal Society of Chemistry. If you are not the intended recipient(s), please (1) notify us immediately by replying to this email, (2) delete all copies from your system, and (3) note that disclosure, distribution, copying or use of this communication is strictly prohibited.

Any advice given by The Royal Society of Chemistry has been carefully formulated but is based on the information available to it. The Royal Society of Chemistry cannot be held responsible for accuracy or completeness of this communication or any attachment. Any views or opinions presented in this email are solely those of the author and do not represent those of The Royal Society of Chemistry. The views expressed in this communication are personal to the sender and unless specifically stated, this e-mail does not constitute any part of an offer or contract. The Royal Society of Chemistry shall not be liable for any resulting damage or loss as a result of the use of this email and/or attachments, or for the consequences of any actions taken on the basis of the information provided. The Royal Society of Chemistry does not warrant that its emails or attachments are Virus-free; The Royal Society of Chemistry has taken reasonable precautions to ensure that no viruses are contained in this email, but does not accept any responsibility once this email has been transmitted. Please rely on your own screening of electronic communication.

More information on The Royal Society of Chemistry can be found on our website: www.rsc.org