Redefining the hepatoprotective potential of Javanese turmeric (Curcuma xanthorrhiza) Kombucha towards the diethylnitrosamine-induced hepatotoxicity of mice

by Ignatius Srianta

Submission date: 12-Jul-2024 01:07PM (UTC+0700) Submission ID: 2415588519 File name: owards_the_diethylnitrosamine-induced_hepatotoxicity_of_mice.pdf (2.17M) Word count: 5862 Character count: 32037



WestminsterResearch http://www.westminster.ac.uk/westminsterresearch

Redefining the hepatoprotective potential of Javanese turmeric (Curcuma xanthorrhiza) Kombucha towards the diethyInitrosamine-induced hepatotoxicity of mice Elok Zubaidah, Eirene Charista Dea, Ella Saparianti, Rhytia Ayu Christianty Putri, Hidayat Sujuti, Ignatius Srianta, Laura Godelive and Ihab Tewfik

This article is © Emerald Publishing Limited and permission has been granted for this version to appear here: <u>http://westminsterresearch.westminster.ac.uk/</u>

The final, published version in Nutrition & Food Science, DOI: 10.1108/NFS-09-2023-0219, 2024 is available at:

https://doi.org/10.1108/NFS-09-2023-0219

This manuscript version is made available under the CC BY-NC 4.0 licence <u>https://creativecommons.org/licenses/by-nc/4.0/</u>

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

Redefining the hepatoprotective potential of Javanese turmeric (*Curcuma xanthorrhiza*)

Kombucha towards the diethylnitrosamine-induced hepatotoxicity of mice

Elok Zubaidah, Eirene Charista Dea, Ella Saparianti and Rhytia Ayu Christianty Putri Department of Food Science and Technology, Faculty of Agricultural Technology, Brawijaya University, Malang, Indonesia

Hidayat Sujuti Department of Biomedical, Faculty of Medicine, Brawijaya University, Malang, Indonesia

Ignatius Srianta and Laura Godelive Department of Food Technology, Widya Mandala Catholic University Surabaya, Surabaya, Indonesia, and

Ihab Tewfik

School of Life Sciences, University of Westminster, London, UK

Abstract

Purpose – This research intended the utilization of Javanese turmeric (0.4% w/v) as a kombucha substrate and analysis of its hepatoprotective activity, in comparison against nonfermented Javanese turmeric beverage (JTB) and black tea kombucha.

Design/methodology/approach – Forty-two healthy male Balb/c mice (two- to three-week-old, 20–30 g) were divided into six groups with seven replicates each. The treatments were normal diet, normal diet þ Javanese turmeric kombucha (JTK), normal diet þ diethylnitrosamine (DEN), DEN þ JTB, DEN þ JTK, DEN þ black tea kombucha. Kombuchas and JTB were given at 0.3 mL/20 g BW/d. DEN was induced intraperitoneally at a dose of 100 mg/kg. Observed biomarkers were blood serum glutamate pyruvate transaminase (SGPT) and serum glutamate oxaloacetate transaminase (SGOT) activity, serum malonaldehyde (MDA), as well as liver histology. Data were analyzed using analysis of variance.

Findings – Among DEN-induced groups, JTK significantly (p < 0.05) diminished the level of blood SGPT, SGOT and serum MDA.JTK also had lower blood SGPT (8.604 6 2.195 U/L) and serum MDA levels

1. Introduction

Chronic liver disease occurs after the liver is exposed to prolonged inflammation. This inflammation can be caused by oxidative stress, alcohol, viral infection, accumulation of cytoplasmic fats or autoimmune disease. Chronic liver disease may alter several biochemical pathways related to nutrients and result in great nutritional impact (Fortes, 2017). Patients with early stages could undergo transplantation, abrasion or resection. However, symptoms may be hard to detect during the early stages and can lead to delayed treatment (Zhang *et al.*, 2021). Therefore, prevention is needed at the early stages to ease the burden and arrest the expansion of the disease.

Diet and dietary constituents seem to play a hepatoprotective role (Elsharkawy *et al.*, 2022). Javanese turmeric is an Indonesian local rhizome that is mostly used to alleviate liver disease (Rahmat *et al.*, 2021). It contained xanthorrhizol, curcuminoid, flavonoid, arturmerone, *a*-turmerone, curcumene, bisacurone, curlone, lactone-germacrone and germacrone (Lukitaningsih *et al.*, 2020). Xanthorrhizol, the most abundant active compound found in Javanese turmeric, is known as a good anti-inflammation compound and exhibits the ability to reduce the activity of blood serum glutamate pyruvate transaminase (SGPT) and serum glutamate oxaloacetate transaminase (SGOT), enzymes that are considered as liver damage markers (Oon *et al.*, 2015). However, these identified bioactive compounds are bound with the plant matrix and other components, therefore they have low bioavailability (Rahmat *et al.*, 2021). Owing to the fermentation process many of these bioactive compounds could be released and become readily bioavailable.

Kombucha is a tea beverage fermented by a symbiotic association of bacteria and yeast (SCOBY) (Miranda *et al.*, 2022). The bacteria and yeast in kombucha are known to produce hepatoprotective components such as glucuronic acid and *D*-saccharic acid 1,4-lactone (DSL). Glucuronic acid can enhance the detoxification of toxicants, increase polyphenol bioavailability and increase the solubility of steroids and vitamin D, whereas DSL may act as a hepatoprotective agent and antioxidant (Martínez-Leal *et al.*, 2018). Recent research on kombucha has focused on using other new materials as substrates (Salafzoon *et al.*, 2018; Zubaidah *et al.*, 2018; Zubaidah *et al.*, 2022). Thus, the objective of this research was to analyze the usage of Javanese turmeric as a kombucha substrate to enhance its hepatoprotective properties. The microbial activity during fermentation was expected to synergize with Javanese turmeric active compounds and enhance their hepatoprotective activity.

2. Materials and methods

2.1 Materials

Javanese turmeric of commercial maturity, commercial kombucha starter, black tea and cane sugar were obtained in Malang, East Java, Indonesia. Javanese turmeric was peeled, cleaned and cut into thin slices. The slices were then dried in a cabinet drier at 60°C for 5–6h and ground into a fine powder. Diethylnitrosamine (DEN) was obtained from the Tokyo Chemical Industry.

2.2 Kombucha and nonfermented solution preparation and analysis

Javanese turmeric solution (0.4% w/v) was made with 2 g of Javanese turmeric powder that was placed into teabags. The concentration was chosen as the optimal treatment after the analysis of total microbial cells, total titratable acidity, pH, total phenolic content and antioxidant activity by Zubaidah *et al.* (2022). Javanese turmeric powder was extracted in boiling water (500 mL) for 5 min, then sweetened with 10% sugar. The solution was put into sterile glass jars and cooled to room temperature (at 30°C). Kombucha starter (10% v/v) was

added aseptically. The jars were covered with sterile cheese cloths and fermentation was carried out at room temperature for 12 days. As a control, black tea kombucha was prepared with the same method at a concentration of 0.4% (Zubaidah *et al.*, 2018). Nonfermented Javanese turmeric beverage (JTB) was prepared with the same concentration and procedure as Javanese turmeric kombucha (JTK) without the addition of a kombucha starter.

2.3 Animal experiment and analysis

Forty-two healthy male BalB/C mice (two- to three-week-old, 20–30 g) were obtained from Biomedic Laboratory, Medical Faculty of Muhammadiyah University, Malang. The mice were divided randomly into six groups with seven replicates each, with Group 1 (normal diet), Group 2 (normal diet þ JTK), Group 3 (normal diet þ DEN), Group 4 (DEN þ JTB), Group 5 (DEN þ JTK) and Group 6 (DEN þ black tea kombucha). Each group shared one housing and the bedding was changed every three days. Mice received exposure to light for 12 h/day. Mice were acclimated for seven days, weighed once every three days, and were given JTK, black tea kombucha and JTB (0.3 mL/20 g BW/day) regularly for three weeks. The mice accessed standard diet (BR2 commercial feed) and water ad libitum throughout the experiment.

Alongside the treatments, DEN (100 mg/kg) was then induced intraperitoneally once per week for two weeks. Mice were then incubated for one week while the treatments were still being given. The mice were then fasted for one day. Twenty-four hours after fasting, the mice were sacrificed using 0.2 mL ketamine injection (0.1 mg/g BW). Blood was withdrawn from the heart during scarification for SGPT and SGOT activity analysis. SGPT and SGOT activity were analyzed using commercial reagent kits (Biosystems S.A, Costa Brava 30, Spain Quality system certified: EN ISO 13485 and EN ISO 9001 standard). Malonaldehyde (MDA; nM/mL) concentration was analyzed in blood serum using the standard spectrophotometric method (Zubaidah *et al.*, 2023) with an MDA kit (Biotect Co, China, ISO 9001:2015, EU2577). The animal study was approved by the Brawijaya University Research Ethics Committee (Ethical Clearance No. 109-KEP-UB-2021).

2.4 Histopathological analysis

Histopathological analysis of liver cells was carried out according to Zubaidah *et al.* (2023). After scarification, livers were taken and fixed in formalin, then dehydrated by 50%-100% ethanol. The livers were dissected, fixed in 5 mm paraffin and stained using hematoxylin and eosin. The stained dissections were observed through a light microscope (40x, 100x and 400x) to analyze the existence of inflammatory cells and apoptotic bodies. Damaged cells were counted.

2.5 Statistical analysis

Data are presented as mean 6 standard deviation (SD) ($n \sqrt{47}$). The data were analyzed by analysis of variance, followed by Fisher's exact test at p < 0.05.

Result and discussion

3.1 $E \square$ ect of treatments on mice behavior

Mice that were given different treatments exhibit different behavior. Group 1 (normal diet) and Group 2 (normal diet b JTK) showed normally active physical activity and normal appetite. In comparison, Group 3 (normal diet b DEN) showed less physical activity and became lethargic. The group also showed a decrease in appetite. DEN is an acute liver toxin. DEN injured the liver and decreased appetite and food consumption (You et al., 2021). The DEN-induced group that was also given JTK, black tea kombucha and JTB was more active compared to Group 3. This group also showed a higher level of appetite compared to Group 3. These might be caused by Javanese turmeric and black tea kombucha chemical components that could increase appetite and repair metabolism (Rahmat et al., 2021; Jayabalan et al., 2014). JTK, JTB and black tea kombucha restored the appetite of mice induced by DEN.

3.2 Serum glutamate pyruvate transaminase and serum glutamate oxalate transaminase enzyme activity

SGPT is an enzyme only present in the liver parenchyma cell cytosol thus its activity specifically indicates liver failure (Desiandura et al., 2022). SGOT is an enzyme found in liver cells cytosol and mitochondria. It also exists in cardiac muscle, skeletal muscle, pancreas and kidney. When the liver was exposed to DEN, the cells lysed and released SGPT and SGOT into the blood circulation (Castro et al., 2015). The evaluation of SGPT and SGOT revealed the hepatoprotective activity of kombucha. Low levels of SGPT and SGOT showed better hepatoprotective activity. The results of SGPT and SGOT activity measurement are shown in Table 1.

The lowest SGPT activity was found in the normal diet b JTK group. The normal SGPT activity value of healthy mice is 5-35 IU/L (Hutagalung et al., 2023). The SGPT activity of the normal diet group (71.8%) was still within range and lower than the normal diet group, which meant JTK enhanced healthy liver condition. The lowest SGOT activity was found in the normal diet group. The SGOT value of the normal diet group was within the normal range of 10-40 IU/L (Hutagalung et al., 2023). The normal diet b JTK group had significantly higher SGOT activity compared to the normal diet group, but still within the normal range. However, the high SGOT activity was not able to specifically indicate liver

Treatments	SGPT (U/L)	%decrease	SGOT (U/L)	%decrease
Normal diet	25.8 ^{ab} 6 9.4	0.0	14.6° 6 11.0	0.0
Normal diet b Javanese turmeric kombucha	7.3° 6 1.2	71.8	23.2 ^{bc} 6 2.9	-59.1
Normal diet b DEN	36.4ª 6 13.6	0.0 ^d	37.1ª 6 3.7	0.0 ^d
DEN b Nonfermented Javanese turmeric beverage	15.2 ^{bc} 6 8.7	58.2 ^b	25.8 ^{abc} 6 4.7	30.4 ^b
DEN p Javanese turmeric kombucha	8.6° 6 2.2	76.4ª	21.2 ^{bc} 6 8.6	42.9ª
DEN b black tea kombucha	18.5 ^{bc} 6 13.0	49.1°	31.8 ^{ab} 6 5.0	14.3°

Table 1. Activities of SGPT and SGOT in blood

Notes: SGPT ¼ serum glutamate pyruvate transaminase; SGOT ¼ serum glutamate oxaloacetate transaminase; DEN ¼ diethylnitrosamine; SGPT and SGOT decrease percentage of normal diet p Javanese turmeric kombucha were calculated based on the SGPT and SGOT value of normal diet. Decrease percentage of DEN b nonfermented Javanese turmeric, DEN b Javanese turmeric kombucha, and DEN b black tea kombucha was calculated based on the SGPT and SGOT value of normal diet p DEN; Data is presented in means 6 SD; data followed by different letters shows a significant difference (p < 0.05) serum of mice (n 1/4 7) Source: Created by authors

damage as SGOT activity is also present in different tissues such as cardiac muscle, skeletal muscle, pancreas and kidney (Desiandura *et al.*, 2022).

The normal diet **þ** DEN group showed the highest SGPT and SGOT activity (Table 1). The SGPT activity of this group was higher than the normal range of healthy mice, whereas the SGOT activity approached the upper limits. This indicated that DEN was able to induce liver damage and cause cell lysis. DEN was able to accumulate free radicals in liver cells and impair enzymes involved in liver DNA reparation. The free radicals harmed DNA and other nucleophilic compounds and resulted in the development of cancer cells (Arboatti *et al.*, 2018). When the liver cells were damaged, SGPT and SGOT were released from the cells and circulated in the bloodstream, therefore the DEN **þ** normal diet group showed high SGPT and SGOT activities (Castro *et al.*, 2015).

DEN **þ** JTB group had 58.2% lower SGPT and 30.4% lower SGOT activity compared to the normal diet **þ** DEN group. Javanese turmeric contains various antioxidant and antiinflammatory compounds such as xanthorrhizol, curcumin, calebin-A, phenolic compounds and triterpenoids. Antioxidants lower the number of free radicals and inhibit liver cytochrome P450 2E1 (CYP2E1), a gene responsible for the induction of microsomal oxidation that increases free radical production. This activity made liver cells produce the right amount of adenosine triphosphate (ATP) and ions, therefore creating a homeostatic condition. This condition decreased osmotic pressure and prevented cell membrane leakage. This activity prevented the release of SGPT and SGOT into the bloodstream (Puteri *et al.*, 2020).

Xanthorrhizol suppressed the phosphorylation of c-Jun N-terminal kinase in the mitogenactivated protein kinases signaling pathway (Oon *et al.*, 2015). The suppression prevented the transcription of cyclooxygenase (COX), inducible nitric oxide synthase (iNOS), c-Fos, and p50, therefore inhibiting the binding of nuclear factor-*k*B (NF-*k*B) and activator protein-1 (AP-1) with DNA. These transcription factors contribute to cell inflammation and apoptosis. Curcumin prevented lipid peroxidation and oxidative stress. It also restored the balance of the Bcl-2/Bax ratio and inhibited transforming growth factor beta (TGF-*b*), which suppressed liver cell apoptosis (Pramono *et al.*, 2018). Calebin-A lowered the regulation of tumor necrosis factoralpha (TNF-*a*), which consequently prevented the NF-*k*B activation (Tyagi *et al.*, 2017). Calebin-A inhibited COX, an enzyme that contributed to inflammation and cytochrome P450 1A2 (CYP1A2) and CYP2D6, which were responsible for transforming xenobiotics into procarcinogenic compounds that may damage liver cells (Oliveira *et al.*, 2015).

DEN þ JTK had 76.4% lower SGPT and 42.9% lower SGOT activity compared to the normal diet þ DEN group. This emphasized that JTK protected the liver from DEN induction. The hepatoprotective activity of JTK was contributed by xanthorrhizol, curcumin, calebin-A, phenolic compounds and terpenoid compounds (Zubaidah *et al.*, 2023). However, the SGPT and SGOT activity of the DEN þ JTK group was also lower than the DEN þ JTB group. This proved that kombucha fermentation enhanced the hepatoprotective activity of the raw material.

During the fermentation of black tea kombucha, yeasts and bacteria formed glucuronic acid, DSL and released complex polyphenol and flavonoid compounds (Wang *et al.*, 2013; Chakravorty *et al.*, 2016). These compounds were also found in JTK. Glucuronic acid increased the polarity of xenobiotics, decreased their toxicity, facilitated liver absorption, metabolism and excretion of xenobiotics through bile and urine. DSL produced by lactic acid bacteria (LAB) and *Gluconacetobacter sp.* inhibited *b*-glucuronidase, an enzyme responsible for the cleavage of glucuronic acid and toxicant. By this inhibition, the toxicant would bind to glucuronic acid and be easier to be excreted (Wang *et al.*, 2013). Microbes also released phenols that were bound to the substrate's matrices. Kombucha yeasts are dominated by *Candida*, a genus that possesses the ability to degrade complex polyphenol and flavonoid

compounds into simpler compounds (Chakravorty et al., 2016), which increases the number and antioxidant activity of the compounds.

DEN b black tea kombucha had 49.1% lower SGPT and 14.3% lower SGOT activity compared to normal diet b DEN. Black tea exhibited good hepatoprotective activity due to its anti-inflammatory and antioxidative activity. Black tea kombucha inhibited liver cell inflammation and apoptosis by maintaining the mitochondrial membrane potential so that the cells did not activate caspase-3, an enzyme that induces cell death (Lee et al., 2019). Black tea kombucha also possessed glucuronic acid and DSL. On the other hand, black tea itself contained various polyphenolic compounds, predominantly catechin. During fermentation, microbes hydrolyzed catechin and converted other polyphenolic compounds into catechin, which increased the antioxidant activity (Jakubczyk et al., 2020).

The SGPT and SGOT activity of DEN b JTK were lower than the DEN b black tea kombucha group. Javanese turmeric contains xanthorrhizol, curcumin, phenolic compounds and triterpenoids which do not exist in black tea (Puteri et al., 2020). During kombucha fermentation, microbes released these hepatoprotective compounds and produced organic acids such as glucuronic acids and DSL by using sugars added to the substrate (Martínez-Leal et al., 2018). It was assumed that there was a synergy between hepatoprotective compounds of Javanese turmeric and microbial activity during fermentation.

3.3 Lipid peroxidation level

MDA was a product of a secondary oxidation process that could be found in the cell membranes. Higher MDA levels indicate higher oxidation levels in the body (Castro et al., 2015). Table 2 shows the result of MDA measurements. The normal diet group resulted in a significantly higher MDA level compared to the normal diet **b** JTK group. This proved that JTK was able to decrease oxidative stress levels in healthy mice. The normal diet b DEN group had the highest MDA level. DEN was metabolized in the liver by P450 monooxygenase, resulting in active ethyl radicals (Mansour et al., 2019). These free radicals increased lipid peroxidation, suppressed antioxidant enzyme activity and increased the number of toxic metabolites such as 4-hydroxynonenal and MDA (Zhang et al., 2021).

DEN b JTB and DEN b JTK performed a lower MDA level compared to the normal diet b DEN group. Javanese turmeric protects the liver from DEN-caused oxidative stress by increasing glutathione reductase, superoxide dismutase, glutathione peroxidase and glutathione S-transferase activity (Lukitaningsih et al., 2020). Xanthorrhizol and curcumin

Treatments	MDA (nanomol/mL)	%decrease
Normal diet	4.7 ^{ab} 6 0.25	0.0
Normal diet b Javanese turmeric kombucha	3.4° 6 0.09	27.1
Normal diet b DEN	5.1° 6 1.00	0.0°
DEN b Nonfermented Javanese turmeric beverage	3.8 ^{bc} 6 0.23	25.6 ^b
DEN b Javanese turmeric kombucha	2.9° 6 0.08	42.9ª
DEN b black tea kombucha	2.9° 6 0.06	42.4ª

Table 2. MDA levels in blood

Notes: MDA ¼ malonaldehyde; DEN ¼ diethylnitrosamine; MDA decrease percentage of normal diet p Javanese turmeric kombucha was calculated based on the MDA value of normal diet MDA decrease percentage of DEN b nonfermented Javanese turmeric, DEN b Javanese turmeric kombucha, and DEN b black tea kombucha was calculated was calculated based on the MDA value of normal diet b DEN; Data is presented in means 6 SD; data followed by different letters shows a significant difference (p < 0.05) serum of mice (n 1/4 7) Source: Created by authors

donated hydrogen to free radicals and chelated oxygen to prevent oxidation (Noreen *et al.*, 2017). Xanthorrhizol can prevent LDL peroxidation due to its phenolic hydroxyl group and bisabolene backbone (Oon *et al.*, 2015). Xanthorrhizol chelated Cu^{2p} , decreased the catalyst of LDL peroxidation and improved the activity of superoxide dismutase (SOD) and catalase that resulted in less MDA formation.

The MDA level of DEN b JTK was less than DEN b JTB. During kombucha fermentation, microbes released enzymes that freed bound antioxidants. Acetic acid bacteria oxidized sugars into aldehydes and organic acids, which donated H^b to free radicals (Hardiwati *et al.*, 2019). LAB produced curcumin/dehydrocurcumin reductase that reduced curcumin into its analog (THC, hexahydrocurcumin and octahydrocurcumin) in the presence of nicotinamide adenine dinucleotide phosphate (Huang *et al.*, 2018). THC stabilized hydroxyl radicals, reduced MDA and increased SOD and catalase better than curcumin (Morales *et al.*, 2015; Trivedi *et al.*, 2020).

DEN **þ** black tea kombucha displayed the lowest MDA level. Nonfermented black tea is known to have good antioxidant activities due to its components (catechin, epicatechin, epigallocatechin, epicatechin gallate, epigallocatechin gallate, theaflavin and thearubigin). Black tea kombucha possessed higher levels of phenolic compounds compared to nonfermented black tea (Jakubczyk *et al.*, 2020). Kombucha also had gluconic acids and thearubigin derivatives that were able to stabilize free radicals. The presence of DSL and ascorbic acids also helped eliminate toxicants which could be metabolized into free radicals inside the body (Morales, 2020). The MDA level of DEN **þ** black tea kombucha showed no significant difference with DEN **þ** JTK and normal diet **þ** JTK.

3.4 Liver histology

In Figure 1, normal diet mice liver and normal diet þ JTK mice liver showed purple cytoplasm, dark nucleus, clear cell lining and distinct central vein. This histology was in concordance with the description of normal mice liver histology (Devaraj *et al.*, 2014). DEN þ normal diet showed ambiguous cell linings, bright pink cytoplasm and loss of nuclei. A group of cells were degraded and replaced by inflammatory cells, which was caused by continuous lobule damage and indicated liver hepatitis (Krishna, 2017). The reddish cells with no nuclei were apoptotic bodies, indicating liver cell death by DEN (Boyd *et al.*, 2020).

The DEN **þ** JTB group showed lesser bile inflammation and apoptotic bodies than the DEN **þ** normal diet group. The DEN **þ** black tea kombucha group underwent bile inflammation and had some apoptotic bodies, although the degraded cell area was not as large as the normal diet **þ** DEN group. DEN **þ** JTK experienced central vein inflammation, but not as serious as other DEN-induced groups. No apoptotic bodies were seen in the DEN **þ** JTK group, indicating that the cells were not seriously injured by DEN and had not yet experienced apoptosis.

Figure 2 shows that the DEN p normal diet group had the highest damaged cell number. DEN could induce DNA methylation which triggers apoptosis (Tolba *et al.*, 2015). DEN induction activated macrophage and neutrophils, which consequently produced proinflammatory cytokine and chemokine. DEN also stimulated the infiltration of inflammatory cells such as lymphocytes, eosinophils and Kupffer cells (Mansour *et al.*, 2019). Liver inflammation can increase vein pressure and decrease intrahepatic nitrite oxide. The suppression of nitrite oxide reduced the ability of vasodilatation and triggered cirrhosis (Vizzutti *et al.*, 2013).

DEN þ JTK and normal diet þ JTK groups had the lowest number of damaged cells, followed by DEN þ JTB and DEN þ black tea kombucha groups. This proved that JTK had better hepatoprotective activity compared to JTB and black tea kombucha. The damaged



(a)





Notes: (a) Normal diet; (b) normal diet + JTK; (c) normal diet + DEN; (d) DEN + nonfermented JTB; (e) DEN + black tea kombucha; (f) DEN + JTK; 400× magnification I = inflammation; A = apoptotic bodies; F = fibrosis**Source:** Created by authors

cell number for DEN þ JTK was nearly the same as the normal diet þ JTK group. This indicated that JTK was able to alleviate liver damage and restore the liver to its normal condition.

4. Conclusions

Fermentation enhanced the hepatoprotective activity of Javanese turmeric due to the release of bioactive compounds and new bioactive compound formation. As evidenced by the least damaged liver-cell numbers, JTK demonstrated better hepatoprotective activity compared to JTB.

Figure 1. Mice liver histology



Figure 2.

Note: DEN = diethylnitrosamine **Source:** Created by authors Damaged cell number on the field of view of

100x magnification

References

- Arboatti, A.S., Lambertucci, F., Sedlmeier, M.G., Pisani, G., Monti, J., Álvarez, M.L., Franc, es, D.E.A., Ronco, M.T. and Carnovale, C.E. (2018), "Diethylnitrosamine increases proliferation in early stages of hepatic carcinogenesis in insulin-treated type 1 diabetic mice", *BioMed Research International*, Vol. 2018.
- Boyd, A., Cain, O., Chauhan, A. and Webb, G.J. (2020), "Medical liver biopsy: background, indications, procedure and histopathology", *Frontline Gastroenterology*, Vol. 11 No. 1, pp. 40-47.
- Castro, C.A., Dias, M.M.S., Silva, K.A., Reis, S.A., Conceição, L.L., Marcon, L.D.N., Moraes, L.F.S. and Peluzio, M.C.G. (2015), "Liver biomarkers and their applications to nutritional interventions in animal studies", in Preedy, V. (Ed.), *Biomarkers in Liver Disease*, Springer, New York, NY, pp. 1-24.
- Chakravorty, S., Bhattacharya, S., Chatzinotas, A., Chakraborty, W., Bhattacharya, D. and Gachhui, R. (2016), "Kombucha tea fermentation: microbial and biochemical dynamics", *International Journal of Food Microbiology*, Vol. 220, pp. 63-72.
- Desiandura, K., Rahayu, A. and Wibisomo, FJ. (2022), "Cat's liver disease detection with SGOT and SGPT evaluation as a gold standard diagnosis", *Indonesian Journal of Tropical and Infectious Disease*, Vol. 10 No. 1, pp. 48-54.
- Devaraj, S., Ismail, S., Ramanathan, S. and Yam, M.F. (2014), "Investigation of antioxidant and hepatoprotective activity of standardized *Curcuma xanthorrhiza* rhizome in carbon tetrachloride-induced hepatic damaged rats", *The Scientific World Journal*, Vol. 2014.
- Elsharkawy, E.E., El-Nisr, N.A., Wahba, N.M. and Elsherif, W.M. (2022), "Hepatotoxicity of methoxychlor and camel milk restoration", *Nutrition and Food Science*, Vol. 52 No. 3, pp. 483-496.
- Fortes, R.C. (2017), "Nutritional implications of chronic liver disease", Journal of Liver Research, Disorders, and Therapy, Vol.3 No.5, pp. 131-133.
- Hardiwati, K.T., Seninha, M., Lay, B.W. and Yanti, L. (2019), "Curcuminoid cider fermented from *Curcuma xanthorrhiza* curcuminoids attenuates gene expression related to obesity-induced inflammation in hypercholesterolaemic rats", *International Food Research Journal*, Vol. 26 No. 3, pp. 859-867.
- Huang, Y., Cao, S., Zhang, Q., Zhang, H., Fan, Y., Qiu, F. and Kang, N. (2018), "Biological and pharmacological effects of hexahydrocurcumin, a metabolite of curcumin", *Archives of Biochemistry and Biophysics*, Vol. 646, pp. 31-37.
- Hutagalung, C.F., Girsang, E., Mutia, M.S., Suhartina, K. and Natali, O. (2023), "Liver protection effect of red rose petal methanol extract in copper-induced liver injury in male Wistar rats", *Biology*, *Medicine*, and Natural Product Chemistry, Vol. 12 No. 2, pp. 507-513.

- Jakubczyk, K., Kałdun, ska, J., Kochman, J. and Janda, K. (2020), "Chemical profile and antioxidant activity of the Kombucha beverage derived from white, green, black and red tea", *Antioxidants*, Vol. 9 No. 5, p. 447.
- Jayabalan, R., Malbaša, R.V., Lon~car, E.S., Vitas, J.S. and Sathishkumar, M. (2014), "A review on kombucha tea—microbiology, composition, fermentation, beneficial effects, toxicity, and tea fungus", Comprehensive Reviews in Food Science and Food Safety, Vol. 13 No. 4, pp. 538-550.
- Krishna, M. (2017), "Patterns of necrosis in liver disease", Clinical Liver Disease, Vol. 10 No. 2, pp. 53-56.
- Lee, C., Kim, J., Wang, S., Sung, S., Kim, N., Lee, H.H., Seo, Y.S. and Jung, Y. (2019), "Hepatoprotective effect of kombucha tea in rodent model of nonalcoholic fatty liver disease/nonalcoholic steatohepatitis", *International Journal of Molecular Sciences*, Vol. 20 No. 9, p. 2369.
- Lukitaningsih, E., Rohman, A., Rafi, M., Nurrulhidayah, A.F. and Windarsih, A. (2020), "In vivo antioxidant activities of *Curcuma longa* and *Curcuma xanthorrhiza*: a review", *Food Research*, Vol. 4 No. 1, pp. 13-19.
- Mansour, D.F., Abdallah, H.M.I., Ibrahim, B.M.M., Hegazy, R.R., Esmail, R.S.E. and Abdel-Salam, L.O. (2019), "The carcinogenic agent diethylnitrosamine induces early oxidative stress, inflammation and proliferation in rat liver, stomach and colon: protective effect of ginger extract", Asian Pacific Journal of Cancer Prevention, Vol. 20 No. 8, pp. 2551-2561.
- Martínez-Leal, J., Su, arez, L.V., Jayabalan, R., Oros, J.H. and Escalante-Aburto, A. (2018), "A review on health benefits of kombucha nutritional compounds and metabolites", *CyTA - Journal of Food*, Vol. 16 No. 1, pp. 390-399.
- Miranda, J.F., Ruiz, L.F., Silva, C.B., Uekane, T.M., Silva, K.A., Gonzalez, A.G.M., Fernandes, F.F. and Lima, A.R. (2022), "Kombucha: a review of substrates, regulations, composition, and biological properties", *Journal of Food Science*, Vol. 87 No. 2, pp. 503-527.
- Morales, D. (2020), "Biological activities of kombucha beverages: the need of clinical evidence", Trends in Food Science and Technology, Vol. 105, pp. 323-333.
- Morales, N.P., Sirijaroonwong, S., Yamanont, P. and Phisalaphong, C. (2015), "Electron paramagnetic resonance study of the free radical scavenging capacity of curcumin and its demethoxy and hydrogenated derivatives", *Biological and Pharmaceutical Bulletin*, Vol. 38 No. 10, pp. 1478-1483.
- Noreen, H., Semmar, N., Farman, M. and McCullagh, J.S.O. (2017), "Measurement of total phenolic content and antioxidant activity of aerial parts of medicinal plant Coronopus didymus", Asian Pacific Journal of Tropical Medicine, Vol. 10 No. 8, pp. 792-801.
- Oliveira, A.L.P., Martinez, S.E., Nagabushnam, K., Majeed, M., Alrushaid, S., Sayre, C.L. and Davies, N. M. (2015), "Calebin A: analytical development for pharmacokinetics study, elucidation of pharmacological activities and content analysis of natural health products", *Journal of Pharmacy* and Pharmaceutical Sciences, Vol. 18 No. 4, pp. 494-514.
- Oon, S.F., Nallappan, M., Tee, T.T., Shohaimi, S., Kassim, N.K., Sa'ariwijaya, M.S.F. and Cheah, Y.H. (2015), "Xanthorrhizol: a review of its pharmacological activities and anticancer properties", *Cancer Cell International*, Vol. 15 No. 1.
- Pramono, S., Arifah, F.H., Pribadi, F.H. and Nugroho, A.E. (2018), "Hepatoprotective activity of *Curcuma xanthorrhiza* Roxb. on paracetamol-induced liver damage in rats and correlation with their chemical compounds", *Thai Journal of Pharmaceutical Sciences*, Vol. 42 No.4, pp. 188-195.
- Puteri, A.I.S., Sandhika, W. and Hasanatuludhhiyah, N. (2020), "Effect of Javanese turmeric (*Curcuma xanthorrhiza*) extract on hepatitis model of alcohol-induced mice", *Jurnal Kedokteran Brawijaya*, Vol. 31 No. 1, pp. 39-42.
- Rahmat, E., Lee, J. and Kang, Y. (2021), "Javanese turmeric (*Curcuma xanthorrhiza* Roxb.): phytochemistry, biotechnology, and pharmacological activities", *Evidence-Based Complementary* and Alternative Medicine, Vol. 2021.

Salafzoon, S., Mahmoodzadeh Hosseini, H. and Halabian, R. (2018), "Evaluation of the antioxidant impact of ginger-based kombucha on the murine breast cancer model", *Journal of Complementary and Integrative Medicine*, Vol. 15 No. 1, pp. 1-8.

- Tolba, R., Kraus, T., Liedtke, C., Schwarz, M. and Weiskirchen, R. (2015), "Diethylnitrosamine (DEN)induced carcinogenic liver injury in mice", *Laboratory Animals*, Vol. 49 No. 1_suppl, pp. 59-69.
- Trivedi, M.K., Panda, P., Sethi, K.K., Gangwar, M., Mondal, S.C. and Jana, S. (2020), "Solid and liquid state characterization of tetrahydrocurcumin using XRPD, FT-IR, DSC, TGA, LC-MS, GC-MS, and NMR and its biological activities", *Journal of Pharmaceutical Analysis*, Vol. 10 No. 4, pp. 334-345.
- Tyagi, A., Prasad, S., Majeed, M. and Aggarwal, B.B. (2017), "Calebin A, a novel component of turmeric, suppresses NF-kB regulated cell survival and inflammatory gene products leading to inhibition of cell growth and chemosensitization", *Phytomedicine*, Vol. 34, pp. 171-181.
- Vizzutti, F., Arena, U., Laffi, G. and Marra, F. (2013), "Acute on chronic liver failure: from pathophysiology to clinical management", *Trends in Anaesthesia and Critical Care*, Vol. 3 No. 3, pp. 122-129.
- Wang, Y., Ji, B., Wu, W., Wang, R., Yang, Z., Zhang, D. and Tian, W. (2013), "Hepatoprotective effects of kombucha tea: identification of functional strains and quantification of functional components", *Journal of the Science of Food and Agriculture*, Vol. 94 No. 2, pp. 265-272.
- You, Y., Zhu, F., Li, Z., Zhang, L., Xie, Y., Chinnathambi, A., Alahmadi, T.A. and Lu, B. (2021), "Phyllanthin prevents diethylnitrosamine (DEN) induced liver carcinogenesis in rats and induces apoptotic cell death in HepG2 cells", *Biomedicine and Pharmacotherapy*, Vol. 137, p.111335.
- Zhang, Y., Li, X. and Li, X. (2021), "Curcumae ameliorates diethylnitrosamine-induced hepatocellular carcinoma via alteration of oxidative stress, inflammation and gut microbiota", *Journal of Inflammation Research*, Vol. 2021 No. 14, pp. 5551-5566.
- Zubaidah, E., Dea, E.C. and Sujuti, H. (2022), "Physicochemical and microbiological characteristics of kombucha based on various concentration of Javanese turmeric (*Curcuma xanthorrhiza*)", *Biocatalysis and Agricultural Biotechnology*, Vol. 44, doi: 10.1016/j.bcab.2022.102467.
- Zubaidah, E., Yurista, S. and Rahmadani, N.R. (2018), "Characteristics of physical, chemical, and microbiological kombucha from various varieties of apples", *IOP Conference Series: Earth and Environmental Science*, Vol. 131 No. 1.
- Zubaidah, E., Susanti, I., Sujuti, H., Martanti, E., Rahayu, A.P., Srianta, I. and Tewfik, I. (2023), "The distinctive hepatoprotective activity of turmeric kombucha (curcuma longa) induced by diethylnitrosamine in Balb/C mice", *Food Bioscience*, Vol. 55, p. 103043.

Further reading

- Aggarwal, B.B., Deb, L. and Prasad, S. (2015), "Curcumin differs from tetrahydrocurcumin for molecular targets, signaling pathways and cellular responses", *Molecules*, Vol. 20 No. 1, pp. 185-205.
- Gurung, R.B., Gong, S.Y., Dhakal, D., Le, T.T., Jung, N.R., Jung, H.J., Oh, T.J. and Sohng, J.K. (2017), "Synthesis of curcumin glycosides with enhanced anticancer properties using one-pot multienzyme glycosylation technique", *Journal of Microbiology and Biotechnology*, Vol. 27 No. 9, pp. 1639-1648.
- Hashish, E.A. and Elgaml, S.A. (2016), "Hepatoprotective and nephroprotective effect of curcumin against copper toxicity in rats", *Indian Journal of Clinical Biochemistry*, Vol. 31 No. 3, pp. 270-277.
- Huh, E., Lim, S., Kim, H.G., Ha, S.K., Park, H.Y., Huh, Y. and Oh, M.S. (2018), "Ginger fermented with: schizosaccharomyces pombe alleviates memory impairment via protecting hippocampal neuronal cells in amyloid beta1-42 plaque injected mice", *Food and Function*, Vol. 9 No. 1, pp. 171-178.

- Imran, M., Nadeem, M., Khan, M.A., Ahmed, S., Imran, A., Amir, R.M., Arshad, M.U., Gilani, S.A., Saeed, F., Rauf, A., Mehmood, Z., Khan, S. and Suleria, H.A.R. (2017), "Curcumin and its allied analogues: epigenetic and health perspectives - a review", *Czech Journal of Food Sciences*, Vol. 35 No. 4, pp. 285-310.
- Jantan, I., Saputri, F.C., Qaisar, M.N. and Buang, F. (2012), "Correlation between chemical composition of curcuma domestica and *Curcuma xanthorrhiza* and their antioxidant effect on human lowdensity lipoprotein oxidation", *Evidence-Based Complementary and Alternative Medicine*, Vol. 2012.
- Kim, M.B., Kim, C., Song, Y. and Hwang, J.K. (2014), "Antihyperglycemic and anti-inflammatory effects of standardized *Curcuma xanthorrhiza* Roxb. Extract and its active compound xanthorrhizol in high-fat diet-induced obese mice", *Evidence-Based Complementary and Alternative Medicine*, Vol. 2014.
- Vitas, J., Vukmanovi, C.S., C~akarevi, C.J., Popovi, C.L. and Malbaša, R. (2020), "Kombucha fermentation of six medicinal herbs: chemical profile and biological activity", *Chemical Industry and Chemical Engineering Quarterly*, Vol. 26 No. 2, pp. 157-170.
- Zhao, F., Gong, Y., Hu, Y., Lu, M., Wang, J., Dong, J., Chen, D., Chen, L., Fu, F. and Qiu, F. (2015), "Curcumin and its major metabolites inhibit the inflammatory response induced by lipopolysaccharide: translocation of nuclear factor-*k*B as potential target", *Molecular Medicine Reports*, Vol. 11 No. 4, pp. 3087-3093.

Corresponding author Elok Zubaidah can be contacted at: elok@ub.ac.id Redefining the hepatoprotective potential of Javanese turmeric (Curcuma xanthorrhiza) Kombucha towards the diethylnitrosamine-induced hepatotoxicity of mice

ORIGINALITY REPORT

% SIMILARITY INDEX **INTERNET SOURCES** PUBLICATIONS STUDENT PAPERS **PRIMARY SOURCES** docslib.org % Internet Source repository.wima.ac.id 1% 2 **Internet Source** Patricia Viera de Oliveira, Afonso Henrique 3 1% da Silva Júnior, Carlos Rafael Silva de Oliveira, Carolina Fagundes Assumpção et al. "Kombucha benefits, risks and regulatory frameworks: A review", Food Chemistry Advances, 2023 Publication

Exclude quotes	On	Exclude matches	< 1%
Exclude bibliography	On		

Redefining the hepatoprotective potential of Javanese turmeric (Curcuma xanthorrhiza) Kombucha towards the diethylnitrosamine-induced hepatotoxicity of mice

GRADEMARK REPORT

FINAL GRADE	GENERAL COMMENTS
/100	
PAGE 1	
PAGE 2	
PAGE 3	
PAGE 4	
PAGE 5	
PAGE 6	
PAGE 7	
PAGE 8	
PAGE 9	
PAGE 10	
PAGE 11	
PAGE 12	
PAGE 13	