

# Induction of High Neutralizing Activity Against Both Omicron BA.2 and Omicron BA.1 by Coronavirus Disease 2019 Messenger RNA Booster Vaccination

*by* Silvia Sutandhio

---

**Submission date:** 02-Aug-2025 01:54PM (UTC+0700)

**Submission ID:** 2724050493

**File name:** jjac159.pdf (136K)

**Word count:** 1990

**Character count:** 10589

Participants were 20 healthy Thai adults aged 18 years and older (mean age 50.66 years, SD 12.95 years) who received 2 doses of AZD1222 vaccination and had no previous or current COVID-19 infection. The participants received mRNA-1273 as a booster (third dose) vaccine at 5–7 months after the second dose of AZD1222. Blood samples were collected at 28 days after the booster. In accordance with the results previously reported by Tjan et al and Yu et al [1, 4] Undetectable (titer < 20) nAbs against both BA.1 and BA.2 were observed among most participants before the booster. At 28 days after the booster the geometric mean titer nAbs significantly increased from 16.6 (95% confidence interval CI], 13.2–21.0) and 11.0 (95% CI, 9.6–12.6) to 548 (95% CI, 415–723) and 324 (95% CI, 214–492) against BA.1 and BA.2, respectively. The results demonstrated that after the mRNA booster, the nAbs titers against BA.2 were slightly lower than those against BA.1 (Figure 1).

These preliminary results indicate that the heterologous booster with mRNA vaccine in AZD1222-primed individuals could induce a robust antibody response that can cross-neutralize both BA.1 and BA.2 omicron variants.

## Notes

**Acknowledgments.** We thank the staff of the Center of Excellence in Clinical Virology and all the participants for helping and supporting this project; and the Ministry of Public Health, Chulabhorn Royal Academy, and Zullig Pharma for providing the vaccines for this study.

**Financial support.** This work was supported by the Health Systems Research Institute, National Research Council of Thailand; Chulalongkorn University (Center of Excellence in Clinical Virology and Second Century Fund); King Chulalongkorn Memorial Hospital; and the National Center for

Genetic Engineering and Biotechnology (grant number P2051613 to T. D.).

**Potential conflicts of interest.** The authors: No reported conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

Suvichada Assawakosri,<sup>1,2</sup> Sittichai Kanokudom,<sup>1,2</sup> Nungpruthai Suntronwong,<sup>1</sup> Jiratchaya Puempha,<sup>1</sup> Thananya Duangchinda,<sup>3</sup> Warangkana Chantima,<sup>4,5</sup> Pattarakul Pakchotanon,<sup>7</sup> Juthathip Mongkolkeha,<sup>4,7</sup> Nissamon Wanlapakorn,<sup>8</sup> Sittisak Honsawek,<sup>2</sup> and Yong Poovorawan<sup>1,8</sup>

<sup>1</sup>Center of Excellence in Clinical Virology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand; <sup>2</sup>Osteoarthritis and Musculoskeletal Research Unit, Faculty of Medicine, Chulalongkorn University, King Chulalongkorn Memorial Hospital, Thai Red Cross Society, Bangkok, Thailand; <sup>3</sup>Molecular Biology of Dengue and Flaviviruses Research Team, National Center for Genetic Engineering and Biotechnology, National Science and Development Agency, Pathum Thani, Thailand; <sup>4</sup>Division of Dengue Hemorrhagic Fever Research, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand; <sup>5</sup>Siriraj Center of Research Excellence in Dengue and Emerging Pathogens, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand; <sup>6</sup>Wellcome Centre for Human Genetics, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom; <sup>7</sup>Chinese Academy of Medical Science, Oxford Institute, University of Oxford, Oxford, United Kingdom; and <sup>8</sup>Fellow of Royal Society of Thailand FRSTI, the Royal Society of Thailand, Bangkok, Thailand

## References

1. Tjan LH, Furukawa K, Kurahashi Y, et al. As well as omicron BA.1, high neutralizing activity against omicron BA.2 can be induced by COVID-19 mRNA booster vaccination. *J Infect Dis* 2022; 226:1482.
2. Assawakosri S, Kanokudom S, Suntronwong N, et al. Neutralizing activities against the omicron variant after a heterologous booster in healthy adults receiving two doses of CoronaVac vaccination. *J Infect Dis* 2022; 226:1372–81.
3. World Health Organization. COVID-19 weekly epidemiological update on COVID-19 - 5 April 2022, edn 86. <https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19-5-april-2022>. Accessed 8 April 2022.

4. Yu J, Collier A-RY, Rowe M, et al. Neutralization of the SARS-CoV-2 omicron BA.1 and BA.2 variants. *N Engl J Med* 2022; 386:1579–80.

Received 21 April 2022; editorial decision 21 April 2022; accepted 22 April 2022; published online 27 April 2022

Correspondence: Y. Poovorawan, MD, Center of Excellence in Clinical Virology, Department of Pediatrics, Faculty of Medicine, Chulalongkorn University, Bangkok, 10330 Thailand (yong.p@chula.ac.th).

The Journal of Infectious Diseases® 2022;226:1480–1482. The Author(s) 2022. Published by Oxford University Press on behalf of Infectious Diseases Society of America. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com  
<https://doi.org/10.1093/infdis/jiac158>

## Induction of High Neutralizing Activity Against Both Omicron BA.2 and Omicron BA.1 by Coronavirus Disease 2019 Messenger RNA Booster Vaccination

TO THE EDITOR—Assawakosri et al [1] reported that heterologous booster vaccines significantly increased binding and neutralizing antibodies (nAbs) against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) variants of concern (VOCs), including the Omicron variant (BA.1) in individuals immunized with 2 doses of CoronaVac, indicating possible vaccine strategies to thwart VOCs. The most recently designated VOC of SARS-CoV-2, Omicron, is associated with an increased risk of reinfection [2]. Among sublineages of Omicron (BA.1, BA.1.1, BA.2, and BA.3), BA.1 has spread to >151 countries and is responsible for the greatly increased number of coronavirus disease 2019 (COVID-19) cases worldwide. However, Omicron BA.2 has also been detected in ≥85 countries and became the dominant lineage in 18 countries by mid-February 2022 [3]. BA.2 has become the next dominant variant worldwide.

A recent epidemiological study in South Africa suggested that the clinical profile of illness caused by BA.1 infection is similar to that caused by BA.2 [4]. Although BA.2 infection has generally been associated with mild illness, its high transmissibility may be

accompanied by high numbers of cases with considerable societal impacts (eg, greater work absences).

In response to the surge in Omicron BA.1 cases, booster vaccination was initiated in many countries. In addition to the study reported by Assawakosri et al [1], our study and others showed that BA.1 escapes 2 doses of BNT162b2 messenger RNA vaccine-induced neutralization and that a booster (third) vaccination is required to induce the nAb against BA.1 [5, 6].

There is, however, limited evidence regarding the effectiveness of a booster vaccination against BA.2 [7, 8]. Therefore, we collected blood samples from 84 physicians at Kobe University Hospital in Kobe, Japan, in January 2022 (median age, 44 years; interquartile range, 33–58 years) about 7 months after they had received 2 BNT162b2 vaccinations and about 2 weeks after their first booster vaccination. We

performed a serum neutralizing assay against BA.2 using authentic virus, as described elsewhere [5]. No participants had a history of SARS-CoV-2 infection. The study was approved by the ethical committee of the Kobe University Graduate School of Medicine (approval code B200200). All participants were recruited and provided written informed consent.

The results demonstrated that, similar to results with BA.1 [5], most participants had no or a very low nAb titer against BA.2 at 7 months after 2 BNT162b2 vaccinations (geometric mean titer, 1.18 [95% confidence interval 1.09–1.27]). However, the titer increased significantly 2 weeks after the booster vaccination (geometric mean titer, 36.44 [95% confidence interval, 30.53–43.50];  $P < .001$ ) (Figure 1).

These results indicate that a booster vaccination could induce neutralizing immunity against Omicron BA.2 (as it

has against BA.1) and that a booster dose of BNT162b2 messenger RNA vaccine induces a high cross-neutralizing response against SARS-CoV-2 variants [5]. This may indicate that booster vaccination is a meaningful approach for the suppression of BA.2 pandemic and can activate memory B cells that produce nAbs recognizing epitopes conserved among SARS-CoV-2 variants.

## Notes

**Acknowledgments.** We gratefully acknowledge Kazuro Sugimura MD, PhD (superintendent of the Hyogo Prefectural Hospital Agency and professor at Kobe University), for giving his full support to this study. We also thank the National Institute of Infectious Disease Japan for providing the severe acute respiratory syndrome coronavirus 2 Omicron BA.2 variant.

**Disclaimer.** The funders had no role in study design, data collection and analysis, the decision to publish, or the preparation of the manuscript.

**Financial support.** This work was supported by the Hyogo Prefectural Government (Y. M.).


**Potential conflicts of interest.** All authors: No reported conflicts. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

Lidya Handayani Tjan, Koichi Furukawa, Yukiya Kurahashi, Silvia Sutandhio, Mitsuhiro Nishimura, Jun Arai, and Yasuko Mori

Division of Clinical Virology, Center for Infectious Diseases, Kobe University Graduate School of Medicine, Kobe, Japan

## References

- Assawakosri S, Kanokudom S, Suntronwong N, et al. Neutralizing activities against the Omicron variant after a heterologous booster in healthy adults receiving two doses of CoronaVac vaccination. *J Infect Dis* 2022; 226:1372–81.



**Figure 1.** Neutralizing antibody (nAb) titers against Omicron BA.2 in BNT162b2 messenger RNA-vaccinated men (n=84) at 7 months after they had received 2 vaccine doses and at 2 weeks after a booster vaccination. Dotted horizontal line represents the limit of detection; solid horizontal lines, the geometric mean titers. Titers were compared by means of the 2-sample Wilcoxon rank sum (Mann-Whitney) test; 2-tailed  $P$  values were calculated.

2. Altarawneh HN, Chemaitelly H, Hasan MR, et al. Protection against the Omicron variant from previous SARS-CoV-2 infection. *N Engl J Med* **2022**; 386:1288–90.
3. World Health Organization (WHO). COVID-19 weekly epidemiological update edition 80. <https://apps.who.int/iris/bitstream/handle/10665/352199/CoV-weekly-sitrep22Feb22-eng.pdf>. Accessed 15 March 2022.
4. Wolter N, Jassat W, Walaza S, et al. Early assessment of the clinical severity of the SARS-CoV-2 omicron variant in South Africa: a data linkage study. *Lancet* **2022**; 399:437–46.
5. Furukawa K, Tjan LH, Kurahashi Y, et al. Acquired neutralizing breadth against SARS-CoV-2 variants including Omicron after three doses of mRNA COVID-19 vaccination and the vaccine efficacy. *JAMA Netw Open* **2022**; 5(5):e2210780.
6. Nemet I, Kliker L, Lustig Y, et al. Third BNT162b2 vaccination neutralization of SARS-CoV-2 Omicron infection. *N Engl J Med* **2022**; 386:492–4.
7. Chen LL, Chu AW, Zhang RR, et al. Serum neutralisation of the SARS-CoV-2 Omicron sublineage BA.2. *Lancet Microbe* **2022**. doi:10.1016/S2666-5247(22)00060-X.
8. Yu J, Collier AY, Rowe M, et al. Neutralization of the SARS-CoV-2 Omicron BA.1 and BA.2 variants. *N Engl J Med* **2022**; 386:1579–80.

Received 07 April 2022; editorial decision 21 April 2022; accepted 24 April 2022; published online 28 April 2022

Correspondence: Yasuko Mori, Division of Clinical Virology, Center for Infectious Diseases, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe, Hyogo 650-0017, Japan ([ymori@med.kobe-u.ac.jp](mailto:ymori@med.kobe-u.ac.jp))

**The Journal of Infectious Diseases**® 2022;226:1481–3  
© The Author(s) 2022. Published by Oxford University Press on behalf of Infectious Diseases Society of America. All rights reserved. For permissions, please e-mail: [journals.permissions@oup.com](mailto:journals.permissions@oup.com)  
<https://doi.org/10.1093/infdis/jiac159>

# Induction of High Neutralizing Activity Against Both Omicron BA.2 and Omicron BA.1 by Coronavirus Disease 2019 Messenger RNA Booster Vaccination

## ORIGINALITY REPORT

6%

SIMILARITY INDEX

3%

INTERNET SOURCES

6%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1

M. L. Houben, L. Bont, B. Wilbrink, M. E. Belderbos, J. L. L. Kimpen, G. H. A. Visser, M. M. Rovers. "Clinical Prediction Rule for RSV Bronchiolitis in Healthy Newborns: Prognostic Birth Cohort Study", PEDIATRICS, 2010

Publication

1%

2

Emanuele Andreano, Ida Paciello, Silvia Marchese, Lorena Donnici et al. "Anatomy of Omicron BA.1 and BA.2 neutralizing antibodies in COVID-19 mRNA vaccinees", Nature Communications, 2022

Publication

1%

3

Alona Kuzmina, Dina Korovin, Ido Cohen Iass, Nofar Atari et al. "Changes within the P681 residue of spike dictate cell fusion and syncytia formation of Delta and Omicron variants of SARS-CoV-2 with no effects on neutralization or infectivity", Heliyon, 2023

Publication

1%

4

Brittany A. Petros, Jacquelyn Turcinovic, Nicole L. Welch, Laura F. White et al. "Early introduction and rise of the Omicron SARS-CoV-2 variant in highly vaccinated university populations", Clinical Infectious Diseases, 2022

Publication

1%

5

[pmc.ncbi.nlm.nih.gov](https://pmc.ncbi.nlm.nih.gov)  
Internet Source

1%

6

S.O Minka, F.H Minka. "A tabulated summary of the evidence on humoral and cellular

1%

responses to the SARS-CoV-2 Omicron VOC,  
as well as vaccine efficacy against this  
variant.", Immunology Letters, 2022  
Publication

---

7

pt.scribd.com  
Internet Source

1 %

---

Exclude quotes    On  
Exclude bibliography    On

Exclude matches    < 1%

# Induction of High Neutralizing Activity Against Both Omicron BA.2 and Omicron BA.1 by Coronavirus Disease 2019 Messenger RNA Booster Vaccination

---

GRADEMARK REPORT

---

FINAL GRADE

GENERAL COMMENTS

/100

---

PAGE 1

---

PAGE 2

---

PAGE 3

---